

The Time–Dependent Born–Oppenheimer Approximation, Crossings, Avoided Crossings, and Some Non-Adiabatic Transitions

George A. Hagedorn

Department of Mathematics, and
Center for Statistical Mechanics, Mathematical Physics
and Theoretical Chemistry

Virginia Tech

Blacksburg, Virginia 24061–0123

USA

hagedorn@math.vt.edu

Nice Quantum Chemistry Workshop

14 Nov. 2008

Outline

1. Semiclassical Wave Packets
2. The Time-Dependent Born-Oppenheimer Approximation
3. Exponentially Small Non-Adiabatic Scattering Transitions
4. Propagation Through Level Crossings and Avoided Crossings

Semiclassical Wave Packets

To state time-dependent results in their most explicit form, we need to discuss semiclassical wave packets $\phi_k(A, B, \hbar, a, \eta, x)$.

- These are generalizations of Harmonic oscillator states.
- They coincide with generalized squeezed states.
- In the molecular context, \hbar will be ϵ^2 .

$\{ \phi_k(A, B, \hbar, a, \eta, x) \}$ is an orthonormal basis of $L^2(\mathbb{R}^d)$ as k ranges over d -dimensional multi-indices.

- $a \in \mathbb{R}^d$ represents a classical position.
- $\eta \in \mathbb{R}^d$ represents a classical momentum.
- A and B are complex invertible $d \times d$ matrices that satisfy $A^t B - B^t A = 0$ and $A^* B + B^* A = 2I$.

The position uncertainty is determined by $\epsilon |A|$, and the momentum uncertainty is determined by $\epsilon |B|$.

$$\begin{aligned} \phi_0(A, B, \hbar, a, \eta, x) &= \pi^{-d/4} \hbar^{-d/4} (\det(A))^{-1/2} \\ &\times \exp \left\{ - (x - a) \cdot BA^{-1}(x - a)/(2\hbar) + i\eta \cdot (x - a)/\hbar \right\}. \end{aligned}$$



There are raising and lowering operators with the same algebraic properties as with the Harmonic oscillator.


Define the Fourier Transform


$$(\mathcal{F}_\hbar f)(\xi) = (2\pi\hbar)^{-d/2} \int_{\mathbb{R}^d} f(x) e^{-i\xi \cdot x/\hbar} dx.$$


Then

$$(\mathcal{F}_\hbar \phi_k(A, B, \hbar, a, \eta, \cdot))(\xi) = e^{-ia \cdot \eta/\hbar} \phi_k(B, A, \hbar, \eta, -a, \xi).$$









If $V(X)$ is smooth and bounded below, then

$$e^{iS(t)/\hbar} \sum_{|k| \leq K} c_k \phi_k(A(t), B(t), \hbar, a(t), \eta(t), X)$$

solves the time-dependent Schrödinger equation

$$i \hbar \frac{\partial \psi}{\partial t} = - \frac{\hbar^2}{2} \Delta_X \psi + V(X) \psi,$$

up to an $O(\hbar^{1/2})$ error.

Here

$$\dot{a}(t) = \eta(t),$$

$$\dot{\eta}(t) = - V^{(1)}(a(t)),$$

$$\dot{A}(t) = i B(t),$$

$$\dot{B}(t) = i V^{(2)}(a(t)) A(t),$$

$$\dot{S}(t) = \frac{\eta(t)^2}{2} - V(a(t)).$$

If $V(X)$ is quadratic, there is no error.

There are many generalizations of this result.

(Time dependent V 's. Higher order in $\hbar^{1/2}$. Approximations with $\exp(-C/\hbar)$ errors from optimal truncation.)

Faou, Gradinaru, and Lubich recently developed a numerical algorithm for solving semiclassical time-dependent Schrödinger equations that is based on these wave packets.

It scales very well as the space dimension and/or the approximation order are increased.

The Time-Dependent Born-Oppenheimer Approximation

Molecular Hamiltonians can be written as

$$H(\epsilon) = -\frac{\epsilon^4}{2} \Delta_X + h(X),$$

where the electron Hamiltonian $h(X)$ depends parametrically on the nuclear configuration X .

We wish to find approximate solutions to

$$i \epsilon^2 \frac{\partial \Psi}{\partial t} = H(\epsilon) \Psi.$$

Born–Oppenheimer Approximations treat the electrons and nuclei separately, while respecting the coupling between them.

STEP 1. For each configuration X of the nuclei, solve the electronic eigenvalue problem.

$$h(X) \Phi(X, x) = E(X) \Phi(X, x).$$

- The various different discrete eigenvalues $E(X)$ that depend continuously on X are called electron energy levels.

STEP 2. Use the semiclassical wave packets for the nuclei with an electron energy level $E(X)$ playing the role of the potential.

Hypotheses

- Assume the resolvent of $h(X)$ is smooth in X .
- Assume $E(X)$ is a non-degenerate level for all X , and let $\Phi(X)$ be an associated normalized eigenvector with phase chosen so $\langle \Phi(X, \cdot), \nabla_X \Phi(X, \cdot) \rangle_{\mathcal{H}_{\text{el}}} = 0$.
- Solve the semiclassical equations of motion

$$\dot{a}(t) = \eta(t),$$

$$\dot{\eta}(t) = -E^{(1)}(a(t)),$$

$$\dot{A}(t) = i B(t),$$

$$\dot{B}(t) = i E^{(2)}(a(t)) A(t),$$

$$\dot{S}(t) = \frac{\eta(t)^2}{2} - E(a(t)).$$

Theorem 1 The time-dependent Schrödinger equation has a solution of the form $\Psi_{N,\epsilon}(X, x, t) + error_{N,\epsilon}$,

where
$$\Psi_{N,\epsilon}(X, x, t) = \sum_{n=0}^N \psi_{n,\epsilon}(X, x, t) \epsilon^n$$

and
$$\|error_{N,\epsilon}\| \leq C_N \epsilon^{N+1}, \quad \text{for } t \in [0, T].$$

The zeroth order term in the expansion is

$$\begin{aligned} & \psi_{0,\epsilon}(X, x, t) \\ &= e^{iS(t)/\epsilon^2} \Phi(X, x) \sum_{|k| \leq K} c_k \phi_k(A(t), B(t), \epsilon^2, a(t), \eta(t), X), \end{aligned}$$

where the c_k and K are arbitrary.

Theorem 2 Under analyticity assumptions on $h(X)$, we can choose $N(\epsilon) = O(\epsilon^{-2})$, such that the Schrödinger equation has a solution of the form $\Psi_{N(\epsilon),\epsilon} + error_\epsilon$,

where $\|error_\epsilon\| \leq C \exp\left(-\frac{\Gamma}{\epsilon^2}\right)$, for $t \in [0, T]$.

Furthermore, given any $b > 0$, this $N(\epsilon)$ can be chosen so that there exist D and $\gamma > 0$, such that

$$\int_{|X-a(t)|>b} \|\Psi_{N(\epsilon),\epsilon}(X, x, t)\|_{\mathcal{H}_{el}}^2 dX \leq D \exp\left(-\frac{\gamma}{\epsilon^2}\right).$$

Remarks

- Semiclassical wave packet techniques handle the nuclear motion.
- An adiabatic expansion handles the electronic states.
- We use the Method of Multiple Scales to separate semiclassical terms from adiabatic terms in the perturbation calculations.
- The analog of Theorem 1 is proven for Coulomb potentials.
- Theorem 2 follows from Theorem 1 and the estimate
$$\|error_{N-1,\epsilon}\| \leq \alpha \beta^N N^{N/2} \epsilon^N$$
by a simple calculation.
- The form of the error estimate in Theorem 2 is optimal. Our approximation ignores tunnelling by the nuclei and non-adiabatic transitions by the electrons.

Remarks (continued)


- There are several other approaches. See, for example,
 - C. Fermanian–Kammerer, P. Gérard.
 - G. Panati, H. Spohn, S. Teufel.
 - C. Lasser, C. Fermanian–Kammerer.
 - A. Martinez, V. Sordoni.

Exponentially Small Non-Adiabatic Transitions

We only have leading order results for small ϵ
when nuclei have 1 degree of freedom
and the electron Hamiltonian is an analytic $n \times n$ matrix.

Example that illustrates rigorous results

$$h(x) = \frac{1}{2} \begin{pmatrix} 1 & \tanh(x) \\ \tanh(x) & -1 \end{pmatrix}$$



Scattering with large negative t asymptotics

$$e^{iS(t)/\epsilon^2} \phi_k(A(t), B, \epsilon^2, a(t), \eta, x) \Phi_{\text{up}}(x).$$

What should we expect?

- The nuclei behave like classical particles (at least for small k).
- The electrons should feel a time-dependent Hamiltonian

$$\tilde{h}(t) = \frac{1}{2} \begin{pmatrix} 1 & \tanh(a(t)) \\ \tanh(a(t)) & -1 \end{pmatrix},$$

and we should simply use the Landau–Zener formula to get the exponentially small transition probability.

- For $\eta = 1$, energy conservation predicts the momentum after the transition to be 1.9566.


This intuitive picture is wrong!

- The transition amplitude is larger than predicted.
- The momentum after the transition is larger than predicted.


Additional Surprises

- For incoming state ϕ_k , the nuclear wave function after the transition is not what one might naively expect.
 - The nuclear wavepacket after transition is a ϕ_0 .
 - The transition amplitude is asymptotically of order


$$\epsilon^{-k} \exp\left(-\alpha/\epsilon^2\right).$$




Position space plot at time $t = -10$ of the probability density for being on the upper energy level.




Momentum space plot at time $t = -10$ of the probability density for being on the upper energy level.




Position space probability density at time $t = 9$.
Lower level plot is multiplied by 3×10^8 .




Momentum space probability density at time $t = 9$.
Lower level plot is multiplied by 3×10^8 .




Position space probability density at time $t = -10$.



Momentum space probability density at time $t = -10$.



Position space probability density at time $t = 9$.
 Plot for the lower level has been multiplied by 10^7 .



Momentum space probability density at time $t = 9$.
 Plot for the lower level has been multiplied by 10^7 .

What's going on, and how do we analyze it?

- We expand $\Psi(x, t)$ in generalized eigenfunctions of $H(\epsilon)$.
- We then do a WKB approximation of the generalized eigenfunctions that is valid for complex x .
- We find that the Landau–Zener formula gives the correct transition amplitude for each generalized eigenfunction. This amplitude behaves roughly like $\exp\left(-\frac{C}{|p|\epsilon^2}\right)$, where p is the incoming momentum.
- So, higher momentum components of the wave function are drastically more likely to experience a transition. We get the correct result by using Landau–Zener for each p and then averaging.

Why do we always get a Gaussian?

- In the formulas, the extra shift in momentum occurs in the exponent.
- In momentum space ϕ_k all have the same exponential factor. The extra shift does not appear in the polynomial that multiplies the exponential.
- For small ϵ , to leading order, the polynomial factor looks like its largest order term near where the Gaussian is concentrated in momentum.
- $\left(\frac{p}{\epsilon}\right)^k \exp\left(-\frac{(p-\eta)^2}{\epsilon^2}\right)$ is approximately ϵ^{-k} times a Gaussian for $\eta \neq 0$.






FIG. 2. Potential-Energy Curves for H_2^- , H_2 , and H_2^+ . A large-scale pullout of this drawing appears at the front of this issue. In general $1s\sigma$ has been omitted from state designations.

Molecular Propagation through Level Crossings

- There are many different types of crossings because of symmetry considerations.
- Generic Minimal Multiplicity quantum mechanical crossings occur on codimension 1, 2, 3, and 5 submanifolds.
- For codimension 1 crossings we have rigorous results through order ϵ^1 .
- For higher codimension crossings we have rigorous results through order ϵ^0
- These results are obtained by using matched asymptotic expansions.
- There are more recent approaches:
C. Fermanian–Kammerer, P. Gérard, C. Lasser, G.–L. Panati, H. Spohn, S. Teufel, Y. Colin de Verdière.





Codimension 1 Example

$$h(X) = \begin{pmatrix} X \sin^2(X) & -X \sin(X) \cos(X) \\ -X \sin(X) \cos(X) & X \cos^2(X) \end{pmatrix}.$$

$$E_{\mathcal{A}}(X) = 0. \quad E_{\mathcal{B}}(X) = X.$$

$$\Phi_{\mathcal{A}}(X) = \begin{pmatrix} \cos(X) \\ \sin(X) \end{pmatrix}. \quad \Phi_{\mathcal{B}}(X) = \begin{pmatrix} -\sin(X) \\ \cos(X) \end{pmatrix}.$$

$$i \epsilon^2 \frac{\partial \Psi}{\partial t} = -\frac{\epsilon^4}{2} \frac{\partial^2 \Psi}{\partial X^2} + h(X) \Psi.$$

Incoming Outer Solution

$$e^{i\eta_0^2 t/(2\epsilon^2)} \phi_0(1 + it, 1, \epsilon^2, \eta_0 t, \eta_0, X) \Phi_{\mathcal{A}}(X) + O(\epsilon^2).$$

Inner Solution

Rewrite the Schrödinger equation in terms of rescaled variables $s = t/\epsilon$ and $Y = (X - \eta_0 t)/\epsilon$. Expand formally for small ϵ .



$$\begin{aligned}
 \Psi(X, t) &= e^{i\eta_0^2 t/(2\epsilon^2)} \phi_0(1 + it, 1, \epsilon^2, \eta_0 t, \eta_0, X) \Phi_{\mathcal{A}}(X) \\
 &\quad + \epsilon \eta_0 \left(\int_{-\infty}^{s+Y/\eta_0} e^{i\eta_0 r^2/2} dr \right) \\
 &\quad \times e^{iS^{\mathcal{B}}(t)/\epsilon^2} \phi_0(A^{\mathcal{B}}(t), B^{\mathcal{B}}, \epsilon^2, a^{\mathcal{B}}(t), \eta^{\mathcal{B}}(t), X) \Phi_{\mathcal{B}}(X) \\
 &\quad + O(\epsilon^{1+\delta})
 \end{aligned}$$

Here,

$$\begin{aligned}
 S^{\mathcal{B}}(t) &= \eta_0^2 t/2 - \eta_0 t^2 + t^3/3, & A^{\mathcal{B}}(t) &= 1 + (i + 1/\eta_0) t, \\
 B^{\mathcal{B}}(t) &= 1 - i/\eta_0, & a^{\mathcal{B}}(t) &= \eta_0 t - t^2/2, & \text{and } \eta^{\mathcal{B}}(t) &= \eta_0 - t.
 \end{aligned}$$

Outgoing Outer Solution

$$\begin{aligned}\Psi(X, t) &= e^{i\eta_0^2 t/(2\epsilon^2)} \phi_0(1 + it, 1, \epsilon^2, \eta_0 t, \eta_0, X) \Phi_{\mathcal{A}}(X) \\ &\quad + \epsilon (1 + i) \pi^{1/2} \eta_0^{1/2} \\ &\quad \quad \times e^{iS^{\mathcal{B}}(t)/\epsilon^2} \phi_0(A^{\mathcal{B}}(t), B^{\mathcal{B}}, \epsilon^2, a^{\mathcal{B}}(t), \eta^{\mathcal{B}}(t), X) \Phi_{\mathcal{B}}(X) \\ &\quad + O(\epsilon^{1+\delta})\end{aligned}$$



Codimension 2 Example

$$h(X) = \begin{pmatrix} X_1 & X_2 \\ X_2 & -X_1 \end{pmatrix}.$$

$$E_{\mathcal{A}}(X) = -(X_1^2 + X_2^2)^{1/2}, \quad E_{\mathcal{B}}(X) = (X_1^2 + X_2^2)^{1/2}.$$

$$\Phi_{\mathcal{A}}(X) = \begin{pmatrix} -\sin(\theta/2) \\ \cos(\theta/2) \end{pmatrix}, \quad \Phi_{\mathcal{B}}(X) = \begin{pmatrix} \cos(\theta/2) \\ \sin(\theta/2) \end{pmatrix}.$$

where $\theta = \tan^{-1}(X_2/X_1)$ with $-\frac{\pi}{2} \leq \theta \leq \frac{3\pi}{2}$.

$$i \epsilon^2 \frac{\partial \Psi}{\partial t} = -\frac{\epsilon^4}{2} \left(\frac{\partial^2 \Psi}{\partial X_1^2} + \frac{\partial^2 \Psi}{\partial X_2^2} \right) + h(X) \Psi.$$

Incoming Outer Solution

$$e^{i S^{\mathcal{A}}(t)/\epsilon^2} \phi_0(A^{\mathcal{A}}(t), B^{\mathcal{A}}(t), \epsilon^2, a^{\mathcal{A}}(t), \eta^{\mathcal{A}}(t), X) \Phi_{\mathcal{A}}(X) + O(\epsilon^2).$$

$$a^{\mathcal{A}}(t) = \begin{pmatrix} \eta_0 t - t^2/2 \\ 0 \end{pmatrix}, \quad \eta^{\mathcal{A}}(t) = \begin{pmatrix} \eta_0 - t \\ 0 \end{pmatrix},$$

$S^{\mathcal{A}}(t)$, $A^{\mathcal{A}}(t)$, $B^{\mathcal{A}}(t)$ fairly complicated, but $A^{\mathcal{A}}(0) = I$.

Inner Solution

Rescale: $s = t/\epsilon$, $Y_1 = (X_1 - \eta_0 t)/\epsilon$, $Y_2 = X_2/\epsilon$.

In the new variables, the leading order inner solution can be written explicitly in terms of parabolic cylinder functions of complex order and complex argument.

Outgoing Outer Solution

$$e^{i S^{\mathcal{A}}(t)/\epsilon^2} \sum_k c_k \phi_k(A^{\mathcal{A}}(t), B^{\mathcal{A}}(t), \epsilon^2, a^{\mathcal{A}}(t), \eta^{\mathcal{A}}(t), X) \Phi_{\mathcal{A}}(X) \\ + e^{i S^{\mathcal{B}}(t)/\epsilon^2} \phi_0(A^{\mathcal{B}}(t), B^{\mathcal{B}}(t), \epsilon^2, a^{\mathcal{B}}(t), \eta^{\mathcal{B}}(t), X) \Phi_{\mathcal{B}}(X) + O(\epsilon^\delta).$$

$$a^{\mathcal{A}}(t) = \begin{pmatrix} \eta_0 t + t^2/2 \\ 0 \end{pmatrix}, \quad \eta^{\mathcal{A}}(t) = \begin{pmatrix} \eta_0 + t \\ 0 \end{pmatrix},$$

$$a^{\mathcal{B}}(t) = \begin{pmatrix} \eta_0 t - t^2/2 \\ 0 \end{pmatrix}, \quad \eta^{\mathcal{B}}(t) = \begin{pmatrix} \eta_0 - t \\ 0 \end{pmatrix},$$


Probability that the system moves to the upper surface is

$$(1 + \pi/\eta_0)^{1/2} + O(\epsilon^\delta).$$


lower electronic level

upper electronic level


$t = -10\epsilon$






$t = 0$



$t = 10\epsilon$



Avoided Crossings



Results for Codimension 2 Avoided Crossings

