

Towards quantitative modelling of biological and complex systems using polarizable force fields.

Jean-Philip Piquemal

Laboratoire de Chimie Théorique, UMR 7616 CNRS, UPMC, Paris 6.

Email: jpp@lct.jussieu.fr; Web: www.lct.jussieu.fr/pagesperso/jpp

In this talk, we will discuss some new results and ongoing developments concerning several polarizable force field methods such as the SIBFA , AMOEBA and GEM approaches. Their capability to compute small intermolecular energy differences as well as their applicability to spectroscopic, molecular dynamics and hybrid QM/MM simulations studies will be adressed.

References

- 1) N. Gresh, G. A. Cisneros, T. A. Darden and J-P Piquemal, J. Chem. Theory. Comput., **2007**, 3, 1960.
- 2) Design of next generation polarizable force fields from ab initio computations: beyond point charges. G. A. Cisneros, T. A. Darden, N. Gresh, P. Reinhardt, O. Parisel, J. Pilmé and J-P Piquemal in Multi-scale Quantum Models for Biocatalysis: Modern Techniques and Applications, for the Book Series: Challenges and Advances in Computational Chemistry and Physics , ed. D. M. York and T.-S. Lee, **2009**, 137-172, Springer Verlag,
- 3) J-P. Piquemal, G. A. Cisneros, P. Reinhardt, N. Gresh and T. A. Darden, J. Chem. Phys., **2006**, 124, 104101.
- 4) J-P. Piquemal, L. Perera, G. A. Cisneros, P. Ren, L. G. Pedersen and T. A. Darden, J. Chem. Phys., **2006**, 125, 054511.