

Smooth branch of travelling waves for the Gross-Pitaevskii equation in \mathbb{R}^2 for small speed

David Chiron and Eliot Pacherie

Université Côte d'Azur, CNRS, LJAD, France

Abstract

We construct a smooth branch of travelling wave solutions for the 2 dimensional Gross-Pitaevskii equations for small speed. These travelling waves exhibit two vortices far away from each other. We also compute the leading order term of the derivatives with respect to the speed. We construct these solutions by an implicit function type argument.

Mathematics Subject Classification (2010): 35C07; 35B20; 35Q40; 35Q56.

1 Introduction and statement of the result

We consider the Gross-Pitaevskii equation

$$0 = (\text{GP})(\mathbf{u}) := i\partial_t \mathbf{u} + \Delta \mathbf{u} - (|\mathbf{u}|^2 - 1)\mathbf{u}$$

in dimension 2 for $\mathbf{u} : \mathbb{R}_t \times \mathbb{R}_x^2 \mapsto \mathbb{C}$. The Gross-Pitaevskii equation is a physical model for Bose-Einstein condensates [10], [18], and is associated with the Ginzburg-Landau energy

$$E(v) := \frac{1}{2} \int_{\mathbb{R}^2} |\nabla v|^2 + \frac{1}{4} \int_{\mathbb{R}^2} (1 - |v|^2)^2.$$

The condition at infinity for (GP) will be

$$|\mathbf{u}| \rightarrow 1 \quad \text{as} \quad |x| \rightarrow +\infty.$$

We look for travelling wave solutions of (GP):

$$\mathbf{u}(t, x) = v(x_1, x_2 + ct)$$

where $x = (x_1, x_2)$ and $c > 0$ is the speed of the travelling wave, which moves along the direction $-\vec{e}_2$. The equation on v is

$$0 = (\text{TW}_c)(v) := -ic\partial_{x_2} v - \Delta v - (1 - |v|^2)v.$$

We want to construct travelling waves for small speed that look like the product of two well-separated vortices. Vortices are stationary solutions of (GP) of degrees $n \in \mathbb{Z}^*$ (see [5] and [14]):

$$V_n(x) = \rho_n(r)e^{in\theta},$$

where $x = re^{i\theta}$, solving

$$\begin{cases} \Delta V_n - (|V_n|^2 - 1)V_n = 0 \\ |V_n| \rightarrow 1 \quad \text{as} \quad |x| \rightarrow \infty. \end{cases}$$

A vortex alone in the plane is a stationary solution of (GP), and vortices might interact when there are several of them. It is expected that if they are far away from each other, their dynamic is governed, at least at first order, by the point vortex system (see [2] and references therein). In particular, a vortex V_1 of degree 1 with a vortex V_{-1} of degree -1 should move at constant speed in the direction orthogonal to the line that connects their centers.

The main result of this paper is the construction of a branch of solution by perturbation of the product of two vortices at any small speed $c > 0$, and the fact that this branch of solution is C^1 with respect to the speed.

Theorem 1.1 *There exists $c_0 > 0$ a small constant such that, for any $0 < c \leq c_0$, there exists a solution of (TW_c) of the form*

$$Q_c = V_1(\cdot - d_c \vec{e}_1) V_{-1}(\cdot + d_c \vec{e}_1) + \Gamma_{c,d_c},$$

where $d_c = \frac{1+o_{c \rightarrow 0}(1)}{c}$ is a continuous function of c . This solution has finite energy ($E(Q_c) < +\infty$) and $Q_c \rightarrow 1$ when $|x| \rightarrow +\infty$.

Furthermore, for all $+\infty \geq p > 2$, there exists $c_0(p) > 0$ such that if $c < c_0(p)$, for the norm

$$\|h\|_{X_p} := \|h\|_{L^p(\mathbb{R}^2)} + \|\nabla h\|_{L^{p-1}(\mathbb{R}^2)}$$

and the space $X_p := \{f \in L^p(\mathbb{R}^2), \nabla f \in L^{p-1}(\mathbb{R}^2)\}$, one has

$$\|\Gamma_{c,d_c}\|_{X_p} = o_{c \rightarrow 0}(1).$$

In addition,

$$c \mapsto Q_c - 1 \in C^1(]0, c_0(p)[, X_p),$$

with the estimate (for $\nu(c) = \frac{1+o_{c \rightarrow 0}(1)}{c^2}$)

$$\|\partial_c Q_c + \nu(c) \partial_d (V_1(\cdot - d \vec{e}_1) V_{-1}(\cdot + d \vec{e}_1))|_{d=d_c}\|_p = o_{c \rightarrow 0} \left(\frac{1}{c^2} \right).$$

Existence of travelling waves solutions for this equation with finite energy has already been proven for small speeds in [3] (see also [4] and [6] for results in dimension 2 and 3). Moreover, the decay at infinity conjectured in [15] has been established in [13]. Here, we use an implicit function argument to construct the solution, using techniques developed in [8] or [16] for instance, displaying a clear understanding of the shape of the solution (see Lemma 3.8 for instance). We show in addition that the constructed branch is C^1 , which is, to the best of our knowledge, the first result of this kind in dimension larger than one.

In the Gross-Pitaevskii equation, vortices play the role of solitons (as we can see in NLS or other such equations). In particular here we show that two vortices interact at long range since the speed c is of order $\frac{1}{d_c}$, the half distance between the vortices. This is due to the slow decay of the vortex: ∇V_1 is of order $\frac{1}{r}$ at infinity due to the phase.

The formal method for this kind of construction is well known. It has been done rigorously in a bounded domain for the Ginzburg Landau equation with no speed ([8]). One of the difficulties here is to find the right functional setting to construct the C^1 branch, in particular with regards to the transport term $ic \partial_{x_2} v$. On the contrary of what is claim in [16], the transport term can not be treated perturbatively. This is why we use another functional setting than [16] or [17] (see Remark 2.11 for more details)

In this paper, we start by doing the construction of the solution to fill these gaps in the case of two vortices in (GP). This construction is also a good introduction to the proof of the differentiability of the branch, which uses many of the same ideas, but with a more technical setting.

We start by reducing the problem to a one dimensional one in section 2. The construction of the travelling wave Q_c is completed in section 3. Furthermore, in subsection 3.2, we show that Q_c has finite energy and we compute some estimates particular to the branch of solutions. Finally section 4 is devoted to the proof of the differentiability of the branch.

We use the scalar product for $f, g \in L^2(\mathbb{R}^2)$,

$$\langle f, g \rangle := \Re \int_{\mathbb{R}^2} f \bar{g},$$

For $X = (X_1, X_2), Y = (Y_1, Y_2) \in \mathbb{C}^2$, we define

$$X.Y := X_1 Y_1 + X_2 Y_2,$$

which is the scalar product if $X, Y \in \mathbb{R}^2$. We use the notation $B(x, r)$ to define the closed ball in \mathbb{R}^2 of center $x \in \mathbb{R}^2$ and radius $r > 0$ for the Euclidean norm. In the estimates, a constant $K > 0$ is a universal constant independent of any parameter of the problem.

Technical lemmas given without proofs here are detailed in the arXiv version (1911.03433). It also contains additional precisions for some estimates.

Acknowledgments . The authors would like to thank Pierre Raphaël for helpful discussions. E.P. is supported by the ERC-2014-CoG 646650 SingWave.

2 Lyapunov-Schmidt reduction

The proof of Theorem 1.1 follows closely the construction done in [8] or [16]. The main idea is to use perturbation methods on an approximate solution.

In subsection 2.1 we define this approximate solution V which consists in two vortices at distance $2d$ from each other. We then look for a solution of (TW_c) as a perturbation of V , with an additive perturbation close to the vortices and a multiplicative one far from them. This is computed in subsection 2.2. We define suitable spaces in subsection 2.3 that we will use to invert the linear part and use a contraction argument. We ask for an orthogonality on the perturbation, and the norms are a little better but more technical than the ones in Theorem 1.1. In particular Γ_{c,d_c} in Theorem 1.1 verifies better estimates which are discussed for instance in Corollary 2.25 and in Lemma 3.8. We invert the linearized operator in Proposition 2.17 and show that the perturbation is a fixed point of a contracting functional in Proposition 2.21. The orthogonality condition create a Lagrangian multiplier (see subsection 2.6), which left us with a problem in one dimension. This multiplier will be cancelled for a good choice of the parameter d in section 3.

2.1 Estimates on vortices

From [14], we can find stationary solution of (GP):

$$V_n(x) = \rho_n(r)e^{in\theta}$$

where $x = re^{i\theta}$, $n \in \mathbb{Z}^*$, solving

$$\begin{cases} \Delta V_n - (|V_n|^2 - 1)V_n = 0 \\ |V_n| \rightarrow 1 \quad \text{as } |x| \rightarrow \infty. \end{cases}$$

These solutions are well understood and, in particular, we have some estimates (see [14] for instance) that we will use. We also know the kernel of the linearized operator around $V_{\pm 1}$ ([7]), which we will need for inverting the linearized operator around the approximate solution V defined using these vortices

$$V(x) := V_1(x - d\vec{e}_1)V_{-1}(x + d\vec{e}_1)$$

where $d > 0$, $x = (x_1, x_2)$. The function V is the product of two vortices with opposite degrees at a distance $2d$ from each other. One vortex alone in \mathbb{R}^2 is a stationary solution, and it is expected that two vortices interact and translate at a constant speed of order $c \simeq \frac{1}{d}$, see [2]. Hence for the two parameters of this problem $c, d > 0$, we let them be free from each other, but with the condition c is of order $1/d$ by imposing that $\frac{1}{2c} < d < \frac{2}{c}$.

We will study in particular areas near the center of each vortices. We will use coordinates adapted to this problem:

$$\begin{aligned} x &= (x_1, x_2) = re^{i\theta}, \\ y &= (y_1, y_2) := x - d\vec{e}_1 = r_1e^{i\theta_1}, \\ z &= (z_1, z_2) := y + 2d\vec{e}_1 = x + d\vec{e}_1 = r_{-1}e^{i\theta_{-1}}, \\ \tilde{r} &:= \min(r_1, r_{-1}). \end{aligned} \tag{2.1}$$

Using y coordinate mean that we are centered around V_1 , and z coordinate for around V_{-1} . Note that we have

$$V(x) = V_1(y)V_{-1}(z)$$

using these notations. If it is not precised, V will be taken in x , V_1 in y and V_{-1} in z . If we compute (TW_c) for V , i.e. $-ic\partial_{x_2}V - \Delta V - (1 - |V|^2)V$, we get

$$(\text{TW}_c)(V) = E - ic\partial_{x_2}V,$$

where we defined

$$E := -\Delta V - (1 - |V|^2)V.$$

We have $V = V_1V_{-1}$ and, by using $-\Delta V_\varepsilon = (1 - |V_\varepsilon|^2)V_\varepsilon$ for $\varepsilon = \pm 1$, we compute

$$E = -2\nabla V_1 \cdot \nabla V_{-1} + V_1V_{-1}(1 - |V_1|^2 + 1 - |V_{-1}|^2 - 1 + |V_1V_{-1}|^2).$$

Hence

$$E = -2\nabla V_1 \cdot \nabla V_{-1} + (1 - |V_1|^2)(1 - |V_{-1}|^2)V_1V_{-1}. \quad (2.2)$$

The rest of this subsection is devoted to estimates on $V, E, \partial_d V$ and $ic\partial_{x_2}V$ using estimates on V_1 and V_{-1} . Let us start with the properties on $V_{\pm 1}$ we need.

Lemma 2.1 ([14]) $V_1(x) = \rho_1(r)e^{i\theta}$ verifies $V_1(0) = 0$, and there exists a constant $\kappa > 0$ such that, for all $r > 0$, $0 < \rho_1(r) < 1$, $\rho_1'(r) > 0$, and

$$\begin{aligned} \rho_1(r) &\sim_{r \rightarrow 0} \kappa r, \\ \rho_1'(r) &= O_{r \rightarrow \infty} \left(\frac{1}{r^3} \right), \\ \rho_1''(r) &= o_{r \rightarrow \infty} \left(\frac{1}{r^3} \right), \\ 1 - |V_1(x)| &= \frac{1}{2r^2} + O_{r \rightarrow \infty} \left(\frac{1}{r^3} \right), \\ \nabla V_1(x) &= iV_1(x) \frac{x^\perp}{r^2} + O_{r \rightarrow \infty} \left(\frac{1}{r^3} \right) \end{aligned}$$

where $x^\perp = (-x_2, x_1)$, $x = re^{i\theta}$. Furthermore we have similar properties for V_{-1} since

$$V_{-1}(x) = \overline{V_1(\overline{x})}.$$

We will use the O notation for convergence independent of any other quantity. Now let us write all the derivatives of a vortex in polar coordinate, which will be useful all along the proof of the results.

Lemma 2.2 We define $u := \frac{\rho_1'(r_1)}{\rho_1(r_1)}$. Then,

$$\begin{aligned} \partial_{x_1} V_1(y) &= \left(\cos(\theta_1)u - \frac{i}{r_1} \sin(\theta_1) \right) V_1, \\ \partial_{x_2} V_1(y) &= \left(\sin(\theta_1)u + \frac{i}{r_1} \cos(\theta_1) \right) V_1, \\ \partial_{x_1 x_1} V_1(y) &= \left(\cos^2(\theta_1)(u^2 + u') + \sin^2(\theta_1) \left(\frac{u}{r_1} - \frac{1}{r_1^2} \right) + 2i \sin(\theta_1) \cos(\theta_1) \left(\frac{1}{r_1^2} - \frac{u}{r_1} \right) \right) V_1, \\ \partial_{x_1 x_2} V_1(y) &= \left(\sin(\theta_1) \cos(\theta_1) \left(u^2 + u' + \frac{1}{r_1^2} - \frac{u}{r_1} \right) - i \cos(2\theta_1) \left(\frac{1}{r_1^2} - \frac{u}{r_1} \right) \right) V_1. \end{aligned}$$

We obtain the derivatives of V_{-1} by changing $i \rightarrow -i$, $y \rightarrow z$, $\theta_1 \rightarrow \theta_{-1}$, $r_1 \rightarrow r_{-1}$ and $V_1 \rightarrow V_{-1}$. We remark in particular that the first derivatives are of first order $\frac{1}{r_1}$ and the second derivatives are of first order $\frac{1}{r_1^2}$ for large values of r_1 . From [14], we can check that, more generally, we have

$$|D^{(n)}V_1(y)| \leq \frac{K(n)}{(1+r_1)^n}. \quad (2.3)$$

Now, we compute some basic estimates on V .

Lemma 2.3 *There exists a universal constant $K > 0$ and a constant $K(d) > 0$ depending only on $d > 1$ such that*

$$\begin{aligned} |1 - V|^2 &\leq \frac{K(d)}{(1+r)^2}, \\ 0 \leq 1 - |V|^2 &\leq \frac{K}{(1+\tilde{r})^2}, \\ |\nabla(|V|)| &\leq \frac{K}{(1+\tilde{r})^3}, \end{aligned}$$

and we have

$$|\nabla V| \leq \frac{K}{(1+\tilde{r})},$$

as well as

$$|\nabla V| \leq \frac{Kd}{(1+\tilde{r})^2},$$

where $\tilde{r} = \min(r_1, r_{-1})$. Furthermore,

$$|\nabla^2 V| \leq \frac{K}{(1+\tilde{r})^2}$$

and

$$|\nabla^2 V| \leq \frac{Kd}{(1+\tilde{r})^3}.$$

We also look at the convergence of some quantities when we are near the center of V_1 and $d \rightarrow \infty$. When we are close to the center of V_1 and d goes to infinity, we expect that the second vortex has no influence.

Lemma 2.4 *As $d \rightarrow \infty$, we have, locally uniformly in \mathbb{R}^2 ,*

$$\begin{aligned} V(\cdot + d\vec{e}_1) &= V_1(\cdot)V_{-1}(\cdot + 2d\vec{e}_1) \rightarrow V_1(\cdot), \\ E(\cdot + d\vec{e}_1) &\rightarrow 0 \end{aligned}$$

and

$$\partial_d V(\cdot + d\vec{e}_1) \rightarrow -\partial_{x_1} V_1(\cdot).$$

We can do a precise computation on the term $ic\partial_{x_2}V$, which appears in $(\text{TW}_c)(V)$.

Lemma 2.5 *There exists a universal constant $C > 0$ (independent of d) such that if $r_1, r_{-1} \geq 1$,*

$$\left| i \frac{\partial_{x_2} V}{V} - 2d \frac{x_1^2 - d^2 - x_2^2}{r_1^2 r_{-1}^2} \right| \leq C \left(\frac{1}{r_1^3} + \frac{1}{r_{-1}^3} \right).$$

Remark that this shows that the first order term of $i \frac{\partial_{x_2} V}{V}$ is real-valued and the dependence on d of this term is explicit.

Finally, we infer estimates on $\partial_d V = \partial_d(V_1(x - d\vec{e}_1)V_{-1}(x + d\vec{e}_1)) = -\partial_{x_1} V_1 V_{-1} + \partial_{x_1} V_{-1} V_1$.

Lemma 2.6 *There exists a constant $K > 0$ such that*

$$\begin{aligned} |\partial_d V| &\leq \frac{K}{(1+\tilde{r})}, \\ |\nabla \partial_d V| &\leq \frac{K}{(1+\tilde{r})^2} \end{aligned}$$

and

$$|\Re(\bar{V} \partial_d V)| \leq \frac{K}{(1+\tilde{r})^3}.$$

Furthermore,

$$|\partial_d^2 V| \leq \frac{K}{(1 + \tilde{r})^2}$$

and

$$|\partial_d^2 \nabla V| \leq \frac{K}{(1 + \tilde{r})^3}.$$

2.2 Setup of the proof

In the same way as in [8] (see also [16]), we will look at a solution of (TW_c) as a perturbation of V of the form

$$v := \eta V(1 + \Psi) + (1 - \eta)Ve^\Psi$$

where $\eta(x) = \tilde{\eta}(r_1) + \tilde{\eta}(r_{-1})$ and $\tilde{\eta}$ is a C^∞ positive cutoff with $\tilde{\eta}(r) = 1$ if $r \leq 1$ and 0 if $r \geq 2$. The perturbation is Ψ and we will also use

$$\Phi := V\Psi.$$

We use such a perturbation because we want it to be additive (in Φ) near the center of the vortices (where $v = V + \Phi$), and multiplicative (in Ψ) far from them (where $v = Ve^\Psi$). We shall require Φ to be bounded (and small) near the vortices. The problem becomes an equation on Ψ , with the following Lemma 2.7, we shall write

$$\eta L(\Phi) + (1 - \eta)VL'(\Psi) + F(\Psi) = 0$$

where L and L' are linear. The main part of the proof of the construction consists of inverting the linearized operator $\eta L(\Phi) + (1 - \eta)VL'(\Psi)$ in suitable spaces, and then use a contraction argument by showing that F is small and conclude on the existence of a solution Ψ by a fixed point theorem.

Lemma 2.7 *The function $v = \eta V(1 + \Psi) + (1 - \eta)Ve^\Psi$ is solution of (TW_c) if and only if*

$$\eta L(\Phi) + (1 - \eta)VL'(\Psi) + F(\Psi) = 0,$$

where $\Phi = V\Psi$,

$$\begin{aligned} L'(\Psi) &:= -\Delta\Psi - 2\frac{\nabla V}{V} \cdot \nabla\Psi + 2|V|^2\Re(\Psi) - ic\partial_{x_2}\Psi, \\ L(\Phi) &:= -\Delta\Phi - (1 - |V|^2)\Phi + 2\Re(\bar{V}\Phi)V - ic\partial_{x_2}\Phi, \\ F(\Psi) &:= E - ic\partial_{x_2}V + V(1 - \eta)(-\nabla\Psi \cdot \nabla\Psi + |V|^2S(\Psi)) + R(\Psi), \end{aligned}$$

with

$$\begin{aligned} E &= -\Delta V - (1 - |V|^2)V, \\ S(\Psi) &:= e^{2\Re(\Psi)} - 1 - 2\Re(\Psi) \end{aligned}$$

and $R(\Psi)$ is a sum of terms at least quadratic in Ψ or Φ localized in the area where $\eta \neq 0$. Furthermore, there exists $C, C_0 > 0$ such that the estimate

$$|R(\Psi)| + |\nabla R(\Psi)| \leq C\|\Phi\|_{C^2(\{\tilde{r} \leq 2\})}^2$$

holds if $\|\Phi\|_{C^2(\mathbb{R}^2)} \leq C_0$ (a constant independent of c), where $\tilde{r} = \min(|x - d\vec{e}_1|, |x + d\vec{e}_1|)$ for $x \in \mathbb{R}^2$. Additionally, $L(\Phi)$ and $L'(\Psi)$ are related by

$$L(\Phi) = (E - ic\partial_{x_2}V)\Psi + VL'(\Psi).$$

The main reason for such a perturbation ansatz is because $V(d\vec{e}_1) = V(-d\vec{e}_1) = 0$, so we can not divide by V as done in L' for instance when we look near the vortices, therefore an additive perturbation is more suitable. But far from the vortices, the perturbation is easier to compute when written multiplicatively with a factorisation by V . Remark also that this allows us to take Ψ to explode at $d\vec{e}_1$ and $-d\vec{e}_1$ as long as $\Phi = V\Psi$ does not. This is needed for the norm we use in subsection 2.3.

As we look for Φ small (it is a perturbation), the conditions $\|\Phi\|_{C^2(\mathbb{R}^2)} \leq C_0$ will always be true. We need them because some of the error terms have an exponential contribution in Ψ , and not only quadratic. We recall that, with our notations, $\nabla\Psi \cdot \nabla\Psi$ is complex-valued.

Remark that the quantity F contains only nonlinear terms and the source term, which is $E - ic\partial_{x_2}V$. Furthermore, contrary to the work [16], the transport term is in the linearized operator, and not considered as an error term in F .

2.3 Setup of the norms

For a given $\sigma \in \mathbb{R}$, we define, similarly as in [8] and [16], for $\Psi = \Psi_1 + i\Psi_2$ and $h = h_1 + ih_2$, the norms

$$\begin{aligned} \|\Psi\|_{*,\sigma,d} &:= \|V\Psi\|_{C^2(\{\tilde{r} \leq 3\})} \\ &+ \|\tilde{r}^{1+\sigma}\Psi_1\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla\Psi_1\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla^2\Psi_1\|_{L^\infty(\{\tilde{r} \geq 2\})} \\ &+ \|\tilde{r}^\sigma\Psi_2\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{1+\sigma}\nabla\Psi_2\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla^2\Psi_2\|_{L^\infty(\{\tilde{r} \geq 2\})}, \\ \|h\|_{**,\sigma,d} &:= \|Vh\|_{C^1(\{\tilde{r} \leq 3\})} \\ &+ \|\tilde{r}^{1+\sigma}h_1\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla h_1\|_{L^\infty(\{\tilde{r} \geq 2\})} \\ &+ \|\tilde{r}^{2+\sigma}h_2\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla h_2\|_{L^\infty(\{\tilde{r} \geq 2\})}, \end{aligned}$$

where $\tilde{r} = \min(r_1, r_{-1})$ (which depends on d). These are the spaces we shall use for the inversion of the linear operator for suitable values of σ .

This norm is not the “natural” energy norm that we could expect, for instance:

$$\|\Phi\|_{H_V}^2 := \int_{\mathbb{R}^2} |\nabla\Phi|^2 + (1 - |V|^2)|\Phi|^2 + \Re(\bar{V}\Phi)^2.$$

In particular, we require different conditions on the decay at infinity (with, in a way, less decay). As a consequence, the decay we have in Theorem 1.1 is not optimal (see [13]). This decay will be recovered later on by showing that the solution has finite energy. The main advantage of the norms $\|\cdot\|_{*,\sigma,d}$ and $\|\cdot\|_{**,\sigma,d}$ is that they will allow us to have uniform estimates on the error, without constants depending on c or d .

We are looking for a solution Ψ on a space of symmetric functions: we suppose that

$$\forall x = (x_1, x_2) \in \mathbb{R}^2, \Psi(x_1, x_2) = \overline{\Psi(x_1, -x_2)} = \Psi(-x_1, x_2)$$

because V and the equation has the same symmetries. With only those symmetries we will not be able to invert the linearized operator because it has a kernel, we also need an orthogonal condition. We define

$$Z_d(x) := \partial_d V(x)(\tilde{\eta}(4r_1) + \tilde{\eta}(4r_{-1})),$$

where $\tilde{\eta}$ is the same function as the one used for v : it is a C^∞ non negative smooth cutoff with $\tilde{\eta}(r) = 1$ if $r \leq 1$ and 0 if $r \geq 2$. In particular $Z_d(x) = 0$ if $\tilde{r} \geq 1/2$, which will make some computations easier. The other interest of the cutoff function is that without it

$$\partial_d V(x) = -\partial_{x_1} V_1 V_{-1} + \partial_{x_1} V_{-1} V_1$$

is not integrable in all \mathbb{R}^2 . We define the Banach spaces we shall use for inverting the linear part:

$$\begin{aligned} \mathcal{E}_{*,\sigma,d} &:= \\ \{\Phi = V\Psi \in C^2(\mathbb{R}^2, \mathbb{C}), \|\Psi\|_{*,\sigma,d} < +\infty; \langle \Phi, Z_d \rangle = 0; \forall x \in \mathbb{R}^2, \Psi(x_1, x_2) = \overline{\Psi(x_1, -x_2)} = \Psi(-x_1, x_2)\}, \\ \mathcal{E}_{**,\sigma',d} &:= \{Vh \in C^1(\mathbb{R}^2, \mathbb{C}), \|h\|_{**,\sigma',d} < +\infty\} \end{aligned}$$

for $\sigma, \sigma' \in \mathbb{R}$. We shall omit the subscript d in the construction and use only $\mathcal{E}_{*,\sigma}$, $\mathcal{E}_{**,\sigma'}$. Remark that $\mathcal{E}_{*,\sigma}$ contains an orthogonality condition as well as the symmetries.

Our first goal is to invert the linearized operator. This is a difficult part, which requires a lot of computations and critical elliptic estimates. The next subsection is devoted to the proof of the elliptic tools use in the proof of the inversion. In particular, our paper diverges here from [16] (see Remark 2.11 thereafter).

2.4 Some elliptic estimates

In this subsection, we provide some tools for elliptic estimate adapted to L^∞ norms.

2.4.1 Weighted L^∞ estimates on a Laplacian problem

Lemma 2.8 For $d \geq 5$, $0 < \alpha < 1$, there exists a constant $K(\sigma) > 0$ such that, for $f \in C^0(\mathbb{R}^2, \mathbb{C})$ such that

$$\forall (x_1, x_2) \in \mathbb{R}^2, \quad f(x_1, x_2) = -f(x_1, -x_2)$$

and with

$$\varepsilon_{f,\alpha} := \|f(x)(1 + \tilde{r})^{2+\alpha}\|_{L^\infty(\mathbb{R}^2)} < +\infty,$$

there exists a unique $C^1(\mathbb{R}^2)$ function ζ such that

$$\Delta \zeta = f$$

in the distribution sense,

$$\forall (x_1, x_2) \in \mathbb{R}^2, \quad \zeta(x_1, x_2) = -\zeta(x_1, -x_2)$$

and ζ satisfies the following two estimates:

$$\forall x \in \mathbb{R}^2, |\zeta(x)| \leq \frac{K(\sigma)\varepsilon_{f,\alpha}}{(1 + \tilde{r})^\alpha}$$

and

$$\forall x \in \mathbb{R}^2, |\nabla \zeta(x)| \leq \frac{K(\sigma)\varepsilon_{f,\alpha}}{(1 + \tilde{r})^{1+\alpha}}.$$

Remark here that for a given function f , if it satisfies two inequalities with different values of $(\varepsilon_{f,\alpha}, \alpha)$, then the associated function ζ satisfies the estimates with both sets of values by uniqueness. Furthermore, with only the hypothesis $f \in C^0(\mathbb{R}^2)$, we do not have $\zeta \in C_{\text{loc}}^2(\mathbb{R}^2)$ a priori.

2.4.2 Fundamental solution for $-\Delta + 2$

We will use the fundamental solution of $-\Delta + 2$. It can be deduce from the fundamental solution of $-\Delta + 1$, which has the following properties.

Lemma 2.9 ([1]) The fundamental solution of $-\Delta + 1$ in \mathbb{R}^2 is $\frac{1}{2\pi}K_0(|\cdot|)$, where K_0 is the modified Bessel function of second kind. It satisfies $K_0 \in C^\infty(\mathbb{R}^{+*})$ and

$$K_0(r) \sim_{r \rightarrow \infty} \sqrt{\frac{\pi}{2r}} e^{-r},$$

$$K_0(r) \sim_{r \rightarrow 0} -\ln(r),$$

$$K_0'(r) \sim_{r \rightarrow \infty} -\sqrt{\frac{\pi}{2r}} e^{-r},$$

$$K_0'(r) \sim_{r \rightarrow 0} \frac{-1}{r},$$

$$\forall r > 0, K_0(r) > 0, K_0'(r) < 0 \quad \text{and} \quad K_0''(r) > 0.$$

Proof The first three equivalents are respectively equations 9.7.2, 9.6.8 and 9.7.4 of [1]. The fourth one can be deduced from equations 9.6.27 and 9.6.9 of [1]. For $\nu \in \mathbb{N}$, K_ν is $C^\infty(\mathbb{R}, \mathbb{R})$ since it solves 9.6.1 of [1] and from the end of 9.6 of [1], we have that K_ν has no zeros. In particular with the asymptotics of 9.6.8, this implies that $K_\nu(r) > 0$. Furthermore, from 9.6.27 of [1], we have $K_0' = -K_1 < 0$ and $K_0'' = -K_1' = \frac{K_0 + K_2}{2} > 0$. \square

We give an elliptic estimate that will be used in the proof of Proposition 2.17.

Lemma 2.10 For any $\alpha > 0$, there exists a constant $C(\alpha) > 0$ such that, for any $d > 1$, if two real-valued functions $\Psi \in H^1(\mathbb{R}^2), h \in C^0(\mathbb{R}^2)$ satisfy in the distribution sense

$$(-\Delta + 2)\Psi = h,$$

and

$$\|(1 + \tilde{r})^\alpha h\|_{L^\infty(\mathbb{R}^2)} < +\infty,$$

then $\Psi \in C^1(\mathbb{R}^2)$ with

$$|\Psi| + |\nabla\Psi| \leq \frac{C(\alpha)\|(1 + \tilde{r})^\alpha h\|_{L^\infty(\mathbb{R}^2)}}{(1 + \tilde{r})^\alpha}.$$

Remark 2.11 Lemma 2.10 is different from the equivalent one of [16] for the gradient, which is equation (5.21) there. They claim that:

for any $0 < \sigma < 1$, there exists $C > 0$ such that, if two real-valued functions $\Psi \in C^1(\mathbb{R}^2), h \in C^0(\mathbb{R}^2)$ satisfy

$$(-\Delta + 2)\Psi = h$$

in the distribution sense, and

$$\|\Psi(1 + \tilde{r})^{1+\sigma}\|_{L^\infty(\mathbb{R}^2)} + \|\nabla\Psi(1 + \tilde{r})^{2+\sigma}\|_{L^\infty(\mathbb{R}^2)} + \|(1 + \tilde{r})^{1+\sigma}h\|_{L^\infty(\mathbb{R}^2)} < +\infty,$$

then

$$|\Psi| \leq \frac{C\|(1 + \tilde{r})^{1+\sigma}h\|_{L^\infty(\mathbb{R}^2)}}{(1 + \tilde{r})^{1+\sigma}}$$

and

$$|\nabla\Psi| \leq \frac{C\|(1 + \tilde{r})^{1+\sigma}h\|_{L^\infty(\mathbb{R}^2)}}{(1 + \tilde{r})^{2+\sigma}}.$$

The main difference they claim would be a stronger decay for the gradient. However, such a result can not hold, because of the following counterexample:

$$\Psi_\varepsilon(x) = \begin{cases} 0 & \text{if } |x| \leq 1/\varepsilon \\ \frac{\sin^2(r)}{(1+r)^{2+\sigma}} & \text{if } |x| \geq 1/\varepsilon. \end{cases}$$

For $\varepsilon > 0$ small enough (in particular such that $\frac{1}{\varepsilon} \gg \frac{1}{c}$, and such that $\frac{1}{\varepsilon}$ is an integer multiple of π , so that Ψ_ε is C^2), we have

$$\|(1 + \tilde{r})^{1+\sigma}h(x)\|_{L^\infty(\mathbb{R}^2)} = \|(1 + \tilde{r})^{1+\sigma}((-\Delta + 2)\Psi)(x)\|_{L^\infty(\mathbb{R}^2)} \leq K\varepsilon$$

and

$$\|(1 + \tilde{r})^{2+\sigma}|\nabla\Psi(x)|\|_{L^\infty(\mathbb{R}^2)} \geq 1/2.$$

Therefore, taking $\varepsilon \rightarrow 0$, we see that the estimate $|\nabla\Psi(x)| \leq \frac{C\|(1+\tilde{r})^{1+\sigma}h\|_{L^\infty(\mathbb{R}^2)}}{(1+\tilde{r})^{2+\sigma}}$ can not hold.

For our proof of the inversion of the linearized operator (Proposition 2.17 below), we did not choose the same norms $\|\cdot\|_{*,\sigma,d}$ and $\|\cdot\|_{**,\sigma',d}$ as in [16] (at the beginning of subsection 2.3). In particular, we require decays on the second derivatives for $\|\cdot\|_{**,\sigma',d}$. Our proof of the inversion of the linearized operator (the equivalent of Lemma 5.1 of [16]) will be different, and will follow more closely the proof of [8].

2.4.3 Estimates for the Gross-Pitaevskii kernels

We are interested here in solving the following equation on ψ , given a source term h and $c \in]0, \sqrt{2}[$:

$$-ic\partial_{x_2}\psi - \Delta\psi + 2\Re\psi = h.$$

It will appear in the inversion of the linearized operator around V . See Lemma 2.15 for the exact result. We give here a way to construct a solution formally. We will highlight all the important quantities, as well as all the difficulties that arise when trying to solve this equation rigorously.

In this subsection, we want to check that a solution of this equation, with $\psi = \psi_1 + i\psi_2$ and $h = h_1 + ih_2$ (where ψ_1, ψ_2, h_1, h_2 are real valued) can be written

$$\psi_1 = K_0 * h_1 + cH, \quad (2.4)$$

with H a function that satisfies

$$\partial_{x_j} H := K_j * h_2,$$

and

$$\partial_{x_j} \psi_2 = G_j - cK_j * h_1, \quad (2.5)$$

where similarly G_j satisfies

$$\partial_{x_k} G_j := (c^2 L_{j,k} - R_{j,k}) * h_2,$$

where, for $j, k \in \{1, 2\}$, $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2$,

$$\begin{aligned} \widehat{K}_0(\xi) &:= \frac{|\xi|^2}{|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2}, \\ \widehat{K}_j(\xi) &:= \frac{\xi_2\xi_j}{|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2}, \\ \widehat{L}_{j,k}(\xi) &:= \frac{\xi_2^2\xi_j\xi_k}{|\xi|^2(|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2)}, \end{aligned}$$

and

$$\widehat{R}_{j,k}(\xi) := \frac{\xi_j\xi_k}{|\xi|^2}.$$

We will check later on that, for continuous and sufficiently decaying functions h , these quantities are well defined, and that H, G_j, ψ_2 can be defined from there derivatives. The Gross-Pitaevskii kernels, $K_0, K_j, L_{j,k}$, and the Riesz kernels $R_{j,k}$ have been studied in [11], and we will recall some of the results obtained there.

We write the system in real and imaginary part:

$$\begin{cases} c\partial_{x_2}\psi_2 - \Delta\psi_1 + 2\psi_1 = h_1 \\ -c\partial_{x_2}\psi_1 - \Delta\psi_2 = h_2. \end{cases}$$

Now, taking the Fourier transform of the system, we have

$$\begin{cases} i\xi_2 c\widehat{\psi}_2 + (|\xi|^2 + 2)\widehat{\psi}_1 = \widehat{h}_1 \\ -i\xi_2 c\widehat{\psi}_1 + |\xi|^2\widehat{\psi}_2 = \widehat{h}_2, \end{cases}$$

and we write it

$$\begin{pmatrix} |\xi|^2 + 2 & ic\xi_2 \\ -ic\xi_2 & |\xi|^2 \end{pmatrix} \begin{pmatrix} \widehat{\psi}_1 \\ \widehat{\psi}_2 \end{pmatrix} = \begin{pmatrix} \widehat{h}_1 \\ \widehat{h}_2 \end{pmatrix}.$$

Here, we suppose that ψ is a tempered distributions and $h \in L^p(\mathbb{R}^2, \mathbb{C})$ for some $p > 1$.

Now, we want to invert the matrix, and for that, we have to divide by its determinant, $|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2$. For $0 < c < \sqrt{2}$, this quantity is zero only for $\xi = 0$. Thus, for $\xi \neq 0$,

$$\begin{pmatrix} \widehat{\psi}_1 \\ \widehat{\psi}_2 \end{pmatrix} = \frac{1}{|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2} \begin{pmatrix} |\xi|^2\widehat{h}_1 - ic\xi_2\widehat{h}_2 \\ (|\xi|^2 + 2)\widehat{h}_2 + ic\xi_2\widehat{h}_1 \end{pmatrix},$$

which implies that

$$\widehat{\psi}_1 = \frac{|\xi|^2\widehat{h}_1}{|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2} + \frac{-ic\xi_2\widehat{h}_2}{|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2}.$$

With the definition of K_0 , we have $\frac{|\xi|^2}{|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2}\widehat{h}_1 = \widehat{K}_0\widehat{h}_1$ and, defining the distribution H by $\widehat{H} = \frac{-ic\xi_2}{|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2}\widehat{h}_2$, we have, for $\xi \neq 0$,

$$\widehat{\partial_{x_j} H} = \frac{\xi_j\xi_2\widehat{h}_2}{|\xi|^4 + 2|\xi|^2 - c^2\xi_2^2} = \widehat{K}_j\widehat{h}_2.$$

Remark that $\frac{-i\xi_2}{|\xi|^4+2|\xi|^2-c^2\xi_2^2} \in L^{3/2}(\mathbb{R}^2, \mathbb{C})$ and thus is a tempered distribution.

Now, we have

$$\widehat{\partial_{x_j}\psi_2} = \frac{i\xi_j(|\xi|^2+2)\widehat{h_2}}{|\xi|^4+2|\xi|^2-c^2\xi_2^2} + \frac{-c\xi_j\xi_2\widehat{h_1}}{|\xi|^4+2|\xi|^2-c^2\xi_2^2}.$$

We check that $\frac{-c\xi_j\xi_2}{|\xi|^4+2|\xi|^2-c^2\xi_2^2}\widehat{h_1} = -c\widehat{K_j}\widehat{h_1}$, and we compute

$$\frac{|\xi|^2+2}{|\xi|^4+2|\xi|^2-c^2\xi_2^2} = \frac{1}{|\xi|^2} \left(1 - \frac{c^2\xi_2^2}{|\xi|^4+2|\xi|^2-c^2\xi_2^2} \right) = \frac{1}{|\xi|^2} - c^2 \frac{\xi_2^2}{|\xi|^2(|\xi|^4+2|\xi|^2-c^2\xi_2^2)},$$

thus, denoting $\widehat{G_j} = \frac{i\xi_j(|\xi|^2+2)}{|\xi|^4+2|\xi|^2-c^2\xi_2^2}\widehat{h_2}$, we have

$$\widehat{\partial_{x_k}G_j} := (c^2\widehat{L_{j,k}} - \widehat{R_{j,k}})\widehat{h_2}.$$

We therefore have that, at least formally, for $\xi \neq 0$, $-ic\partial_{x_2}\psi - \Delta\psi + 2\Re(\psi) - h(\xi) = 0$. We deduce that there exists $P \in \mathbb{C}[X_1, X_2]$ such that $-ic\partial_{x_2}\psi - \Delta\psi + 2\Re(\psi) - h = P$. Now, if the function ψ and h are such that the left hand side is bounded and goes to 0 at infinity, this implies that $P = 0$. This will hold under a condition on h (which will be $\int_{\mathbb{R}^2} h_2 = 0$ and some decay estimates, that ψ will inherit). Another remark is that ψ is here in part defined through its derivatives, and we need an argument to construct a primitive. See Lemma 2.15 for a rigorous proof of this construction. Remark that $-ic\partial_{x_2}\psi - \Delta\psi + 2\Re(\psi) = 0$ has some nonzero or unbounded polynomial solutions, for instance $\psi = i$ or $\psi = ix_2 - \frac{c}{2}$.

The kernels K_0, K_j and $L_{j,k}$ have been studied in details in [11], [12] and [13]. In particular, we recall the following result.

Theorem 2.12 ([11], Theorems 5 and 6) *For $\mathcal{K} \in \{K_0, K_j, L_{j,k}\}$ and any $0 < c_0 < \sqrt{2}$, there exist a constant $K(c_0) > 0$ such that, for all $0 < c < c_0$,*

$$|\mathcal{K}(x)| \leq \frac{K(c_0)}{|x|^{1/2}(1+|x|)^{3/2}}$$

and

$$|\nabla\mathcal{K}(x)| \leq \frac{K(c_0)}{|x|^{3/2}(1+|x|)^{3/2}}.$$

Proof This is the main result of Theorems 5 and 6 of [11]. We added the fact that the constant K is uniform in c , given that c is small. This can be easily shown by following the proof of Theorem 5 and 6 of [11], and verifying that the constants depends only on weighed L^∞ norms on $\widehat{\mathcal{K}}$ and its first derivatives, which are uniforms in c if $c > 0$ is small. The condition $c < c_0$ is taken in order to avoid $c \rightarrow \sqrt{2}$, where this does not hold (the singularity near $\xi = 0$ of $\widehat{\mathcal{K}}$ changes of order at the limit). Furthermore, the factor 1/2 for the growth near $x = 0$ is not at all optimal, but we will not require more here.

Remark that the speed in [11] is in the direction \vec{e}_1 , whereas it is in the direction \vec{e}_2 in our case, which explains the swap between ξ_2 and ξ_1 in the two papers. \square

We recall that $\tilde{r} = \min(r_1, r_{-1})$ with $r_{\pm 1} = |x \mp d\vec{e}_1|$. We give some estimates of convolution with these kernels.

Lemma 2.13 *Take $K \in \{K_0, K_j, L_{j,k}\}$ and $h \in C^0(\mathbb{R}^2, \mathbb{R})$, and suppose that, for some $\alpha > 0$, $\|h(1+\tilde{r})^\alpha\|_{L^\infty(\mathbb{R}^2)} < +\infty$. Then, for any $0 < \alpha' < \alpha$, there exists $C(\alpha, \alpha') > 0$ such that, for $0 < c < 1$, if either*

$$- \alpha < 2$$

$$- 2 < \alpha < 3, \forall (x_1, x_2) \in \mathbb{R}^2, h(-x_1, x_2) = h(x_1, x_2) \text{ and } \int_{\mathbb{R}^2} h = 0,$$

then

$$|K * h| \leq \frac{C(\alpha, \alpha')\|h(1+\tilde{r})^\alpha\|_{L^\infty(\mathbb{R}^2)}}{(1+\tilde{r})^{\alpha'}}.$$

Furthermore, if $\alpha < 3$ (without any other conditions), then

$$|\nabla K * h| \leq \frac{C(\alpha, \alpha')\|h(1+\tilde{r})^\alpha\|_{L^\infty(\mathbb{R}^2)}}{(1+\tilde{r})^{\alpha'}}.$$

The symmetry and $\int_{\mathbb{R}^2} h = 0$ in the case $2 < \alpha < 3$ could be removed, if we suppose instead that $\int_{\{x_1 \geq 0\}} h = \int_{\{x_1 \leq 0\}} h = 0$. In particular, if we suppose that $\forall (x_1, x_2) \in \mathbb{R}^2, h(x_1, x_2) = -h(x_1, -x_2)$, then the condition $\int_{\mathbb{R}^2} h = 0$ is automatically satisfied. We complete these estimates with some for $R_{j,k}$.

Lemma 2.14 *Take $h \in C^1(\mathbb{R}^2, \mathbb{R})$ with $\forall x = (x_1, x_2) \in \mathbb{R}^2, h(-x_1, x_2) = h(x_1, x_2)$, and suppose that for some $\alpha > 0$, $\|h(1 + \tilde{r})^\alpha\|_{L^\infty(\mathbb{R}^2)} + \|\nabla h(1 + \tilde{r})^\alpha\|_{L^\infty(\mathbb{R}^2)} < +\infty$. Then, for any $0 < \alpha' < \alpha$, for $0 < c < 1$, if either*

$$- \alpha < 2$$

$$- 2 < \alpha < 3 \text{ and } \int_{\mathbb{R}^2} h = 0,$$

then, there exists $C(\alpha, \alpha') > 0$ such that

$$|R_{j,k} * h| \leq \frac{C(\alpha, \alpha') (\|h(1 + \tilde{r})^\alpha\|_{L^\infty(\mathbb{R}^2)} + \|\nabla h(1 + \tilde{r})^\alpha\|_{L^\infty(\mathbb{R}^2)})}{(1 + \tilde{r})^{\alpha'}}.$$

We can now solve the problem

$$\begin{aligned} -ic\partial_{x_2}\psi - \Delta\psi + 2\Re(\psi) &= h, \\ \int_{\mathbb{R}^2} \Im(h) &= 0 \end{aligned}$$

in some suitable spaces. We define the norms, for $\sigma, \sigma' \in \mathbb{R}$,

$$\begin{aligned} \|\psi\|_{\otimes, \sigma, \infty} &:= \|(1 + \tilde{r})^{1+\sigma}\psi_1\|_{L^\infty(\mathbb{R}^2)} + \|(1 + \tilde{r})^{2+\sigma}\nabla\psi_1\|_{L^\infty(\mathbb{R}^2)} \\ &+ \|(1 + \tilde{r})^{2+\sigma}\nabla^2\psi_1\|_{L^\infty(\mathbb{R}^2)} + \|(1 + \tilde{r})^\sigma\psi_2\|_{L^\infty(\mathbb{R}^2)} \\ &+ \|(1 + \tilde{r})^{1+\sigma}\nabla\psi_2\|_{L^\infty(\mathbb{R}^2)} + \|(1 + \tilde{r})^{2+\sigma}\nabla^2\psi_2\|_{L^\infty(\mathbb{R}^2)} \end{aligned}$$

and

$$\begin{aligned} \|h\|_{\otimes \otimes, \sigma', \infty} &:= \|(1 + \tilde{r})^{1+\sigma'}h_1\|_{L^\infty(\mathbb{R}^2)} + \|(1 + \tilde{r})^{2+\sigma'}\nabla h_1\|_{L^\infty(\mathbb{R}^2)} \\ &+ \|(1 + \tilde{r})^{2+\sigma'}h_2\|_{L^\infty(\mathbb{R}^2)} + \|(1 + \tilde{r})^{2+\sigma'}\nabla h_2\|_{L^\infty(\mathbb{R}^2)}, \end{aligned}$$

as well as the spaces

$$\mathcal{E}_{\otimes \otimes, \sigma}^\infty := \{\psi \in C^2(\mathbb{R}^2, \mathbb{C}), \|\psi\|_{\otimes, \sigma, \infty} < +\infty, \forall (x_1, x_2) \in \mathbb{R}^2, \psi(x_1, x_2) = \psi(-x_1, x_2)\},$$

and

$$\mathcal{E}_{\otimes \otimes, \sigma'}^\infty := \{h \in C^1(\mathbb{R}^2, \mathbb{C}), \|h\|_{\otimes \otimes, \sigma', \infty} < +\infty, \forall (x_1, x_2) \in \mathbb{R}^2, h(x_1, x_2) = h(-x_1, x_2)\}.$$

The norms $\|\cdot\|_{\otimes, \sigma, \infty}$ and $\|\cdot\|_{*, \sigma}$ differ only on $\{\tilde{r} \leq 3\}$, and $\mathcal{E}_{\otimes \otimes, \sigma}^\infty$ has one less symmetry than $\mathcal{E}_{*, \sigma}$, but they are equivalent at infinity in position. Same remarks hold for $\|\cdot\|_{\otimes \otimes, \sigma', \infty}$ and $\|\cdot\|_{**, \sigma'}$ and their associated spaces. Remark that if $\chi \geq 0$ is a smooth cutoff function with value 0 on $\{\tilde{r} \leq R/2\}$ and 1 on $\{\tilde{r} \geq R\}$, then for any $\sigma \in \mathbb{R}$,

$$\|\psi\|_{*, \sigma} \leq K(R, \sigma) \|V\psi\|_{C^2(\{\tilde{r} \leq R\})} + K\|\chi\psi\|_{\otimes, \sigma, \infty}. \quad (2.6)$$

Lemma 2.15 *Given $1 > \sigma' > \sigma > 0$, there exists $K_1(\sigma, \sigma') > 0$ such that, for any $h \in \mathcal{E}_{\otimes \otimes, \sigma'}^\infty$ with $\int_{\mathbb{R}^2} \Im(h) = 0$ and $0 < c < 1$, there exists a unique solution to the problem*

$$-ic\partial_{x_2}\psi - \Delta\psi + 2\Re(\psi) = h,$$

in $\mathcal{E}_{\otimes \otimes, \sigma}^\infty$. This solution $\psi \in \mathcal{E}_{\otimes \otimes, \sigma}^\infty$ satisfies

$$\|\psi\|_{\otimes, \sigma, \infty} \leq K_1(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}.$$

Furthermore, if instead $\sigma \in]-1, 0[$ and $1 > \sigma' > \sigma$, there exists then $K_2(\sigma, \sigma') > 0$ such that, for any $h \in \mathcal{E}_{\otimes \otimes, \sigma'}^\infty$ with $\forall (x_1, x_2) \in \mathbb{R}^2, h(x_1, x_2) = \overline{h(x_1, -x_2)}$, there exists a unique solution to the problem

$$-ic\partial_{x_2}\psi - \Delta\psi + 2\Re(\psi) = h$$

in $\{\Psi \in \mathcal{E}_{\otimes \otimes, \sigma}^\infty, \forall (x_1, x_2) \in \mathbb{R}^2, \Psi(x_1, x_2) = \overline{\Psi(x_1, -x_2)}\}$. This solution $\psi \in \mathcal{E}_{\otimes \otimes, \sigma}^\infty$ satisfies

$$\|\psi\|_{\otimes, \sigma, \infty} \leq K_2(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}.$$

The case $\sigma \in]-1, 0[$ is particular and such a norm will be used only in Lemma 2.18 (if $\|\psi\|_{\otimes, \sigma, \infty} < +\infty$ for $\sigma < 0$, the function ψ is not necessarily bounded for instance). Remark that the condition $\int_{\mathbb{R}^2} \mathfrak{I}m(h) = 0$ is automatically satisfied if $\forall (x_1, x_2) \in \mathbb{R}^2, h(x_1, x_2) = \overline{h(x_1, -x_2)}$.

Proof For $1 > \sigma' > \sigma > -1$, we write in real and imaginary parts $h = h_1 + ih_2$. We define, for $j \in \{1, 2\}$,

$$\Psi_{1,j} := K_0 * \partial_{x_j} h_1 + cK_j * h_2.$$

If $1 > \sigma' > \sigma > 0$, since $\partial_{x_j} h_1, h_2 \in L^1(\mathbb{R}^2)$ (because $\sigma' > 0$ and $h \in \mathcal{E}_{\otimes, \sigma'}^\infty$), and $\int_{\mathbb{R}^2} h_2 = \int_{\mathbb{R}^2} \partial_{x_2} h_1 = 0$, by Lemma 2.13 (applied with $0 < \alpha = 2 + \sigma' < 3, 0 < \alpha' = 2 + \sigma < \alpha$), the function $\Psi_{1,2}$ is well defined and satisfies

$$|\nabla \Psi_{1,2}| + |\Psi_{1,2}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}.$$

This result still holds if $\sigma \in]-1, 0[$ and $1 > \sigma' > \sigma$, since $0 < \alpha = 2 + \sigma' < 3, 0 < \alpha' = 2 + \sigma < \alpha$. We check, still with Lemma 2.13 (applied with $0 < \alpha = 2 + \sigma' < 3, 0 < \alpha' = 2 + \sigma < \alpha$), that $\Psi_{1,1}$ is well defined and

$$|\nabla \Psi_{1,1}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}.$$

If $\sigma \in]-1, 0[$, we have $|\Psi_{1,1}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}$ by Lemma 2.13 ($2 + \sigma < 2$). But since $\partial_{x_1} h_1$ is not even in x_1 , we can not apply Lemma 2.13 to estimate $\Psi_{1,1}$ with the same decay in the case $\sigma > 0$. However, we check that the estimate holds if $|x + d\vec{e}_1| \leq 1$ or $|x - d\vec{e}_1| \leq 1$, and that otherwise

$$|\Psi_{1,1}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}} + \left| K(x + d\vec{e}_1) \int_{\{y_1 \leq 0\}} \partial_{x_1} h(y) dy + K(x - d\vec{e}_1) \int_{\{y_1 \geq 0\}} \partial_{x_1} h(y) dy \right|.$$

Since

$$\int_{\{y_1 \leq 0\}} \partial_{x_1} h(y) dy = - \int_{\{y_1 \geq 0\}} \partial_{x_1} h(y) dy = \int_{\mathbb{R}} h(0, y_2) dy_2,$$

and

$$\left| \int_{\mathbb{R}} h(0, y_2) dy_2 \right| \leq \int_{\mathbb{R}} \frac{\|h\|_{\otimes, \sigma', \infty}}{(1 + \tilde{r})^{1+\sigma'}} dy_2 \leq c^\sigma \|h\|_{\otimes, \sigma', \infty} \int_{\mathbb{R}} \frac{dy_2}{(1 + |y_2|)^{1+\sigma'-\sigma}} \leq K(\sigma, \sigma') c^\sigma \|h\|_{\otimes, \sigma', \infty},$$

we have

$$\begin{aligned} & \left| K(x + d\vec{e}_1) \int_{\{y_1 \leq 0\}} \partial_{x_1} h(y) dy + K(x - d\vec{e}_1) \int_{\{y_1 \geq 0\}} \partial_{x_1} h(y) dy \right| \\ & \leq K(\sigma, \sigma') |K(x + d\vec{e}_1) - K(x - d\vec{e}_1)| c^\sigma \|h\|_{\otimes, \sigma', \infty}. \end{aligned}$$

By Theorem 2.12, if $|x + d\vec{e}_1|, |x - d\vec{e}_1| \geq 1$,

$$|K(x + d\vec{e}_1) - K(x - d\vec{e}_1)| \leq \frac{K}{(1 + |x + d\vec{e}_1|)^2} + \frac{K}{(1 + |x - d\vec{e}_1|)^2} \leq \frac{K}{(1 + \tilde{r})^2},$$

and, if $\tilde{r} \leq 3d$,

$$|K(x + d\vec{e}_1) - K(x - d\vec{e}_1)| \leq \frac{K}{(1 + \tilde{r})^2} \leq \frac{Kd}{(1 + \tilde{r})^3},$$

or if $\tilde{r} \geq 3d$,

$$|K(x + d\vec{e}_1) - K(x - d\vec{e}_1)| \leq Kd \sup_{\nu \in [-d, d]} |\nabla K(x + \nu \vec{e}_1)| \leq \frac{Kd}{(1 + \tilde{r})^3},$$

therefore, by interpolation,

$$|K(x + d\vec{e}_1) - K(x - d\vec{e}_1)| \leq \left(\frac{K}{(1 + \tilde{r})^2} \right)^{1-\sigma} \times \left(\frac{Kd}{(1 + \tilde{r})^3} \right)^\sigma \leq \frac{Kd^\sigma}{(1 + \tilde{r})^{2+\sigma}}.$$

We deduce

$$\begin{aligned}
& \left| K(x + d\vec{e}_1) \int_{\{y_1 \leq 0\}} \partial_{x_1} h(y) dy + K(x - d\vec{e}_1) \int_{\{y_1 \geq 0\}} \partial_{x_1} h(y) dy \right| \\
& \leq K(\sigma, \sigma') |K(x + d\vec{e}_1) - K(x - d\vec{e}_1)| c^\sigma \|h\|_{\otimes \otimes, \sigma', \infty} \\
& \leq \frac{K(\sigma, \sigma') (dc)^\sigma}{(1 + \tilde{r})^{2+\sigma}} \|h\|_{\otimes \otimes, \sigma', \infty} \\
& \leq \frac{K(\sigma, \sigma')}{(1 + \tilde{r})^{2+\sigma}} \|h\|_{\otimes \otimes, \sigma', \infty}.
\end{aligned}$$

Combining the previous estimates, we conclude that, for $j \in \{1, 2\}$,

$$|\nabla \Psi_{1,j}| + |\Psi_{1,j}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}. \quad (2.7)$$

Let us show that $\Psi_{1,j} \in C^1(\mathbb{R}^2, \mathbb{C})$ by dominated convergence theorem (it is not clear at this point that $\nabla \Psi_{1,j}$ is continuous). For $x, \varepsilon \in \mathbb{R}^2$,

$$\begin{aligned}
\nabla \Psi_{1,j}(x + \varepsilon) - \nabla \Psi_{1,j}(x) &= \int_{\mathbb{R}^2} \nabla K_0(y) (\partial_{x_j} h_1(x + \varepsilon - y) - \partial_{x_j} h_1(x - y)) dy, \\
&+ c \int_{\mathbb{R}^2} \nabla K_j(y) (h_2(x + \varepsilon - y) - h_2(x - y)) dy.
\end{aligned}$$

We check that for any $y \in \mathbb{R}^2$, $\partial_{x_j} h_1(x + \varepsilon - y) - \partial_{x_j} h_1(x - y) \rightarrow 0$, $h_2(x + \varepsilon - y) - h_2(x - y) \rightarrow 0$ pointwise when $|\varepsilon| \rightarrow 0$ (by continuity of $\partial_{x_j} h_1$ and h_2), and

$$\begin{aligned}
& |\nabla K_0(y) (\partial_{x_j} h_1(x + \varepsilon - y) - \partial_{x_j} h_1(x - y))| \\
& + c |\nabla K_j(y) (h_2(x + \varepsilon - y) - h_2(x - y))| \\
& \leq K(\sigma) \frac{|\nabla K_0(y)|}{(1 + \tilde{r}(x - y))^{2+\sigma'}} \|\partial_{x_j} h_1(1 + \tilde{r})^{2+\sigma'}\|_{L^\infty(\mathbb{R}^2)} \\
& + K(\sigma) \frac{c |\nabla K_j(y)|}{(1 + \tilde{r}(x - y))^{2+\sigma'}} \|h_2(1 + \tilde{r})^{2+\sigma'}\|_{L^\infty(\mathbb{R}^2)} \\
& \leq K(\sigma, x) \frac{|\nabla K_0(y)|}{(1 + \tilde{r}(y))^{2+\sigma'}} \|\partial_{x_j} h_1(1 + \tilde{r})^{2+\sigma'}\|_{L^\infty(\mathbb{R}^2)} \\
& + K(\sigma, x) \frac{c |\nabla K_j(y)|}{(1 + \tilde{r}(y))^{2+\sigma'}} \|h_2(1 + \tilde{r})^{2+\sigma'}\|_{L^\infty(\mathbb{R}^2)} \in L^1(\mathbb{R}^2)
\end{aligned}$$

for $|\varepsilon| \leq 1$ and a constant $K(\sigma, x) > 0$, giving the domination.

Now, we check, by taking their Fourier transforms, that $\partial_{x_1} \Psi_{1,2} = \partial_{x_2} \Psi_{1,1} \in L^2(\mathbb{R}^2, \mathbb{C})$ (see the computations at the beginning of subsection 2.4.3), and thus the integral of the vector field $\begin{pmatrix} \Psi_{1,1} \\ \Psi_{1,2} \end{pmatrix}$ on any closed curve of \mathbb{R}^2 is 0. For a large constant $D > 0$, taking, for $x_1 \in \mathbb{R}$, the path

$$\{(x_1, y), y \in [-D, D]\} \cup \{Y = (y_1, y_2) \in \mathbb{R}^2, |(x_1, 0) - Y| = D, y_1 \geq 0\},$$

since $|\Psi_{1,2}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}$ and

$$\int_{\{Y=(y_1, y_2) \in \mathbb{R}^2, |(x_1, 0) - Y| = D, y_1 \geq 0\}} |\Psi_{1,2}| \leq \frac{K(c, \sigma, \sigma', h)}{D^{1+\sigma}} \rightarrow 0$$

when $D \rightarrow \infty$ (since $1 + \sigma > 0$), we deduce that

$$\int_{-\infty}^{+\infty} \Psi_{1,2}(x_1, y_2) dy_2 = 0. \quad (2.8)$$

We then define for $(x_1, x_2) \in \mathbb{R}^2$,

$$\psi_1(x_1, x_2) = \int_{+\infty}^{x_2} \Psi_{1,2}(x_1, y_2) dy_2,$$

and thus, if $x_2 < 0$,

$$\psi_1(x_1, x_2) = \int_{-\infty}^{x_2} \Psi_{1,2}(x_1, y_2) dy_2.$$

With (2.7), we check that $\psi_1 \in C^1(\mathbb{R}^2, \mathbb{C})$, and by simple integration from infinity using the equations above (with $\tilde{r} = \min(|x - d_c \vec{e}_1|, |x + d_c \vec{e}_1|)$, and since $1 + \sigma > 0$), that

$$|\psi_1| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes\otimes, \sigma', \infty}}{(1 + \tilde{r})^{1+\sigma}}.$$

Furthermore, we check that

$$\partial_{x_2} \psi_1 = \Psi_{1,2} \in C^1(\mathbb{R}^2, \mathbb{C}),$$

and (by taking their Fourier transforms)

$$\partial_{x_1} \psi_1 = \Psi_{1,1} \in C^1(\mathbb{R}^2, \mathbb{C}),$$

therefore $\psi_1 \in C^2(\mathbb{R}^2, \mathbb{C})$, and by (2.7),

$$|\nabla \psi_1| \leq |\Psi_{1,1}| + |\Psi_{1,2}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes\otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}.$$

For $j, k \in \{1, 2\}$, we have $\partial_{x_j x_k}^2 \psi_1 = \partial_{x_j} \Psi_{1,k}$, thus, by (2.7),

$$|\nabla^2 \psi_1| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes\otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}.$$

Now, we define

$$\Psi_{2,j,k} := (c^2 L_{j,k} - R_{j,k}) * h_2 - c K_j * \partial_{x_k} h_1.$$

In the case $1 > \sigma' > \sigma > 0$, $\partial_{x_k} h_1, h_2 \in L^1(\mathbb{R}^2)$ and since $\int_{\mathbb{R}^2} h_2 = \int_{\mathbb{R}^2} \partial_{x_k} h_1 = 0$, by Lemmas 2.13 and 2.14 (for $\alpha = 2 + \sigma' < 3$, $\alpha' = 2 + \sigma < \alpha$, and the same variant for $K_j * \partial_{x_1} h_1$ as in the proof of (2.7)), this function is well defined in $L^\infty(\mathbb{R}^2, \mathbb{C})$, and satisfies,

$$|\Psi_{2,j,k}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes\otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}. \quad (2.9)$$

We check, as for the proof of (2.7), that this result holds if $\sigma \in]-1, 0[$ and $1 > \sigma' > \sigma$.

Remark here that we do not have $\Psi_{2,j,k} \in C^1(\mathbb{R}^2, \mathbb{C})$, since in Lemma 2.14, the estimate on $R_{j,k} * h_2$ uses ∇h_2 in the norm (showing that $\Psi_{2,j,k} \in C^1(\mathbb{R}^2, \mathbb{C})$ would require estimates on $\nabla^2 h_2$). However, we have that $\Psi_{2,j,k} \in C^0(\mathbb{R}^2, \mathbb{C})$ by dominated convergence and continuity of h_2 and $\partial_{x_k} h_1$ (as for $\nabla \Psi_{1,j}$). Furthermore, we check (by taking their Fourier transforms) that $\partial_{x_1} \Psi_{2,j,2} = \partial_{x_2} \Psi_{2,j,1}$ in the distribution sense. We infer that the integral of $\begin{pmatrix} \Psi_{2,j,1} \\ \Psi_{2,j,2} \end{pmatrix}$ on any bounded closed curve of \mathbb{R}^2 is 0. Indeed, taking χ_n a mollifier sequence, then $\chi_n * \Psi_{2,j,1}, \chi_n * \Psi_{2,j,2} \in C^1(\mathbb{R}^2, \mathbb{C})$,

$$\partial_{x_1} (\chi_n * \Psi_{2,j,2}) - \partial_{x_2} (\chi_n * \Psi_{2,j,1}) = \chi_n * (\partial_{x_1} \Psi_{2,j,2} - \partial_{x_2} \Psi_{2,j,1}) = 0,$$

therefore, for any closed curve \mathcal{C} , the integral of the field $\begin{pmatrix} \chi_n * \Psi_{2,j,1} \\ \chi_n * \Psi_{2,j,2} \end{pmatrix}$ is 0. Using $\chi_n * \Psi_{2,j,k} \rightarrow \Psi_{2,j,k}$ pointwise (by continuity of $\Psi_{2,j,k}$) and the domination

$$\|\chi_n * \Psi_{2,j,1}\|_{L^\infty(\mathbb{R}^2)} \leq \|\Psi_{2,j,1}\|_{L^\infty(\mathbb{R}^2)} < +\infty,$$

we infer that this result holds for $\begin{pmatrix} \Psi_{2,j,1} \\ \Psi_{2,j,2} \end{pmatrix}$. We deduce, as for the proof of (2.8), that

$$\int_{-\infty}^{+\infty} \Psi_{2,j,2}(x_1, y_2) dy_2 = 0. \quad (2.10)$$

We then define for $(x_1, x_2) \in \mathbb{R}^2, j \in \{1, 2\}$,

$$\Psi_{2,j}(x_1, x_2) = \int_{+\infty}^{x_2} \Psi_{2,j,2}(x_1, y_2) dy_2,$$

and if $x_2 < 0$, by (2.10),

$$\Psi_{2,j}(x_1, x_2) = \int_{-\infty}^{x_2} \Psi_{2,j,2}(x_1, y_2) dy_2.$$

With arguments similar to the proof for $\Psi_{1,j}$, we check that $\Psi_{2,j} \in C^1(\mathbb{R}^2, \mathbb{C})$ with $\partial_{x_k} \Psi_{2,j} = \Psi_{2,j,k}$,

$$|\Psi_{2,j}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}}{(1 + \tilde{r})^{1+\sigma}},$$

as well as

$$|\nabla \Psi_{2,j}| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}}.$$

Finally, since $\partial_{x_1} \Psi_{2,2} = \Psi_{2,2,1} = \Psi_{2,1,2} = \partial_{x_2} \Psi_{2,1} \in L^2(\mathbb{R}^2, \mathbb{C})$ (by taking their Fourier transforms, it follows from $R_{j,k} = R_{k,j}$, $L_{j,k} = L_{k,j}$ and $\hat{K}_j \xi_k = \hat{K}_k \xi_j$), we have, as before, that

$$\int_{-\infty}^{+\infty} \Psi_{2,2}(x_1, y_2) dy_2 = 0.$$

We define

$$\psi_2(x_1, x_2) = \int_{+\infty}^{x_2} \Psi_{2,2}(x_1, y_2) dy_2,$$

and thus, if $x_2 < 0$,

$$\psi_2(x_1, x_2) = \int_{-\infty}^{x_2} \Psi_{2,2}(x_1, y_2) dy_2.$$

We check, as previously, by integration from infinity, that $\psi_2 \in C^2(\mathbb{R}^2, \mathbb{C})$, $\partial_{x_j x_k}^2 \psi_2 = \Psi_{2,j,k}$, and

$$|\nabla^2 \psi_2| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}}{(1 + \tilde{r})^{2+\sigma}},$$

$$|\nabla \psi_2| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}}{(1 + \tilde{r})^{1+\sigma}},$$

as well as (if $\sigma > 0$)

$$|\psi_2| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}}{(1 + \tilde{r})^\sigma}.$$

Remark that if h satisfies $\forall (x_1, x_2) \in \mathbb{R}^2, h(x_1, x_2) = \overline{h(x_1, -x_2)}$, then by the definition of ψ_1 and ψ_2 above, for $\psi = \psi_1 + i\psi_2$, we have that $\forall (x_1, x_2) \in \mathbb{R}^2, \psi(x_1, x_2) = \overline{\psi(x_1, -x_2)}$. Therefore, in the case $\sigma \in]-1, 0[$, since $\forall (x_1, x_2) \in \mathbb{R}^2, \psi_2(x_1, x_2) = -\psi_2(x_1, -x_2)$, we have $\psi_2(x_1, 0) = 0$, and we integrate $\nabla \psi_2$ from the line $\{x_2 = 0\}$ instead of infinity to show that $|\psi_2| \leq \frac{K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}}{(1 + \tilde{r})^\sigma}$.

We deduce that, in either cases, $\psi = \psi_1 + i\psi_2 \in \mathcal{E}_{\otimes, \sigma}^\infty$, and it satisfies

$$\|\psi\|_{\otimes, \sigma, \infty} \leq K(\sigma, \sigma') \|h\|_{\otimes \otimes, \sigma', \infty}.$$

Now, let us show that $-ic\partial_{x_2} \psi - \Delta \psi + 2\mathfrak{R}\mathfrak{e}(\psi) = h$. From the computations at the beginning of subsection 2.4.3, we check that the Fourier transform (in the distribution sense) of both side of the equation are equals on $\{\xi \in \mathbb{R}^2, \xi \neq 0\}$ (remark that they are both in $L^p(\mathbb{R}^2, \mathbb{C})$ for some $p > 2$ large enough). This implies that

$$\text{Supp} \left(-ic\partial_{x_2} \psi - \widehat{\Delta \psi} + 2\mathfrak{R}\mathfrak{e}(\psi) - h \right) \subset \{0\},$$

and thus $-ic\partial_{x_2}\psi - \Delta\psi + 2\Re(\psi) - h = P \in \mathbb{C}[X_1, X_2]$. With the decay estimates on ψ and h , we check that P is bounded and goes to 0 at infinity (since $\sigma, \sigma' > -1$), thus $P = 0$.

Finally, if $\tilde{\psi} \in \mathcal{E}_{\otimes, \sigma}^\infty$ satisfies $-ic\partial_{x_2}\tilde{\psi} - \Delta\tilde{\psi} + 2\Re(\tilde{\psi}) = h$, then $\psi - \tilde{\psi} \in C^2(\mathbb{R}^2, \mathbb{C})$ and

$$(-ic\partial_{x_2} - \Delta + 2\Re)(\psi - \tilde{\psi}) = 0.$$

With the computations at the beginning of subsection 2.4.3, since $\psi - \tilde{\psi}$ is a tempered distribution, we check that $\text{Supp } \psi - \tilde{\psi} \subset \{0\}$, therefore $\psi - \tilde{\psi} = P \in \mathbb{C}[X_1, X_2]$. If $\sigma > 0$, since $\psi - \tilde{\psi}$ goes to 0 at infinity, $P = 0$. If $\sigma \in]-1, 0[$, then $P = i\lambda$ for some $\lambda \in \mathbb{R}$ ($\Re(\psi - \tilde{\psi}) \rightarrow 0$ at infinity and $\tilde{r}^{-\sigma}\Im(\psi - \tilde{\psi})$ is bounded), and by the symmetry on $\psi, \tilde{\psi}$ we have in that case, $\lambda = 0$. This shows the uniqueness of a solution in $\mathcal{E}_{\otimes, \sigma}^\infty$ (with the symmetry if $\sigma \in]-1, 0[$), and thus concludes the proof of this lemma. \square

2.5 Reduction of the problem

2.5.1 Inversion of the linearized operator

One of the key element in the inversion of the linearized operator is the computation of the kernel for only one vortex. The kernel of the linearized operator around one vortex has been studied in [7], with the following result.

Theorem 2.16 (Theorem 1.2 of [7]) *Consider the linearized operator around one vortex of degree $\varepsilon = \pm 1$,*

$$L_{V_\varepsilon}(\Phi) := -\Delta\Phi - (1 - |V_\varepsilon|^2)\Phi + 2\Re(\overline{V_\varepsilon}\Phi)\overline{V_\varepsilon}.$$

Suppose that

$$\|\Phi\|_{H_{V_\varepsilon}}^2 := \int_{\mathbb{R}^2} |\nabla\Phi|^2 + (1 - |V_\varepsilon|^2)|\Phi|^2 + \Re(\overline{V_\varepsilon}\Phi) < +\infty$$

and

$$L_{V_\varepsilon}(\Phi) = 0.$$

Then, there exist two constants $c_1, c_2 \in \mathbb{R}$ such that

$$\Phi = c_1\partial_{x_1}V_\varepsilon + c_2\partial_{x_2}V_\varepsilon.$$

This result describes the kernel of L_{V_ε} that will appear in the proof of Proposition 2.17. It shows that the kernel in $H_{V_\varepsilon} := \{\Phi \in H_{\text{loc}}^1(\mathbb{R}^2), \|\Phi\|_{H_{V_\varepsilon}} < +\infty\}$ contains only the two elements we expect: $\partial_{x_1}V_\varepsilon, \partial_{x_2}V_\varepsilon$, which are due to the invariance by translation of (GP). One of the directions will be killed by the symmetry and the other one by the orthogonality.

Now, we shall invert the linear part $\eta L(\Phi) + (1 - \eta)V L'(\Psi)$. We recall that $\Phi = V\Psi$. We first state an a priori estimate result. We recall the definitions, for $\sigma, \sigma' \in]0, 1[$,

$$\mathcal{E}_{*, \sigma, d} =$$

$$\{\Phi = V\Psi \in C^2(\mathbb{R}^2, \mathbb{C}), \|\Psi\|_{*, \sigma, d} < +\infty; \langle \Phi, Z_d \rangle = 0; \forall x \in \mathbb{R}^2, \Psi(x_1, x_2) = \overline{\Psi(x_1, -x_2)} = \Psi(-x_1, x_2)\}$$

and

$$\mathcal{E}_{**, \sigma', d} = \{Vh \in C^1(\mathbb{R}^2, \mathbb{C}), \|h\|_{**, \sigma', d} < +\infty\}.$$

Proposition 2.17 *For $1 > \sigma' > \sigma > 0$, consider the problem, in the distribution sense*

$$\begin{cases} \eta L(\Phi) + (1 - \eta)V L'(\Psi) = Vh \\ \Phi = V\Psi \in \mathcal{E}_{*, \sigma}, Vh \in \mathcal{E}_{**, \sigma'}. \end{cases}$$

Then, there exist constants $c_0(\sigma, \sigma') > 0$ small and $C(\sigma, \sigma') > 0$ depending only on σ and σ' , such that, for any solution of this problem with $0 < c \leq c_0(\sigma, \sigma')$, $\frac{1}{2} < cd < 2$, it holds

$$\|\Psi\|_{*, \sigma, d} \leq C(\sigma, \sigma')\|h\|_{**, \sigma', d}.$$

Proof This proof is similar to the ones done in [8] for the inversion of their linearized operator. The main difference is that we have a transport term. Fix $1 > \sigma' > \sigma > 0$. We argue by contradiction. Suppose that for given $1 > \sigma' > \sigma > 0$, there is no threshold $c_0(\sigma, \sigma') > 0$ such that, if $0 < c \leq c_0(\sigma, \sigma')$ we have $\|\Psi\|_{*,\sigma,d} \leq C(\sigma, \sigma') \|h\|_{**,\sigma',d}$. We can then find a sequence of $c_n \rightarrow 0$ (and so $d_n \rightarrow \infty$), functions $\Phi_n = V\Psi_n \in \mathcal{E}_{*,\sigma}$ and $Vh_n \in \mathcal{E}_{**,\sigma'}$ solutions of the problem and such that

$$\|\Psi_n\|_{*,\sigma,d_n} = 1$$

and

$$\|h_n\|_{**,\sigma',d_n} \rightarrow 0.$$

We look in the region $\Sigma := \{x_1 \geq 0\}$ thanks to the symmetry $\Psi(x_1, x_2) = \Psi(-x_1, x_2)$. The orthogonality condition of $\mathcal{E}_{*,\sigma}$ becomes $2\Re \int_{\Sigma} \overline{\Phi_n} Z_{d_n} = 0$.

Step 1. Inner estimates.

The problem can be written (using $VL'(\Psi_n) = -(E - ic_n \partial_{x_2} V)\Psi_n + L(\Phi_n)$ from Lemma 2.7) as

$$Vh_n = L(\Phi_n) - (1 - \eta)(E - ic_n \partial_{x_2} V)\Psi_n.$$

First, we recall that V and E are depending on n . The sequence $(\Phi_n(\cdot + d_n \vec{e}_1^\rightarrow))_{n \in \mathbb{N}}$ is equicontinuous and bounded ($1 = \|\Psi_n\|_{*,\sigma,d}$ controls Φ_n and its derivatives in $L^\infty(\mathbb{R}^2)$ uniformly in n).

Such a function Φ_n , as a solution of

$$\Delta \Phi_n = -(1 - |V|^2)\Phi_n + 2\Re \epsilon(\bar{V}\Phi_n)V - ic \partial_{x_2} \Phi_n - (1 - \eta)(E - ic_n \partial_{x_2} V)\Psi_n - Vh_n \quad (2.11)$$

in the distribution sense, by Theorem 8.8 of [9] is $H_{\text{loc}}^2(\mathbb{R}^2)$ (since the right hand side is $C^0(\mathbb{R}^2)$). Furthermore, still by Theorem 8.8 of [9], we have, for $x \in \mathbb{R}^2$,

$$\|\Phi_n\|_{H^2(B(x,1))} \leq K(\|\Phi_n\|_{H^1(B(x,2))} + \|\Delta \Phi_n\|_{L^2(B(x,2))}).$$

By $\|\Psi_n\|_{*,\sigma,d} = 1$, the quantities $\|\Phi_n\|_{L^\infty(B(x,2))}$, $\|\nabla \Phi_n\|_{L^\infty(B(x,2))}$ and $\|\Delta \Phi_n\|_{L^\infty(B(x,2))}$ are bounded by a constant independent of n . Therefore, $(\Phi_n)_{n \in \mathbb{N}}$ is bounded in $H_{\text{loc}}^2(\mathbb{R}^2)$.

We deduce, by compact embedding, that there exists a function Φ such that $\Phi_n(\cdot + d_n \vec{e}_1^\rightarrow) \rightarrow \Phi$ in $H_{\text{loc}}^1(\mathbb{R}^2)$ (up to a subsequence).

Now, since $L(\Phi_n) = -\Delta \Phi_n - (1 - |V|^2)\Phi_n + 2\Re \epsilon(\bar{V}\Phi_n)V - ic \partial_{x_2} \Phi_n$, we have, in the weak sense,

$$\Delta \Phi_n + Vh_n = -(1 - |V|^2)\Phi_n + 2\Re \epsilon(\bar{V}\Phi_n)V - ic_n \partial_{x_2} \Phi_n - (1 - \eta)(E - ic_n \partial_{x_2} V)\Psi_n,$$

therefore $\Delta \Phi_n(\cdot + d_n \vec{e}_1^\rightarrow) + Vh_n(\cdot + d_n \vec{e}_1^\rightarrow)$ is equicontinuous and bounded uniformly and then, by Ascoli's Theorem, up to a subsequence converges to a limit l in $\mathcal{C}_{\text{loc}}^0(\mathbb{R}^2)$. Since $Vh_n(\cdot + d_n \vec{e}_1^\rightarrow) \rightarrow 0$ in $\mathcal{C}_{\text{loc}}^0(\mathbb{R}^2)$ by $\|h_n\|_{**,\sigma',d} \rightarrow 0$ and $\Delta \Phi_n(\cdot + d_n \vec{e}_1^\rightarrow) \rightarrow \Delta \Phi$ in the distribution sense, this limit must be $\Delta \Phi$ (in the $H_{\text{loc}}^{-1}(\mathbb{R}^2)$ sense).

We have locally uniformly that $Vh_n(\cdot + d_n \vec{e}_1^\rightarrow) \rightarrow 0$ because $\|h_n\|_{**,\sigma',d} \rightarrow 0$ and $|V| \leq 1$, and we have, from Lemma 2.4, that $E(y + d_n \vec{e}_1^\rightarrow) \rightarrow 0$ and $V(y + d_n \vec{e}_1^\rightarrow) \rightarrow V_1(y)$ when $n \rightarrow \infty$ locally uniformly. Lastly, $\partial_{x_2} \Phi_n$ and $(1 - \eta)\partial_{x_2} V \Psi_n$ are uniformly bounded in \mathbb{R}^2 independently of n . Therefore when we take the locally uniform limit when $d_n \rightarrow \infty$ in

$$(Vh_n)(y + d_n \vec{e}_1^\rightarrow) = (L(\Phi_n))(y + d_n \vec{e}_1^\rightarrow) - ((1 - \eta)(E - ic_n \partial_{x_2} V)\Psi_n)(y + d_n \vec{e}_1^\rightarrow),$$

we have (in the distribution sense)

$$L_{V_1}(\Phi) = 0.$$

Using $\partial_d V(\cdot + d \vec{e}_1^\rightarrow) \rightarrow -\partial_{x_1} V_1(\cdot)$ locally uniformly from Lemma 2.4, we show that

$$0 = 2\Re \int_{\Sigma} \overline{\Phi_n} Z_d \rightarrow 2\langle \Phi | \tilde{\eta}(\cdot/4) \partial_{x_1} V_1 \rangle$$

since Z_d is compactly supported around 0 when we take the equation in $y + d_n \vec{e}_1$. The problem at the limit $n \rightarrow \infty$ becomes (in the $H_{\text{loc}}^{-1}(\mathbb{R}^2)$ sense)

$$\begin{cases} L_{V_1}(\Phi) = 0 \\ \langle \Phi | \tilde{\eta}(\frac{\cdot}{4}) \partial_{x_1} V_1 \rangle = 0, \end{cases}$$

with $\Phi = V_1 \Psi$ (since $V(y + d \vec{e}_1) \rightarrow V_1(y)$ from Lemma 2.4). We can check that $\|\Phi\|_{H_{V_1}} < +\infty$, therefore, we can apply Theorem 2.16. We deduce that

$$\Phi = c_1 \partial_{x_1} V_1 + c_2 \partial_{x_2} V_1$$

for some constants $c_1, c_2 \in \mathbb{R}$.

Since $\forall x \in \mathbb{R}^2, \Psi_n(x_1, x_2) = \overline{\Psi_n(x_1, -x_2)}$, we have $\forall y \in \mathbb{R}^2, \Phi(y_1, y_2) = \overline{\Phi(y_1, -y_2)}$. The function $\partial_{x_1} V_1$ enjoys also this symmetry, therefore so does $c_2 \partial_{x_2} V_1$. It is possible only if $c_2 = 0$. The orthogonality condition then imposes

$$c_1 \int_{\Sigma} |\partial_{x_1} V_1(y)|^2 \tilde{\eta}\left(\frac{y}{4}\right) dy = 0,$$

implying that $c_1 = 0$. Hence

$$\Phi_n(\cdot + d_n \vec{e}_1) \rightarrow 0$$

in $\mathcal{C}_{\text{loc}}^1(\mathbb{R}^2)$. By equation (2.11) and standard elliptic estimates, this convergence also hold in $\mathcal{C}_{\text{loc}}^2(\mathbb{R}^2)$. The same proof works for the z coordinate (around the center of the -1 vortex). As a consequence, for any $R > 0$, we have

$$\|\Phi_n\|_{L^\infty(\{\tilde{r} \leq R\})} + \|\nabla \Phi_n\|_{L^\infty(\{\tilde{r} \leq R\})} + \|\nabla^2 \Phi_n\|_{L^\infty(\{\tilde{r} \leq R\})} \rightarrow 0 \quad (2.12)$$

as $n \rightarrow \infty$. With this result, to obtain a contradiction (which will be $\|\Psi_n\|_{*,\sigma,d} \rightarrow 0$) we still need to have estimates near the infinity in space.

Step 2. Outer computations.

Thanks to the previous step, we can take a cutoff to look only at the infinity in space. For $R \geq 4$, we define χ_R a smooth cutoff function with value $\chi_R(x) = 1$ if $\tilde{r} \geq R$ and $\chi_R(x) = 0$ if $\tilde{r} \leq \frac{R}{2}$, with $|\nabla \chi_R| \leq \frac{4}{R}$. We then define

$$\tilde{\Psi}_n := \chi_R \Psi_n,$$

$$\tilde{h}_n := \chi_R h_n$$

and we choose χ_R such that $\tilde{\Psi}_n$ and \tilde{h}_n enjoy the same symmetries than Ψ_n and h_n respectively. We compute on $\mathbb{R}^2 \setminus (B(d_n \vec{e}_1, R) \cup B(-d_n \vec{e}_1, R))$:

$$\nabla \tilde{\Psi}_n = \nabla \chi_R \Psi_n + \chi_R \nabla \Psi_n = \nabla \Psi_n,$$

$$\Delta \tilde{\Psi}_n = \Delta \chi_R \Psi_n + 2 \nabla \chi_R \nabla \Psi_n + \chi_R \Delta \Psi_n = \Delta \Psi_n.$$

We deduce that $\tilde{\Psi}_n \in \mathcal{E}_{\otimes, \sigma}^\infty$ and $\tilde{h}_n \in \mathcal{E}_{\otimes, \sigma'}^\infty$ by (2.6), since $\tilde{\Psi}_n \in C^2(B(d_n \vec{e}_1, R) \cup B(-d_n \vec{e}_1, R), \mathbb{C})$, $\tilde{h}_n \in C^1(B(d_n \vec{e}_1, R) \cup B(-d_n \vec{e}_1, R), \mathbb{C})$ and, outside of $B(d_n \vec{e}_1, R) \cup B(-d_n \vec{e}_1, R)$, $\tilde{\Psi}_n = \Psi_n$ with $\|\Psi_n\|_{*,\sigma,d_n} = 1$, as well as $\tilde{h}_n = h_n$, with $\|h_n\|_{**, \sigma', d_n} \rightarrow 0$ when $n \rightarrow \infty$. In particular,

$$\|\tilde{h}_n\|_{\otimes, \sigma', \infty} = o_{n \rightarrow \infty}^R(1),$$

where $o_{n \rightarrow \infty}^R(1)$ is a sequence that, for fixed $R \geq 4$, goes to 0 when $n \rightarrow \infty$ (it also depends on σ and σ').

Since $\chi_R = 1$ on $\mathbb{R}^2 \setminus (B(d_n \vec{e}_1, R) \cup B(-d_n \vec{e}_1, R))$, we have there $L'(\tilde{\Psi}_n) = \tilde{h}_n$. Therefore, we can write that in \mathbb{R}^2 that $L'(\tilde{\Psi}_n) = \tilde{h}_n + \text{Loc}(\tilde{\Psi}_n)$, with

$$\text{Loc}(\tilde{\Psi}_n) := -\frac{\chi_R \eta}{V} L(V \tilde{\Psi}_n) + (1 - \eta)(L'(\chi_R \tilde{\Psi}_n) - \chi_R L'(\tilde{\Psi}_n)),$$

a term that is supported in $\mathbb{R}^2 \setminus (B(d_n \vec{e}_1, R) \cup B(-d_n \vec{e}_1, R))$. By (2.12) and $\|h_n\|_{**, \sigma', d_n} \rightarrow 0$ when $n \rightarrow \infty$, it satisfies

$$\begin{aligned} & \|\text{Loc}(\Psi_n)\|_{\otimes \otimes, \sigma', \infty} \\ & \leq K(R) \|\text{Loc}(\Psi_n)\|_{C^1(\mathbb{R}^2 \setminus (B(d_n \vec{e}_1, R) \cup B(-d_n \vec{e}_1, R)))} \\ & \leq K(R) \|\Phi_n\|_{C^2(\mathbb{R}^2 \setminus (B(d_n \vec{e}_1, R) \cup B(-d_n \vec{e}_1, R)))} \\ & = o_{n \rightarrow \infty}^R(1). \end{aligned}$$

We recall that $L'(\Psi) = -\Delta \Psi - 2 \frac{\nabla V}{V} \cdot \nabla \Psi + 2|V|^2 \Re \mathfrak{e}(\Psi) - ic \partial_{x_2} \Psi$, therefore

$$-\Delta \tilde{\Psi}_n - ic \partial_{x_2} \tilde{\Psi}_n + 2 \Re \mathfrak{e}(\tilde{\Psi}_n) = \tilde{h}_n + \text{Loc}(\Psi_n) + 2 \frac{\nabla V}{V} \cdot \nabla \tilde{\Psi}_n + 2(1 - |V|^2) \Re \mathfrak{e}(\tilde{\Psi}_n). \quad (2.13)$$

We define

$$\tilde{h}'_n := \tilde{h}_n + \text{Loc}(\Psi_n) + 2 \frac{\nabla V}{V} \cdot \nabla \tilde{\Psi}_n + 2(1 - |V|^2) \Re \mathfrak{e}(\tilde{\Psi}_n).$$

We check that $\tilde{h}'_n \in \mathcal{E}_{\otimes \otimes, \sigma'}^\infty$ with

$$\|\tilde{h}'_n\|_{\otimes \otimes, \sigma', \infty} \leq o_{n \rightarrow \infty}^R(1) + o_{R \rightarrow \infty}(1),$$

where $o_{R \rightarrow \infty}(1)$ is a quantity that goes to 0 when $R \rightarrow \infty$ (in particular, independently of n). By Lemma 2.15, (the condition $\int_{\mathbb{R}^2} \Im \mathfrak{m}(\tilde{h}'_n) = 0$ is a consequence of the symmetries on \tilde{h}_n and $\tilde{\Psi}_n$), this implies, with equation (2.13) (and since $\tilde{\Psi}_n \in \mathcal{E}_{\otimes, \sigma}^\infty$), that

$$\|\tilde{\Psi}_n\|_{\otimes, \sigma, \infty} \leq o_{n \rightarrow \infty}^R(1) + o_{R \rightarrow \infty}(1). \quad (2.14)$$

Step 3. Conclusion.

We have $\|\Psi_n\|_{*, \sigma, d_n} \leq K(R) \|\Phi_n\|_{C^2(\{\tilde{r} \leq R\})} + K \|\tilde{\Psi}_n\|_{*, \sigma, d_n}$ by (2.6), therefore, with equations (2.12) and (2.14),

$$\|\Psi_n\|_{*, \sigma, d_n} \leq o_{n \rightarrow \infty}^R(1) + o_{R \rightarrow \infty}(1).$$

If we take R large enough (depending on σ, σ') so that $o_{R \rightarrow \infty}(1) \leq 1/10$ and then n large enough (depending on R, σ and σ') so that $o_{n \rightarrow \infty}^R(1) \leq 1/10$, we have, for n large, $\|\Psi_n\|_{*, \sigma, d_n} \leq 1/5$, which is in contradiction with

$$\|\Psi_n\|_{*, \sigma, d_n} = 1.$$

□

2.5.2 Existence of a solution

At this point, we do not have existence of a solution to the linear problem

$$\begin{cases} \eta L(\Phi) + (1 - \eta) V L'(\Psi) = V h \\ \Phi \in \mathcal{E}_{*, \sigma}, V h \in \mathcal{E}_{**, \sigma'}, \end{cases}$$

only an a priori estimate. The existence of a solution is done in Proposition 2.20, its proof being the purpose of this subsection. In [8], the existence proof is done using mainly the fact that the domain is bounded. We provide here a proof of existence by approximation on balls of large radii for a particular Hilbertian norm. Given $c > 0$ and $a > 10/c^2$, we define

$$H_a := \left\{ \Phi = Q_c \Psi \in H_{\text{loc}}^1(B(0, a)), \|\Phi\|_{H_a}^2 := \|\Phi\|_{H^1(\{\tilde{r} \leq 3\})}^2 + \int_{\{\tilde{r} \geq 2\} \cap \{r \leq a\}} |\nabla \Psi|^2 + \Re \mathfrak{e}^2(\Psi) + \frac{\Im \mathfrak{m}^2(\Psi)}{(1+r)^{5/2}} \right\},$$

and we also allow $a = +\infty$. We first state a result on functions in H_∞ .

Lemma 2.18 *There exists $c_0 > 0$ such that, for $0 < c < c_0$, $0 < \sigma < \sigma' < 1$, $Vh \in \mathcal{E}_{**, \sigma'}$, if a function $\Phi \in H_\infty \cap C^1(\mathbb{R}^2)$ satisfies, in the weak sense,*

$$\eta L(\Phi) + (1 - \eta)VL'(\Psi) = Vh,$$

and $\Phi = V\Psi$, $\langle V\Psi, Z_d \rangle = 0; \forall x \in \mathbb{R}^2, \Psi(x_1, x_2) = \overline{\Psi(x_1, -x_2)} = \Psi(-x_1, x_2)$, then

$$\Phi \in \mathcal{E}_{*, \sigma}.$$

The next step is to construct a solution on a large ball in the space H_a .

Lemma 2.19 *For $0 < \sigma' < 1$, there exists $c_0(\sigma') > 0$ such that, for $0 < c < c_0(\sigma')$, there exists $a_0(c, \sigma') > \frac{10}{c^2}$ such that, for $Vh \in \mathcal{E}_{**, \sigma'}$, $a > a_0(c, \sigma')$, the problem*

$$\begin{cases} \eta L(\Phi) + (1 - \eta)VL'(\Psi) = Vh & \text{in } B(0, a) \\ \Phi \in H_a, \Phi = V\Psi, \langle V\Psi, Z_d \rangle = 0; \forall x \in B(0, a), \Psi(x_1, x_2) = \overline{\Psi(x_1, -x_2)} = \Psi(-x_1, x_2) \\ \Phi = 0 & \text{on } \partial B(0, a) \\ \langle Vh, Z_d \rangle = 0 \end{cases}$$

admits a unique solution, and furthermore, there exists $K(\sigma', c) > 0$ independent of a such that

$$\|\Phi\|_{H_a} \leq K(\sigma', c)\|h\|_{**, \sigma'}.$$

Here, $a > 10/c^2$ is not necessary, the condition $a > 10/c$ should be enough. However, this simplifies some estimates in the proof, and it will be enough for us here. Here, we require $\langle Vh, Z_d \rangle = 0$ in order to apply the Fredholm alternative in $\{\varphi \in H_0^1(B(0, a)), \langle \varphi, Z_d \rangle = 0\}$ to show the existence of a solution.

Proof We argue by contradiction on the estimation. Assuming the existence, take any $0 < \sigma' < 1$, and choose $c_0(\sigma') > 0$ smaller than the one from Proposition 2.17, and $0 < c < c_0(\sigma')$. Suppose that there exists a sequence $a_n > \frac{10}{c^2}$, $a_n \rightarrow \infty$, functions $\Phi_n \in H_{a_n}$, $\Phi_n = 0$ on $\partial B(0, a_n)$ and $Vh_n \in \mathcal{E}_{**, \sigma'}$ such that $\|\Phi_n\|_{H_{a_n}} = 1$, $\|h_n\|_{**, \sigma'} \rightarrow 0$ and $\eta L(\Phi_n) + (1 - \eta)VL'(\Psi_n) = Vh_n$ on $B(0, a_n)$. In particular, remark here that c is independent of n , only the size of the ball grows. Our goal is to show that $\|\Phi_n\|_{H_{a_n}} = o_{n \rightarrow \infty}^c(1)$, where $o_{n \rightarrow \infty}^c(1)$ is a quantity going to 0 when $n \rightarrow \infty$ at fixed c , which leads to the contradiction.

Following the same arguments as in step 1 of the proof of Proposition 2.17, we check that $\Phi_n \rightarrow \Phi$ in $C_{\text{loc}}^2(\mathbb{R}^2)$ and $\eta L(\Phi) + (1 - \eta)VL'(\Psi) = 0$ in \mathbb{R}^2 . Furthermore, it is easy to check that, since $\|\Phi_n\|_{H_{a_n}} = 1$, we have $\|\Phi\|_{H_\infty} \leq 1$. Then, by Lemma 2.18, since the orthogonality and the symmetries pass at the limit, this implies that $\Phi \in \mathcal{E}_{*, \sigma}$ for any $0 < \sigma < \sigma'$, and therefore, by Proposition 2.17, $\Phi = 0$.

We deduce that $\|\Phi_n\|_{C^2(B(0, 10/c^2))} = o_{n \rightarrow \infty}^c(1)$. Now, we have the system on $\tilde{\Psi}_n = \tilde{\Psi}_1 + i\tilde{\Psi}_2$ (with $\tilde{h}_n = \tilde{h}_1 + i\tilde{h}_2$):

$$\begin{cases} \Delta \tilde{\Psi}_1 - 2\tilde{\Psi}_1 - c\partial_{x_2} \tilde{\Psi}_2 = -\tilde{h}_1 - 2\Re \left(\frac{\nabla V}{V} \cdot \nabla \tilde{\Psi}_n \right) + \text{Loc}_1(\Psi_n) - 2(1 - |V|^2)\tilde{\Psi}_1 \\ \Delta \tilde{\Psi}_2 + c\partial_{x_2} \tilde{\Psi}_1 = -\tilde{h}_2 - 2\Im \left(\frac{\nabla V}{V} \cdot \nabla \tilde{\Psi}_n \right) + \text{Loc}_2(\Psi_n). \end{cases}$$

Now, multiplying the first equation by $\tilde{\Psi}_1$ and integrating on $\Omega = B(0, a) \setminus B(0, 5/c^2)$, we infer by Cauchy-Schwarz

$$\|\nabla \tilde{\Psi}_1\|_{L^2(\Omega)}^2 + 2\|\tilde{\Psi}_1\|_{L^2(\Omega)}^2 \leq$$

$$c\|\nabla \tilde{\Psi}_2\|_{L^2(\Omega)}\|\tilde{\Psi}_1\|_{L^2(\Omega)} + (o_{c \rightarrow 0}(1) + o_{n \rightarrow \infty}^c(1))(\|\nabla \tilde{\Psi}_n\|_{L^2(\Omega)} + \|\tilde{\Psi}_1\|_{L^2(\Omega)}) + o_{n \rightarrow \infty}^c(1).$$

Now, we multiply the second equation by $\tilde{\Psi}_2$, and we integrate on Ω . By integration by parts, we check that

$$\|\nabla \tilde{\Psi}_2\|_{L^2(\Omega)}^2 \leq o_{n \rightarrow \infty}^c(1) + c\|\tilde{\Psi}_1\|_{L^2(\Omega)}\|\nabla \tilde{\Psi}_2\|_{L^2(\Omega)} + o_{c \rightarrow 0}^{\sigma'}(1)\|\nabla \tilde{\Psi}_2\|_{L^2(\Omega)},$$

and by a Hardy-type inequality,

$$\left| \int_{\Omega} \frac{|\tilde{\Psi}_2|^2}{(1 + |x|)^{5/2}} \right| \leq o_{n \rightarrow \infty}^c(1) + o_{c \rightarrow 0}(1).$$

Together with $\|\Phi_n\|_{C^2(B(0,10/c^2))} = o_{n \rightarrow \infty}^c(1)$, this is in contradiction with $\|\Phi_n\|_{H_{a_n}} = 1$.

This concludes the proof of the estimation. Now, for the existence, we argue by Fredholm's alternative in $\{\varphi \in H_0^1(B(0,a)), \langle \varphi, Z_d \rangle = 0\}$, and we remark that the norms $\|\cdot\|_{H_a}$ and $\|\cdot\|_{H^1}$ are equivalent on $B(0,a)$. By Riesz's representation theorem, the elliptic equation $\eta L(\Phi) + (1-\eta)VL'(\Psi) = Vh$ can be rewritten in the operational form $\Phi + \mathcal{K}(\Phi) = \mathcal{S}(h)$ where \mathcal{K} is a compact operator in $H_0^1(B(0,a))$, and it has no kernel in H_a (i.e. in $\{\varphi \in H_0^1(B(0,a)), \langle \varphi, Z_d \rangle = 0\}$) by the estimation we just showed. Therefore, there exists a unique solution $\Phi \in H_a$, and it then satisfies

$$\|\Phi\|_{H_a} \leq K(\sigma', c)\|h\|_{**,\sigma'}.$$

□

Proposition 2.20 *Consider the problem, for $0 < \sigma < \sigma' < 1$,*

$$\begin{cases} \eta L(\Phi) + (1-\eta)VL'(\Psi) = Vh \\ Vh \in \mathcal{E}_{**,\sigma'}, \langle Vh, Z_d \rangle = 0. \end{cases}$$

*Then, there exist constants $c_0(\sigma, \sigma') > 0$ small and $C(\sigma, \sigma') > 0$ depending only on σ, σ' , such that, for $0 < c \leq c_0(\sigma, \sigma')$ and $Vh \in \mathcal{E}_{**,\sigma'}$ with $\langle Vh, Z_d \rangle = 0$, there exists $\Phi \in \mathcal{E}_{*,\sigma}$, $\Phi = V\Psi$ solution of this problem, with*

$$\|\Psi\|_{*,\sigma,d} \leq C(\sigma, \sigma')\|h\|_{**,\sigma',d}.$$

Proof By Lemma 2.19, For $a > a_0(c, \sigma')$, there exists a solution to the problem

$$\begin{cases} \eta L(\Phi_a) + (1-\eta)VL'(\Psi_a) = Vh & \text{on } B(0,a) \\ \Phi_a \in H_a, \Phi_a = V\Psi_a, \langle V\Psi_a, Z_d \rangle = 0; \forall x \in B(0,a), \Psi_a(x_1, x_2) = \overline{\Psi_a(x_1, -x_2)} = \Psi_a(-x_1, x_2) \\ \Phi_a = 0 & \text{on } \partial B(0,a) \\ \langle h, Z_d \rangle = 0 \end{cases}$$

with $\|\Phi_a\|_{H_a} \leq K(\sigma', c)\|h\|_{**,\sigma'}$. Taking a sequence of values $a_n > a_0$ going to infinity, we can construct by a diagonal argument a function $\Phi \in H_{\text{loc}}^1(\mathbb{R}^2)$ which satisfies in the distribution sense

$$\eta L(\Phi) + (1-\eta)VL'(\Phi) = Vh$$

(hence $\Phi \in C^2(\mathbb{R}^2)$ by standard elliptic arguments), such that

$$\|\Phi\|_{H_\infty} \leq \limsup_{n \rightarrow \infty} \|\Phi_n\|_{H_{a_n}} \leq K(\sigma', c)\|h\|_{**,\sigma'},$$

thus $\Phi \in H_\infty$, and $\Phi = V\Psi$, $\langle V\Psi, Z_d \rangle = 0; \forall x \in \mathbb{R}^2, \Psi(x_1, x_2) = \overline{\Psi(x_1, -x_2)} = \Psi(-x_1, x_2)$. From Lemma 2.18, we deduce that $\Phi \in \mathcal{E}_{*,\sigma}$, and is thus a solution to the problem. Furthermore, by Proposition 2.17, $\|\Psi\|_{*,\sigma,d} \leq C(\sigma, \sigma')\|h\|_{**,\sigma',d}$. Still by Proposition 2.17, this solution is unique in $\mathcal{E}_{*,\sigma'}$. □

2.5.3 Estimates for the contraction in the orthogonal space

We showed in Proposition 2.20 that the operator $\eta L(\cdot) + (1-\eta)VL'(\cdot/V)$ is invertible from $\mathcal{E}_{**,\sigma',d} \cap \{\langle \cdot, Z_d \rangle = 0\}$ to $\mathcal{E}_{*,\sigma,d}$. The operator $(\eta L(\cdot) + (1-\eta)VL'(\cdot/V))^{-1}$ is the one that, for a given $Vh \in \mathcal{E}_{**,\sigma',d}$ such that $\langle Vh, Z_d \rangle = 0$, returns the unique function $\Phi = V\Psi \in \mathcal{E}_{*,\sigma,d}$ such that $\eta L(\Phi) + (1-\eta)VL'(\Psi) = Vh$, and this function satisfies the estimate $\|\Psi\|_{*,\sigma,d} \leq C(\sigma, \sigma')\|h\|_{**,\sigma',d}$.

Now, we define (for any $\Phi \in C^0(\mathbb{R}^2, \mathbb{C})$)

$$\Pi_d^\perp(\Phi) := \Phi - \langle \Phi, Z_d \rangle \frac{Z_d}{\|Z_d\|_{L^2(\mathbb{R}^2)}^2},$$

the projection on the orthogonal of Z_d . We want to apply a fixed-point theorem on the functional

$$(\eta L(\cdot) + (1-\eta)VL'(\cdot/V))^{-1}(\Pi_d^\perp(-F(\cdot/V))) : \mathcal{E}_{*,\sigma} \rightarrow \mathcal{E}_{*,\sigma},$$

and for that we need some estimates on the function $\Pi_d^\perp oF(./V) : \mathcal{E}_{*,\sigma} \rightarrow \{Vh \in \mathcal{E}_{**,\sigma'}, \langle Vh, Z_d \rangle = 0\}$. The function F contains the source term $E - ic\partial_{x_2} V$ and nonlinear terms. The source term requires a precise computation (see Lemma 2.22) to show its smallness in the spaces of invertibility. The nonlinear terms will be small if we do the contraction in an area with small Ψ (which is the case since we will do it in the space of function $\Phi = V\Psi \in \mathcal{E}_{*,\sigma}$ such that $\|\Psi\|_{*,\sigma,d} \leq K_0(\sigma, \sigma')c^{1-\sigma'}$ for a well chosen constant $K_0(\sigma, \sigma') > 0$). This subsection is devoted to the proof of the following result.

Proposition 2.21 *For $0 < \sigma < \sigma' < 1$, there exist constants $K_0(\sigma, \sigma'), c_0(\sigma, \sigma') > 0$ depending only on σ, σ' such that for $0 < c < c_0(\sigma, \sigma')$, the function (from $\mathcal{E}_{*,\sigma,d}$ to $\mathcal{E}_{*,\sigma,d}$)*

$$\Phi \mapsto (\eta L(\cdot) + (1 - \eta)V L'(\cdot/V))^{-1}(\Pi_d^\perp(-F(\Phi/V)))$$

is a contraction in the space of functions $\Phi = V\Psi \in \mathcal{E}_{*,\sigma,d}$ such that $\|\Psi\|_{*,\sigma,d} \leq K_0(\sigma, \sigma')c^{1-\sigma'}$. As such, by the contraction mapping theorem, it admits a unique fixed point $\Phi \in \mathcal{E}_{*,\sigma,d}$ in $\{\Phi \in \mathcal{E}_{*,\sigma,d}, \|\Psi\|_{*,\sigma,d} \leq K_0(\sigma, \sigma')c^{1-\sigma'}\}$, and there exists $\lambda(c, d) \in \mathbb{R}$ such that

$$\eta L(\Phi) + (1 - \eta)V L'(\Psi) + F(\Psi) = \lambda(c, d)Z_d$$

in the distribution sense.

We recall that, from the definition of $\mathcal{E}_{*,\sigma,d}$ in subsection 2.3, $\Phi \in \mathcal{E}_{*,\sigma,d}$ implies that $\langle \Phi, Z_d \rangle = 0$, which is the origin of the fact that $\eta L(\Phi) + (1 - \eta)V L'(\Psi) + F(\Psi)$ is not zero, but only proportional to Z_d .

We start with some estimates on the terms contained in $F(\Psi)$. These are done in the following three lemmas.

Lemma 2.22 *For any $0 < \sigma' < 1$, there exists a constant $C_1(\sigma') > 0$ depending only on σ' such that*

$$\left\| \frac{ic\partial_{x_2} V}{V} \right\|_{**,\sigma',d} + \left\| \frac{E}{V} \right\|_{**,\sigma',d} \leq C_1(\sigma')c^{1-\sigma'}.$$

Proof We have defined the norm

$$\|h\|_{**,\sigma',d} = \|Vh\|_{C^1(\{\tilde{r} \leq 3\})} + \|\tilde{r}^{1+\sigma'} h_1\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma'} h_2\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma'} \nabla h\|_{L^\infty(\{\tilde{r} \geq 2\})},$$

thus we separate two areas for the computation: the first one is where $\tilde{r} \leq 3$ which will be easy and then far from the vortices, i.e. in $\{\tilde{r} \geq 2\}$, where the division by V is not a problem.

Step 1. Estimates for E .

In (2.2), we showed that

$$E = (1 - |V_1|^2)(1 - |V_{-1}|^2)V_1 V_{-1} - 2\nabla V_1 \cdot \nabla V_{-1}.$$

Near V_1 , i.e. in $B(d\vec{e}_1, 3)$, we have from Lemma 2.1,

$$\|(1 - |V_{-1}|^2)\|_{C^1(\{r_1 \leq 3\})} \leq Kc^2 \quad \text{and} \quad \|\nabla V_{-1}\|_{C^1(\{r_1 \leq 3\})} \leq Kc,$$

hence

$$\left\| \frac{E}{V} \right\|_{C^1(\{r_1 \leq 3\})} \leq Kc \leq o_{c \rightarrow 0}^{\sigma'}(1)c^{1-\sigma'}, \quad (2.15)$$

where $o_{c \rightarrow 0}^{\sigma'}(1)$ is a quantity that for a fixed $\sigma' > 0$, goes to 0 when $c \rightarrow 0$. By symmetry, the result holds in the area where $\tilde{r} \leq 3$.

We now turn to the estimates for $\tilde{r} \geq 2$. The first term $(1 - |V_1|^2)(1 - |V_{-1}|^2)$ of $\frac{E}{V}$ is real valued. Using the definition of r_1 and r_{-1} from (2.1), in the right half-plane, where $r_1 \leq r_{-1}$ and $r_{-1} \geq d \geq \frac{K}{c}$, we have from Lemma 2.1

$$\|r_1^{1+\sigma'}(1 - |V_1|^2)(1 - |V_{-1}|^2)\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq K \left\| \frac{1}{r_1^{1-\sigma'} r_{-1}^2} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})}$$

and

$$\|r_1^2(1 - |V_1|^2)(1 - |V_{-1}|^2)\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq K.$$

In this area, $\frac{1}{r_{-1}^2} \leq Kc^2$ and $\frac{1}{r_1^{1-\sigma'}} \leq \frac{1}{2^{1-\sigma'}}$, thus

$$\|r_1^{1+\sigma'}(1 - |V_1|^2)(1 - |V_{-1}|^2)\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq K(\sigma')c^2 \leq o_{c \rightarrow 0}^{\sigma'}(1)c^{1-\sigma'}.$$

By symmetry, the same result holds for the other half-plane, hence

$$\|\tilde{r}^{1+\sigma'}(1 - |V_1|^2)(1 - |V_{-1}|^2)\|_{L^\infty(\{\tilde{r} \geq 2\})} \leq o_{c \rightarrow 0}^{\sigma'}(1)c^{1-\sigma'}. \quad (2.16)$$

From Lemma 2.1, we have

$$\nabla V_\varepsilon(x) = i\varepsilon V_\varepsilon(x) \frac{x^\perp}{r^2} + O\left(\frac{1}{r^3}\right),$$

hence

$$\frac{\nabla V_1 \cdot \nabla V_{-1}}{V_1 V_{-1}} = \frac{y^\perp \cdot z^\perp}{r_1^2 r_{-1}^2} + O\left(\frac{1}{r_1^3 r_{-1}}\right) + O\left(\frac{1}{r_{-1}^3 r_1}\right).$$

Remark that the first term is real-valued. We compute first in the right half-plane, where $r_1 \leq r_{-1}$ and $r_{-1} \geq d \geq \frac{K}{c}$,

$$\left\| r_1^{1+\sigma'} \frac{y^\perp \cdot z^\perp}{r_1^2 r_{-1}^2} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq \left\| \frac{r_1^{1+\sigma'}}{r_1 r_{-1}} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})}.$$

Since

$$\frac{r_1^{1+\sigma'}}{r_1 r_{-1}} = \left(\frac{r_1}{r_{-1}}\right)^{\sigma'} \frac{1}{r_{-1}^{1-\sigma'}} \leq K(\sigma')c^{1-\sigma'},$$

we deduce

$$\left\| r_1^{1+\sigma'} \frac{y^\perp \cdot z^\perp}{r_1^2 r_{-1}^2} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq K(\sigma')c^{1-\sigma'}$$

and by symmetry,

$$\left\| \tilde{r}^{1+\sigma'} \frac{y^\perp \cdot z^\perp}{r_1^2 r_{-1}^2} \right\|_{L^\infty(\{\tilde{r} \geq 2\})} \leq K(\sigma')c^{1-\sigma'}. \quad (2.17)$$

For the last two terms $O\left(\frac{1}{r_1^3 r_{-1}}\right) + O\left(\frac{1}{r_{-1}^3 r_1}\right)$, we will show that in the right half-plane

$$\left\| r_1^{2+\sigma'} \frac{1}{r_1^3 r_{-1}} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} + \left\| r_1^{2+\sigma'} \frac{1}{r_{-1}^3 r_1} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq o_{c \rightarrow 0}^{\sigma'}(1)c^{1-\sigma'}. \quad (2.18)$$

This immediately implies

$$\left\| r_1^{1+\sigma'} \frac{1}{r_1^3 r_{-1}} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} + \left\| r_1^{1+\sigma'} \frac{1}{r_{-1}^3 r_1} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq o_{c \rightarrow 0}^{\sigma'}(1)c^{1-\sigma'}. \quad (2.19)$$

We compute in the right half-plane where $r_1 \leq r_{-1}$ and $r_{-1} \geq d \geq \frac{K}{c}$, $\frac{1}{r_{-1}} \leq Kc$ and $\frac{1}{r_1^{1-\sigma'}} \leq K(\sigma')$, thus

$$r_1^{2+\sigma'} \frac{1}{r_1^3 r_{-1}} = \frac{1}{r_1^{1-\sigma'} r_{-1}} \leq Kc \leq o_{c \rightarrow 0}^{\sigma'}(1)c^{1-\sigma'}.$$

Furthermore, still in the right half-plane,

$$r_1^{2+\sigma'} \frac{1}{r_{-1}^3 r_1} = \left(\frac{r_1}{r_{-1}}\right)^{1+\sigma'} \frac{1}{r_{-1}^{2-\sigma'}} \leq K(\sigma')c^{2-\sigma'} \leq o_{c \rightarrow 0}^{\sigma'}(1)c^{1-\sigma'}.$$

Gathering (2.18) to (2.19) and using the symmetry for the left half-plane, we deduce with the previous estimates (2.15), (2.16), (2.17) that

$$\left\| V \left(\frac{E}{V} \right) \right\|_{C^1(\{\tilde{r} \leq 3\})} + \left\| \tilde{r}^{1+\sigma'} \Re \left(\frac{E}{V} \right) \right\|_{L^\infty(\{\tilde{r} \geq 2\})} + \left\| \tilde{r}^{2+\sigma'} \Im \left(\frac{E}{V} \right) \right\|_{L^\infty(\{\tilde{r} \geq 2\})} \leq K(\sigma') c^{1-\sigma'}.$$

Now, for the estimate on $\nabla \left(\frac{E}{V} \right)$ in $\{\tilde{r} \geq 2\}$, we have from Lemma 2.1, for $\tilde{r} \geq 2$,

$$|\nabla((1 - |V_1|^2)(1 - |V_{-1}|^2))| \leq |\nabla|V_1|^2(1 - |V_{-1}|^2)| + |(1 - |V_1|^2)\nabla|V_{-1}|^2| \leq \frac{K}{r_1^3 r_{-1}^2} + \frac{K}{r_1^2 r_{-1}^3},$$

and

$$\left| \nabla \left(\frac{\nabla V_1 \cdot \nabla V_{-1}}{V_1 V_{-1}} \right) \right| \leq \left| \nabla \left(\frac{\nabla V_1}{V_1} \right) \cdot \frac{\nabla V_{-1}}{V_{-1}} \right| + \left| \frac{\nabla V_1}{V_1} \cdot \nabla \left(\frac{\nabla V_{-1}}{V_{-1}} \right) \right| \leq \frac{K}{r_1^2 r_{-1}} + \frac{K}{r_1 r_{-1}^2},$$

thus, with similar estimates as previously, we deduce

$$\left\| \tilde{r}^{2+\sigma'} \nabla \left(\frac{E}{V} \right) \right\|_{L^\infty(\{\tilde{r} \geq 2\})} \leq K(\sigma') c^{1-\sigma'}. \quad (2.20)$$

This concludes the proof of

$$\left\| \frac{E}{V} \right\|_{**,\sigma',d} \leq C'_1(\sigma') c^{1-\sigma'}$$

for some constant $C'_1(\sigma') > 0$ depending only on σ' .

Step 2. Estimates for $ic \frac{\partial_{x_2} V}{V}$.

First, near the vortices, we have $|\partial_{x_2} V| + |\nabla \partial_{x_2} V| \leq K$ a universal constant, therefore

$$\left\| ic \frac{\partial_{x_2} V}{V} V \right\|_{C^1(\{\tilde{r} \leq 3\})} \leq Kc \leq o_{c \rightarrow 0}^{\sigma'}(1) c^{1-\sigma'}.$$

We now turn to the estimate for $\tilde{r} \geq 2$. Recall Lemma 2.5, stating that for a universal constant $C > 0$, since $r_1, r_{-1} \geq 2$,

$$\left| ic \frac{\partial_{x_2} V}{V} - 2cd \frac{x_1^2 - d^2 - x_2^2}{r_1^2 r_{-1}^2} \right| \leq C \left(\frac{c}{r_1^3} + \frac{c}{r_{-1}^3} \right).$$

Remark that $2cd \frac{x_1^2 - d^2 - x_2^2}{r_1^2 r_{-1}^2}$ is real-valued. Using that $cd \leq 2$, that

$$|x_1^2 - d^2| = |(x_1 - d)(x_1 + d)| \leq r_1 r_{-1}$$

and also that $x_2^2 \leq r_1 r_{-1}$, we deduce that in the right half-plane, where $r_1 \leq r_{-1}$ and $r_{-1} \geq d \geq \frac{K}{c}$,

$$\left\| r_1^{1+\sigma'} 2cd \frac{x_1^2 - d^2 - x_2^2}{r_1^2 r_{-1}^2} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq K \left\| \frac{r_1^{1+\sigma'}}{r_1 r_{-1}} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})},$$

and since we have

$$\frac{r_1^{1+\sigma'}}{r_1 r_{-1}} = \left(\frac{r_1}{r_{-1}} \right)^{\sigma'} \frac{1}{r_{-1}^{1-\sigma'}} \leq K(\sigma') c^{1-\sigma'},$$

we infer

$$\left\| 2r_1^{1+\sigma'} cd \frac{x_1^2 - d^2 - x_2^2}{r_1^2 r_{-1}^2} \right\|_{L^\infty(\{2 \leq r_1 \leq r_{-1}\})} \leq K(\sigma') c^{1-\sigma'}.$$

It is easy to check that in the right half-plane

$$r_1^{2+\sigma'} \left(\frac{c}{r_1^3} + \frac{c}{r_{-1}^3} \right) \leq Kc \leq o_{c \rightarrow 0}^{\sigma'}(1) c^{1-\sigma'},$$

and therefore by symmetry for the left half-plane,

$$\begin{aligned} & \left\| V \left(ic \frac{\partial_{x_2} V}{V} \right) \right\|_{C^1(\{\tilde{r} \leq 3\})} + \left\| \tilde{r}^{1+\sigma'} \Re \left(ic \frac{\partial_{x_2} V}{V} \right) \right\|_{L^\infty(\{\tilde{r} \geq 2\})} \\ & + \left\| \tilde{r}^{2+\sigma'} \Im \left(ic \frac{\partial_{x_2} V}{V} \right) \right\|_{L^\infty(\{\tilde{r} \geq 2\})} \\ & \leq K(\sigma') c^{1-\sigma'}. \end{aligned}$$

From we can also check (using Lemma 2.3) that, if $\tilde{r} \geq 1$,

$$\left| \nabla \left(ic \frac{\partial_{x_2} V}{V} - 2cd \frac{x_1^2 - d^2 - x_2^2}{r_1^2 r_{-1}^2} \right) \right| \leq K \left(\frac{c}{r_1^3} + \frac{c}{r_{-1}^3} \right).$$

With $\left| \nabla \left(\frac{1}{r_{\pm 1}} \right) \right| \leq \frac{K}{r_{\pm 1}^2}$ if $\tilde{r} \geq 1$ and similar computations as previously, we check that

$$\left| \nabla \left(2cd \frac{x_1^2 - d^2 - x_2^2}{r_1^2 r_{-1}^2} \right) \right| \leq K(\sigma') c^{1-\sigma'}.$$

Therefore, there exists $C_1''(\sigma') > 0$ such that

$$\left\| ic \frac{\partial_{x_2} V}{V} \right\|_{**, \sigma', d} \leq C_1''(\sigma') c^{1-\sigma'}.$$

We conclude by taking $C_1(\sigma') = \max(C_1'(\sigma'), C_1''(\sigma'))$. \square

Lemma 2.23 For $0 < \sigma < \sigma' < 1$, for $\Phi = V\Psi, \Phi' = V\Psi' \in \mathcal{E}_{*,\sigma,d}$ such that $\|\Psi\|_{*,\sigma,d}, \|\Psi'\|_{*,\sigma,d} \leq C_0$ with C_0 defined in Lemma 2.7, if there exists $K(\sigma, \sigma') > 0$ such that $\|\Psi\|_{*,\sigma,d}, \|\Psi'\|_{*,\sigma,d} \leq K(\sigma, \sigma') c^{1-\sigma'}$, then

$$\left\| \frac{R(\Psi)}{V} \right\|_{**, \sigma', d} \leq o_{c \rightarrow 0}^{\sigma'}(1) c^{1-\sigma'}$$

and

$$\left\| \frac{R(\Psi') - R(\Psi)}{V} \right\|_{**, \sigma', d} \leq o_{c \rightarrow 0}^{\sigma'}(1) \|\Psi' - \Psi\|_{*,\sigma,d},$$

where the $o_{c \rightarrow 0}^{\sigma, \sigma'}(1)$ is a quantity that, for fixed σ and σ' , goes to 0 when $c \rightarrow 0$.

Lemma 2.24 For $0 < \sigma < \sigma' < 1$, for $\Phi = V\Psi, \Phi' = V\Psi' \in \mathcal{E}_{*,\sigma,d}$ such that $\|\Psi\|_{*,\sigma,d}, \|\Psi'\|_{*,\sigma,d} \leq C_0$ with C_0 defined in Lemma 2.7, if there exists $K(\sigma, \sigma') > 0$ such that $\|\Psi\|_{*,\sigma,d}, \|\Psi'\|_{*,\sigma,d} \leq K(\sigma, \sigma') c^{1-\sigma'}$, then

$$\begin{aligned} & \|(1 - \eta)(-\nabla\Psi \cdot \nabla\Psi + |V|^2 S(\Psi))\|_{**, \sigma', d} \leq o_{c \rightarrow 0}^{\sigma, \sigma'}(1) c^{1-\sigma'}, \\ & \|(1 - \eta)(-\nabla\Psi' \cdot \nabla\Psi' + \nabla\Psi \cdot \nabla\Psi + |V|^2 (S(\Psi') - S(\Psi)))\|_{**, \sigma', d} \leq o_{c \rightarrow 0}^{\sigma, \sigma'}(1) \|\Psi' - \Psi\|_{*,\sigma,d}. \end{aligned}$$

Now, we end the proof of Proposition 2.21

Proof (of Proposition 2.21) We take the constants $C(\sigma, \sigma')$ defined in Proposition 2.17 and $C_1(\sigma')$ from Lemma 2.22. We then define $K_0(\sigma, \sigma') := C(\sigma, \sigma')(C_1(\sigma') + 1)$.

To apply the contraction mapping theorem, we need to show that for $\Phi = V\Psi, \Phi' = V\Psi' \in \mathcal{E}_{*,\sigma,d}$ with

$$\|\Psi\|_{*,\sigma,d}, \|\Psi'\|_{*,\sigma,d} \leq K_0(\sigma, \sigma') c^{1-\sigma'},$$

we have for small $c > 0$,

$$\left\| \frac{F(\Psi)}{V} \right\|_{**, \sigma', d} \leq \frac{K_0(\sigma, \sigma')}{C(\sigma, \sigma')} c^{1-\sigma'} \quad (2.21)$$

and

$$\left\| \frac{F(\Psi') - F(\Psi)}{V} \right\|_{**,\sigma',d} \leq o_{c \rightarrow 0}^{\sigma,\sigma'}(1) \|\Psi' - \Psi\|_{*,\sigma,d}. \quad (2.22)$$

If these estimates hold, using Proposition 2.17, we have that the closed ball $B_{\|\cdot\|_{*,\sigma,d}}(0, K_0(\sigma, \sigma')c^{1-\sigma'})$ is stable by $\Phi \mapsto V(\eta L(V) + (1-\eta)V L'(\cdot))^{-1}(\Pi_d^\perp(-F(\Phi/V)))$ and this operator is a contraction in the ball (for c small enough, depending on σ, σ'), hence we can apply the contraction mapping theorem.

From Lemma 2.7, we have

$$F(\Psi) = E - ic\partial_{x_2}V + V(1-\eta)(-\nabla\Psi \cdot \nabla\Psi + |V|^2S(\Psi)) + R(\Psi).$$

By Lemmas 2.22 to 2.24, we have, given that c is small enough (depending only on σ, σ'), that both (2.21) and (2.22) hold. Therefore, defining $c_0(\sigma, \sigma') > 0$ small enough such that all the required conditions on c are satisfied if $c < c_0(\sigma, \sigma')$, we end the proof of Proposition 2.21.

We have therefore constructed a function $\Phi = V\Psi \in \mathcal{E}_{*,\sigma,d}$ such that

$$\Phi = (\eta L(\cdot) + (1-\eta)V L'(\cdot/V))^{-1}(\Pi_d^\perp(-F(\Phi/V))).$$

Therefore, by definition of the operator $(\eta L(\cdot) + (1-\eta)V L'(\cdot/V))^{-1}$, we have, in the distribution sense,

$$\eta L(\Phi) + (1-\eta)V L'(\Psi) = \Pi_d^\perp(-F(\Phi/V)),$$

and thus, there exists $\lambda(c, d) \in \mathbb{R}$ such that

$$\eta L(\Phi) + (1-\eta)V L'(\Psi) + F(\Psi) = \lambda(c, d)Z_d. \quad \square$$

At this point, we have the existence of a function $\Phi = V\Psi \in \mathcal{E}_{*,\sigma,d}$ depending on c, d and a priori σ, σ' , such that $\|\Psi\|_{*,\sigma,d} \leq K(\sigma, \sigma')c^{1-\sigma'}$ and

$$\eta L(\Phi) + (1-\eta)V L'(\Psi) + F(\Psi) = \lambda(c, d)Z_d \quad (2.23)$$

in the distribution sense for some $\lambda(c, d) \in \mathbb{R}$. By using elliptic regularity, we show easily that $\Phi \in C^\infty(\mathbb{R}^2, \mathbb{C})$ and that (2.23) is verified in the strong sense. The goal is now to show that we can take $\lambda(c, d) = 0$ for a good choice of d , but first we need a better estimate on Φ using the parameters σ and σ' . We denote by $\Phi_{\sigma,\sigma'} = V\Psi_{\sigma,\sigma'}$ the solution obtained by Proposition 2.21 for the values $\sigma < \sigma'$.

Corollary 2.25 *For $0 < \sigma_1 < \sigma'_1 < 1$, $0 < \sigma_2 < \sigma'_2 < 1$, there exists $c_0(\sigma_1, \sigma'_1, \sigma_2, \sigma'_2) > 0$ such that for $0 < c < c_0(\sigma_1, \sigma'_1, \sigma_2, \sigma'_2)$, $\Phi_{\sigma_1, \sigma'_1} = V\Psi_{\sigma_1, \sigma'_1} = V\Psi_{\sigma_2, \sigma'_2} = \Phi_{\sigma_2, \sigma'_2}$. We can thus take any values of σ, σ' with $\sigma < \sigma'$ and the estimate*

$$\|\Psi\|_{*,\sigma,d} \leq K(\sigma, \sigma')c^{1-\sigma'}$$

holds for $0 < c < c_0(\sigma, \sigma')$. In particular, for c small enough,

$$\|\Phi\|_{C^2(\{\tilde{r} \leq 3\})} \leq Kc^{3/4}.$$

Proof This is because for $\sigma_1 < \sigma_2$, $\mathcal{E}_{*,\sigma_2} \subset \mathcal{E}_{*,\sigma_1}$ hence the fixed point for σ_2 (for any $\sigma'_2 > \sigma_2$) yields the same value of Ψ as the fixed point for σ_1 for c small enough (for any $\sigma'_1 > \sigma_1$). In particular, this implies also that $\lambda(c, d)$ is independent of σ, σ' (for c small enough). \square

2.6 Estimation on the Lagrange multiplier $\lambda(c, d)$

To finish the construction of a solution of (TW_c) , we need to find a link between d and c such that $\lambda(c, d) = 0$ in (2.23). Here, we give an estimate of $\lambda(c, d)$ for small values of c .

Proposition 2.26 For $\lambda(c, d), \Phi = V\Psi$ defined in the equation of Proposition 2.21, namely

$$\eta L(\Phi) + (1 - \eta)VL'(\Psi) + F(\Psi) = \lambda(c, d)Z_d,$$

we have, for any $0 < \sigma < 1$,

$$\lambda(c, d) \int_{\mathbb{R}^2} |\partial_d V|^2 \eta = \pi \left(\frac{1}{d} - c \right) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

We will take the scalar product of $\eta L(\Phi) + (1 - \eta)VL'(\Psi) + F(\Psi) - \lambda(c, d)Z_d$ with $\partial_d V$. We will show in the proof that in the term $\langle \eta L(\Phi) + (1 - \eta)VL'(\Psi) + F(\Psi), \partial_d V \rangle$, the largest contribution come from the source term $E - ic\partial_{x_2} V$ in $F(\Psi)$. We will show that $\langle E, \partial_d V \rangle \simeq \frac{\pi}{d}$ and $\langle -ic\partial_{x_2} V, \partial_d V \rangle \simeq -\pi c$, so that, at the leading order, $\lambda(c, d) \sim K \left(\frac{1}{d} - c \right)$. In the proof, steps 1, 2 and 7 show that the terms other than $E - ic\partial_{x_2} V$ are of lower order, and steps 3-6 compute exactly the contribution of these leading order terms.

Proof Recall from Lemma 2.7 that $L(\Phi) = (E - ic\partial_{x_2} V)\Psi + VL'(\Psi)$, hence we write the equation under the form

$$L(\Phi) - (1 - \eta)(E - ic\partial_{x_2} V)\Psi + F(\Psi) = \lambda(c, d)Z_d.$$

We want to take the scalar product with $\partial_d V$. We will compute the terms $(1 - \eta)E\Psi$ (step 1), $F(\Psi)$ (steps 2 to 6) and in step 7 we will show that we can do an integration by parts for $\langle L(\Phi), Z_d \rangle$ and compute its contribution.

We have by definition $Z_d = \eta\partial_d V$, hence

$$\langle Z_d, \partial_d V \rangle = \int_{\mathbb{R}^2} |\partial_d V|^2 \eta$$

which is finite and independent of d since $\eta = 0$ outside $\{\tilde{r} \leq 2\}$. Recall that $\|\Psi\|_{*,\sigma} \leq K(\sigma, \sigma')c^{1-\sigma'}$ where

$$\begin{aligned} \|\Psi\|_{*,\sigma} &= \|V\Psi\|_{C^2(\{\tilde{r} \leq 3\})} + \|\tilde{r}^{1+\sigma}\Psi_1\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla\Psi_1\|_{L^\infty(\{\tilde{r} \geq 2\})} \\ &+ \|\tilde{r}^\sigma\Psi_2\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{1+\sigma}\nabla\Psi_2\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla^2\Psi\|_{L^\infty(\{\tilde{r} \geq 2\})}, \end{aligned}$$

which we will heavily use with several values of σ, σ' in the following computations, in particular for $\sigma \in]0, 1[$, the estimate

$$\|\Psi\|_{*,\sigma/2,d} \leq K(\sigma)c^{1-\sigma}.$$

Step 1. We have $\langle (1 - \eta)(E - ic\partial_{x_2} V)\Psi, \partial_d V \rangle = O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$.

From Lemma 2.6, we have

$$|\partial_d V| \leq \frac{K}{1 + \tilde{r}}. \quad (2.24)$$

In (2.2), we showed that

$$E = -2\nabla V_1 \cdot \nabla V_{-1} + (1 - |V_1|^2)(1 - |V_{-1}|^2)V_1 V_{-1},$$

hence, with Lemmas 2.1 and 2.5 (estimating $ic\partial_{x_2} V$ as in step 2 of the proof of Lemma 2.22), we have

$$|E - ic\partial_{x_2} V| \leq \frac{Kc}{1 + \tilde{r}}$$

by using $|\nabla V_1| \leq \frac{K}{1 + \tilde{r}}$, $|\nabla V_{-1}| \leq \frac{K}{d} \leq Kc$ and $|1 - |V_{-1}|^2| \leq Kc^2$ in the right half-plane and the symmetric estimate in the other one. We also have, in $\{1 - \eta \neq 0\}$,

$$|\Psi| \leq K \frac{\|\Psi\|_{*,\sigma/2,d}}{(1 + \tilde{r})^{\sigma/2}} \leq \frac{K(\sigma)c^{1-\sigma}}{(1 + \tilde{r})^{\sigma/2}},$$

hence

$$|\langle (1 - \eta)(E - ic\partial_{x_2} V)\Psi, \partial_d V \rangle| \leq K(\sigma) \int_{\mathbb{R}^2} \frac{c^{2-\sigma}}{(1 + \tilde{r})^{2+\sigma/2}} = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Step 2. We have $\langle F(\Psi), \partial_d V \rangle = \langle E - ic\partial_{x_2} V, \partial_d V \rangle + O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$.

In this step, we want to show that the nonlinear terms in $F(\Psi)$ are negligible. Recall that

$$F(\Psi) = E - ic\partial_{x_2} V + R(\Psi) + V(1 - \eta)(-\nabla\Psi \cdot \nabla\Psi + |V|^2 S(\Psi)).$$

We first show that

$$\langle R(\Psi), \partial_d V \rangle = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Indeed, $R(\Psi)$ is localized in $\{\tilde{r} \leq 2\}$ and $|R(\Psi)| \leq C\|\Phi\|_{C^1(\{\tilde{r} \leq 3\})}^2$ (since $\|\Psi\|_{*,\sigma,d} \leq C_0$, see Lemma 2.7), and using that in $\{\tilde{r} \leq 3\}$, $|\Phi| + |\nabla\Phi| \leq K(\sigma)c^{1-\sigma/2}$ yields

$$|R(\Psi)| \leq c\|\partial_{x_2}\Phi\|_{C^0(\{\tilde{r} \leq 3\})} + C\|\Phi\|_{C^1(\{\tilde{r} \leq 3\})}^2 = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Now, we use $\|\Psi\|_{*,\sigma/2,d} \leq K(\sigma)c^{1-\sigma}$ to estimate, in $\{1 - \eta \neq 0\}$,

$$|\nabla\Psi \cdot \nabla\Psi| \leq \frac{K\|\Psi\|_{*,\sigma,d}^2}{(1 + \tilde{r})^{2+\sigma}} \leq \frac{K(\sigma)c^{2-\sigma}}{(1 + \tilde{r})^{2+\sigma}},$$

therefore

$$|\langle -\nabla\Psi \cdot \nabla\Psi V(1 - \eta), \partial_d V \rangle| \leq Kc^{2-\sigma} \int_{\mathbb{R}^2} \frac{1}{(1 + \tilde{r})^{3+\sigma}} = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

The same argument can be made for

$$|\langle -|V|^2 S(\Psi) V(1 - \eta), \partial_d V \rangle| = O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$$

by using $S(\Psi) = e^{2\Re(\Psi)} - 1 - 2\Re(\Psi)$ and the fact that it is real-valued.

Step 3. We have $\langle E - ic\partial_{x_2} V, \partial_d V \rangle = -2 \int_{\{x_1 \geq 0\}} \Re((E - ic\partial_{x_2} V) \overline{\partial_{x_1} V_1 V_{-1}}) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$.

The goal of this step is to simplify the computation by using the symmetry. By symmetry, we can only look in the right half-plane:

$$\langle E - ic\partial_{x_2} V, \partial_d V \rangle = 2 \int_{\{x_1 \geq 0\}} \Re((E - ic\partial_{x_2} V) \overline{\partial_d V}).$$

Recall that $\partial_d V = -\partial_{x_1} V_1 V_{-1} + \partial_{x_1} V_{-1} V_1$, hence we need to show that

$$\int_{\{x_1 \geq 0\}} \Re((E - ic\partial_{x_2} V) \overline{\partial_{x_1} V_{-1} V_1}) = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

We compute

$$\begin{aligned} \int_{\{x_1 \geq 0\}} \Re((E - ic\partial_{x_2} V) \overline{\partial_{x_1} V_{-1} V_1}) &= \int_{\{x_1 \geq 0\}} \Re\left(\left(\frac{E - ic\partial_{x_2} V}{V} |V|^2\right) \overline{\frac{\partial_{x_1} V_{-1}}{V_{-1}}}\right) \\ &= \int_{\{x_1 \geq 0\}} \Re\left(\frac{E - ic\partial_{x_2} V}{V} |V|^2\right) \Re\left(\frac{\partial_{x_1} V_{-1}}{V_{-1}}\right) \\ &\quad + \int_{\{x_1 \geq 0\}} \Im\left(\frac{E - ic\partial_{x_2} V}{V} |V|^2\right) \Im\left(\frac{\partial_{x_1} V_{-1}}{V_{-1}}\right). \end{aligned}$$

In the right half-plane, we have $d \leq r_{-1}$ and $\tilde{r} \leq r_1$, hence

$$\begin{aligned} \left| \Re\left(\frac{\partial_{x_1} V_{-1}}{V_{-1}}\right) \right| &\leq \frac{K}{r_{-1}^3} \leq \frac{Kc^{1-\sigma/2}}{(1 + \tilde{r})^{2+\sigma/2}}, \\ \left| \Im\left(\frac{\partial_{x_1} V_{-1}}{V_{-1}}\right) \right| &\leq \frac{K}{r_{-1}} \leq \frac{Kc^{1-\sigma/2}}{(1 + \tilde{r})^{\sigma/2}}, \end{aligned}$$

from Lemma 2.1. Moreover,

$$\left| \Re \left(\frac{E - ic\partial_{x_2} V}{V} |V|^2 \right) \right| \leq \frac{Kc^{1-\sigma/2}}{(1+\tilde{r})^{1+\sigma/2}},$$

$$\left| \Im \left(\frac{E - ic\partial_{x_2} V}{V} |V|^2 \right) \right| \leq \frac{Kc^{1-\sigma/2}}{(1+\tilde{r})^{2+\sigma/2}},$$

from Lemma 2.22. We thus deduce that

$$\left| \int_{\{x_1 \geq 0\}} \Re \left((E - ic\partial_{x_2} V) \overline{\partial_{x_1} V_{-1} V_{-1}} \right) \right| \leq Kc^{1-\sigma/2} \int_{\mathbb{R}^2} \frac{c^{1-\sigma/2}}{(1+\tilde{r})^{2+\sigma}} = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Step 4. We have

$$\int_{\{x_1 \geq 0\}} \Re \left(E \overline{\partial_{x_1} V_{-1} V_{-1}} \right) = -2 \int_{\{x_1 \geq 0\}} \Re \left(\partial_{x_2} V_{-1} \overline{\partial_{x_1} V_{-1} \partial_{x_2} V_{-1} V_{-1}} \right) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

The goal of this step is to compute the part of E that produces the higher order term. Recall from (2.2) that

$$E = -2\nabla V_1 \cdot \nabla V_{-1} + (1 - |V_1|^2)(1 - |V_{-1}|^2)V_1 V_{-1}$$

and since

$$|(1 - |V_1|^2)(1 - |V_{-1}|^2)| \leq \frac{Kc^2}{(1+\tilde{r})^2}$$

by Lemma 2.1, we deduce

$$\int_{\{x_1 \geq 0\}} \Re \left((1 - |V_1|^2)(1 - |V_{-1}|^2) V_1 V_{-1} \overline{\partial_{x_1} V_{-1} V_{-1}} \right) = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Now we compute the first contribution from $-2\nabla V_1 \cdot \nabla V_{-1} = -2\partial_{x_1} V_1 \partial_{x_1} V_{-1} - 2\partial_{x_2} V_1 \partial_{x_2} V_{-1}$,

$$\int_{\{x_1 \geq 0\}} \Re \left((-2\partial_{x_1} V_1 \partial_{x_1} V_{-1}) \overline{\partial_{x_1} V_{-1} V_{-1}} \right) = -2 \int_{\{x_1 \geq 0\}} |\partial_{x_1} V_1|^2 \Re \left(\partial_{x_1} V_{-1} \overline{V_{-1}} \right).$$

From Lemma 2.1 we have

$$\Re \left(\partial_{x_1} V_{-1} \overline{V_{-1}} \right) = O \left(\frac{1}{r_{-1}^3} \right)$$

since the main part in $\partial_{x_1} V_{-1} \overline{V_{-1}}$ is purely imaginary. Using $r_1 \leq r_{-1}$ and $r_{-1} \geq d \geq \frac{K}{c}$ in the right half-plane, we have $\frac{1}{r_{-1}^3} \leq \frac{Kc^{2-\sigma}}{(1+\tilde{r})^{1+\sigma}}$ and, noting that $|\partial_{x_1} V_1|^2 \leq \frac{K}{(1+\tilde{r})^2}$, we obtain

$$\int_{\{x_1 \geq 0\}} |\partial_{x_1} V_1|^2 |\Re \left(\partial_{x_1} V_{-1} \overline{V_{-1}} \right)| \leq Kc^{2-\sigma} \int_{\{x_1 \geq 0\}} \frac{1}{(1+\tilde{r})^{3+\sigma}} = O_{c \rightarrow 0}(c^{5/4}).$$

Finally, the second contribution from $-2\nabla V_1 \cdot \nabla V_{-1}$ is

$$\int_{\{x_1 \geq 0\}} \Re \left((-2\partial_{x_2} V_1 \partial_{x_2} V_{-1}) \overline{\partial_{x_1} V_{-1} V_{-1}} \right) = -2 \int_{\{x_1 \geq 0\}} \Re \left(\partial_{x_2} V_1 \overline{\partial_{x_1} V_{-1} \partial_{x_2} V_{-1} V_{-1}} \right)$$

which concludes the proof of this step.

Step 5. We have $\int_{\{x_1 \geq 0\}} \Re \left(E \overline{\partial_{x_1} V_{-1} V_{-1}} \right) = \frac{\pi}{d} + O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$.

By Lemma 2.1, we have

$$\partial_{x_2} V_{-1} \overline{V_{-1}} = -i |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} + O\left(\frac{1}{r_{-1}^3}\right).$$

The $O\left(\frac{1}{r_{-1}^3}\right)$ yielding a term which is a $O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$ as in step 4, therefore

$$\int_{\{x_1 \geq 0\}} \Re \left((-2\partial_{x_2} V_1 \partial_{x_2} \overline{V_{-1}}) \overline{\partial_{x_1} V_1 V_{-1}} \right) = 2 \int_{\{x_1 \geq 0\}} \Re \left(i \partial_{x_2} V_1 \overline{\partial_{x_1} V_1} \right) |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Now we compute in polar coordinate around $d\vec{e}_1^\rightarrow$, writing $V_1 = \rho_1(r_1)e^{i\theta_1}$. From Lemma 2.2, we have

$$\begin{aligned} \partial_{x_1} V_1 &= \left(\cos(\theta_1) \frac{\rho_1'(r_1)}{\rho_1(r_1)} - \frac{i}{r_1} \sin(\theta_1) \right) V_1, \\ \partial_{x_2} V_1 &= \left(\sin(\theta_1) \frac{\rho_1'(r_1)}{\rho_1(r_1)} + \frac{i}{r_1} \cos(\theta_1) \right) V_1. \end{aligned}$$

We then compute

$$\Re \left(i \partial_{x_2} V_1 \overline{\partial_{x_1} V_1} \right) = -|V_1|^2 \left(\cos^2(\theta_1) \frac{\rho_1'}{r_1 \rho_1} + \sin^2(\theta_1) \frac{\rho_1'}{r_1 \rho_1} \right) = -|V_1|^2 \frac{\rho_1'}{r_1 \rho_1}.$$

From Lemma 2.1, we have $\rho_1'(r_1) = O_{r_1 \rightarrow \infty}\left(\frac{1}{r_1^3}\right)$. As a consequence

$$\begin{aligned} & \left| \int_{\{x_1 \geq 0\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} - \int_{\{r_1 \leq d^{1/2}\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} \right| \\ & \leq K c^{2-\sigma} \int_{\{r_1 \geq d^{1/2}\}} \frac{1}{(1+\tilde{r})^{2+2\sigma}} \end{aligned}$$

because when $x_1 \geq 0$ and $r_1 \geq d^{1/2}$, we have $\left| |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} \right| \leq \frac{K c^{2-\sigma}}{(1+\tilde{r})^{2+2\sigma}}$. We deduce that

$$\int_{\{x_1 \geq 0\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} = \int_{\{r_1 \leq d^{1/2}\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

In the ball $\{r_1 \leq d^{1/2}\}$, we have

$$r_{-1}^2 = 4d^2 \left(1 + O_{d \rightarrow \infty} \left(\frac{1}{d} \right) \right) \quad \text{and} \quad |V_{-1}|^2 = 1 + O\left(\frac{1}{d^2}\right)$$

therefore

$$\int_{\{x_1 \geq 0\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} = \frac{1}{4d^2} \int_{\{r_1 \leq d^{1/2}\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} (y_1 + 2d) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Since $y_1 = r_1 \cos(\theta_1)$, by integration in polar coordinates we have

$$\int_{\{r_1 \leq d^{1/2}\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} y_1 = 0$$

hence

$$\int_{\{x_1 \geq 0\}} \Re \left(E \overline{\partial_{x_1} V_1 V_{-1}} \right) = \frac{1}{d} \int_{\{r_1 \leq d^{1/2}\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Remark that $|V_1|^2 = \rho_1^2$ hence

$$\int_{\{r_1 \leq d^{1/2}\}} |V_1|^2 \frac{\rho_1'}{r_1 \rho_1} = 2\pi \int_0^{d^{1/2}} \rho_1 \rho_1' dr_1 = \pi [\rho_1^2]_0^{d^{1/2}} = \pi + O_{d \rightarrow \infty} \left(\frac{1}{d} \right)$$

Since $\rho_1 = 1 + O\left(\frac{1}{r_1^2}\right)$ when $r_1 \rightarrow \infty$ and $\rho_1(0) = 0$ by Lemma 2.1. Therefore, as claimed,

$$\int_{\{x_1 \geq 0\}} \Re \left(E \overline{\partial_{x_1} V_1 V_{-1}} \right) = \frac{\pi}{d} + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Notice that we have shown in particular that

$$\int_{\mathbb{R}^2} \Re \left(i \partial_{x_2} V_1 \overline{\partial_{x_1} V_1} \right) |V_{-1}|^2 = -\pi + O_{c \rightarrow 0}^\sigma(c^{1-\sigma}). \quad (2.25)$$

Step 6. We have $\int_{\{x_1 \geq 0\}} \Re \left(-ic \partial_{x_2} V \overline{\partial_{x_1} V_1 V_{-1}} \right) = -\pi c + O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$.

We are left with the computation of

$$\begin{aligned} & \int_{\{x_1 \geq 0\}} \Re \left(-ic \partial_{x_2} V \overline{\partial_{x_1} V_1 V_{-1}} \right) = \\ & \int_{\{x_1 \geq 0\}} \Re \left(-ic \partial_{x_2} V_1 \overline{\partial_{x_1} V_1} \right) |V_{-1}|^2 + \int_{\{x_1 \geq 0\}} \Re \left(-ic \partial_{x_2} V_{-1} V_1 \overline{\partial_{x_1} V_1 V_{-1}} \right) \end{aligned} \quad (2.26)$$

since $\partial_{x_2} V = \partial_{x_2} V_1 V_{-1} + \partial_{x_2} V_{-1} V_1$. For the second term in (2.26), we compute

$$-c \int_{\{x_1 \geq 0\}} \Re \left(i \partial_{x_2} V_{-1} V_1 \overline{\partial_{x_1} V_1 V_{-1}} \right) = c \int_{\{x_1 \geq 0\}} \Re \left(\overline{\partial_{x_1} V_1 V_1} \right) |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} + O_{c \rightarrow 0}^\sigma(c^{2-s})$$

in view of the relation

$$i \partial_{x_2} V_{-1} \overline{V_{-1}} = -|V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} + O\left(\frac{1}{r_{-1}^3}\right)$$

from Lemma 2.1 and the fact that $\int_{\{x_1 \geq 0\}} c O\left(\frac{1}{r_{-1}^3}\right) = O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$ (as in step 4). Now recall from Lemma 2.2 that

$$\partial_{x_1} V_1 = \left(\cos(\theta_1) \frac{\rho_1'(r_1)}{\rho_1(r_1)} - \frac{i}{r_1} \sin(\theta_1) \right) V_1$$

therefore

$$\Re \left(\overline{\partial_{x_1} V_1 V_1} \right) = \cos(\theta_1) \frac{\rho_1'}{\rho_1} |V_1|^2.$$

In particular, $|\Re \left(\overline{\partial_{x_1} V_1 V_1} \right)| \leq \frac{K}{1+r_1^3}$ is integrable. Furthermore, $\left| |V_{-1}|^2 \frac{y_1 + 2d}{r_{-1}^2} \right| = O_{c \rightarrow 0}(c)$ in the right half-plane, therefore

$$-c \int_{\{x_1 \geq 0\}} \Re \left(i \partial_{x_2} V_{-1} V_1 \overline{\partial_{x_1} V_1 V_{-1}} \right) = O_{c \rightarrow 0}(c^2) = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

The first contribution in (2.26) is

$$c \int_{\{x_1 \geq 0\}} \Re \left(i \partial_{x_2} V_1 \overline{\partial_{x_1} V_1} \right) |V_{-1}|^2 = c \int_{\{x_1 \geq 0\}} \Re \left(i \partial_{x_2} V_1 \overline{\partial_{x_1} V_1} \right) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$$

using that $|V_{-1}|^2 = 1 + O\left(\frac{1}{r_{-1}^2}\right)$. From (2.25), we have

$$\int_{\{x_1 \geq 0\}} \Re \left(i \partial_{x_2} V_1 \overline{\partial_{x_1} V_1} \right) = -\pi + O_{c \rightarrow 0}^\sigma(c^{1-\sigma}).$$

This concludes the proof of step 6, and combining step 4, 5 and 6 we deduce

$$\int_{\{x_1 \geq 0\}} \Re \left((E - ic \partial_{x_2} V) \overline{\partial_{x_1} V_1 V_{-1}} \right) = \pi \left(\frac{1}{d} - c \right) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Step 7. We have $\langle L(\Phi), \partial_d V \rangle = O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$.

We want to compute, by integration by parts, that

$$\langle L(\Phi), \partial_d V \rangle = \langle \Phi, L(\partial_d V) \rangle.$$

First, we recall that the left hand side is well defined, because we showed in the previous steps that all the other terms are bounded, therefore this one is also bounded. We have

$$\int_{B(0,R)} \Re(\Delta \Phi \overline{\partial_d V}) = \int_{\partial B(0,R)} \Re(\nabla \Phi \overline{\partial_d V}) \cdot \vec{n} - \Re(\Phi \overline{\nabla \partial_d V}) \cdot \vec{n} + \int_{B(0,R)} \Re(\Phi \Delta \overline{\partial_d V}),$$

and

$$|\Re(\nabla \Phi \overline{\partial_d V})| + |\Re(\Phi \overline{\nabla \partial_d V})| \leq \frac{K}{(1+\tilde{r})^{2+1/2}},$$

therefore

$$\int_{\partial B(0,R)} \Re(\nabla \Phi \overline{\partial_d V}) \cdot \vec{n} - \Re(\Phi \overline{\nabla \partial_d V}) \cdot \vec{n} = o_{R \rightarrow \infty}(1)$$

and the integration by parts holds.

Recall that

$$L(h) = -\Delta h - (1 - |V|^2)h + 2\Re(\bar{V}h)V - ic\partial_{x_2}h$$

and

$$L_{V_1}(h) = -\Delta h - (1 - |V_1|^2)h + 2\Re(\bar{V}_1 h)V_1.$$

From Lemma 2.6 and $\|\Psi\|_{*,\sigma/2} \leq K(\sigma)c^{1-\sigma}$, we check easily that

$$|\langle \Phi, -ic\partial_{x_2}\partial_d V \rangle| \leq \int_{\mathbb{R}^2} \frac{K(\sigma)c^{2-\sigma}}{(1+\tilde{r})^{2+\sigma/2}} = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

We therefore focus on the remaining part, with the operator

$$\tilde{L}(h) := -\Delta h - (1 - |V|^2)h + 2\Re(\bar{V}h)V - ic\partial_{x_2}h.$$

We remark that we have $L_{V_1}(\partial_{x_1}V_1) = 0$, since $\partial_{x_1}(-\Delta V_1 - (1 - |V_1|^2)V_1) = 0$. Recall that $\partial_d V = -\partial_{x_1}V_1V_{-1} + \partial_{x_1}V_{-1}V_1$ and let us compute

$$\tilde{L}(V_{-1}h) = L_{V_1}(h)V_{-1} - \Delta(V_{-1}h) + \Delta hV_{-1} + (|V|^2 - |V_1|^2)hV_{-1} + 2\Re(\bar{V}_1 h)(1 - |V_{-1}|^2)V,$$

therefore, using the equation on V_{-1} ,

$$\tilde{L}(V_{-1}h) = L_{V_1}(h)V_{-1} - 2\nabla V_{-1} \cdot \nabla h + (1 - |V_{-1}|^2)(1 - |V_1|^2)V_{-1}h + 2\Re(\bar{V}_1 h)(1 - |V_{-1}|^2)V.$$

Taking $h = \partial_{x_1}V_1$ then yields

$$\tilde{L}(V_{-1}\partial_{x_1}V_1) = -2\nabla V_{-1} \cdot \nabla \partial_{x_1}V_1 + (1 - |V_{-1}|^2)(1 - |V_1|^2)V_{-1}\partial_{x_1}V_1 + 2\Re(\bar{V}_1 \partial_{x_1}V_1)(1 - |V_{-1}|^2)V.$$

Remark that $|\nabla V_{-1} \cdot \nabla \partial_{x_1}V_1| \leq \frac{K}{(1+r_1)(1+r_{-1})^2}$, $|(1 - |V_{-1}|^2)(1 - |V_1|^2)V_{-1}\partial_{x_1}V_1| \leq \frac{K}{(1+r_1)^3(1+r_{-1})^2}$ and $|2\Re(\bar{V}_1 \partial_{x_1}V_1)(1 - |V_{-1}|^2)V| \leq \frac{K}{(1+r_1)^3(1+r_{-1})^2}$ for a universal constant $K > 0$ by Lemma 2.1, therefore

$$\langle \Phi, \tilde{L}(\partial_{x_1}V_1V_{-1}) \rangle = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Exchanging the roles of V_1 and V_{-1} , we have similarly

$$\tilde{L}(V_1\partial_{x_1}V_{-1}) = -2\nabla V_1 \cdot \nabla \partial_{x_1}V_{-1} + (1 - |V_{-1}|^2)(1 - |V_1|^2)V_1\partial_{x_1}V_{-1}.$$

We then conclude that

$$\langle \tilde{L}(\Phi), \partial_d V \rangle = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}),$$

which end the proof of this step. Notice that we have shown

$$|L(\partial_d V)| \leq \frac{Kc}{(1+\tilde{r})^2} \quad (2.27)$$

because $\frac{1}{(1+r_1)(1+r_{-1})} \leq \frac{Kc}{(1+\tilde{r})}$ in the whole space.

Step 8. Conclusion.

Adding all the results obtained in steps 1 to 7, we deduce

$$\lambda(c, d) \int_{\mathbb{R}^2} |\partial_d V|^2 \eta = \pi \left(\frac{1}{d} - c \right) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

□

At this point, we cannot conclude that there exists d such that $\lambda(c, d) = 0$. For that, we need to show that the $O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$ is continuous with respect to c and d . This will be shown in section 3.

3 Construction and properties of the travelling wave

Given $0 < \sigma < \sigma' < 1$, $d, c > 0$ satisfying $\frac{1}{2c} < d < \frac{2}{c}$ and $c < c_0(\sigma, \sigma')$ defined in Proposition 2.21, we define $\Phi_{c,d} = V\Psi_{c,d} \in \mathcal{E}_{*,\sigma,d}$ the function constructed by the contraction mapping theorem in Proposition 2.21. From Corollary 2.25, for any $0 < \sigma < \sigma' < 1$, this function satisfies, for $c < c_0(\sigma, \sigma')$, that

$$\|\Psi_{c,d}\|_{*,\sigma,d} \leq K(\sigma, \sigma')c^{1-\sigma'}.$$

With equation (2.23) and Proposition 2.26, if we show that $\Phi_{c,d}$ is a continuous function of c and d , then there exists $c_0 > 0$ such that, for any $0 < c < c_0$, by the intermediate value theorem, there exists $d_c > 0$ such that $\lambda(c, d_c) = 0$. This would conclude the construction of the travelling wave, and is done in subsection 3.1. In subsection 3.2, we compute some estimates on Q_c which will be useful for understanding the linearized operator around Q_c . We also show there that Q_c is a travelling wave solution with finite energy.

3.1 Proof that $\Phi_{c,d}$ is a C^1 function of c and d

To end the construction of the travelling wave, we only need the continuity of $\Phi_{c,d}$ with respect to c and d . But for the construction of the C^1 branch of travelling wave in section 4, we need its differentiability.

3.1.1 Setup of the problem

From Proposition 2.21, the function $\Phi_{c,d}$ is defined by the implicit equation on $\mathcal{E}_{*,\sigma,d}$

$$(\eta L(\cdot) + (1-\eta)V L'(\cdot/V))^{-1}(\Pi_d^\perp(-F(\Phi_{c,d}/V))) + \Phi_{c,d} = 0,$$

where $(\eta L(\cdot) + (1-\eta)V L'(\cdot/V))^{-1}$ is the linear operator from $\mathcal{E}_{**,\sigma',d} \cap \{\langle \cdot, Z_d \rangle = 0\}$ to $\mathcal{E}_{*,\sigma,d}$, that, for a function $Vh \in \mathcal{E}_{**,\sigma',d}$ with $\langle Vh, Z_d \rangle = 0$, yields the unique function $\Phi = V\Psi \in \mathcal{E}_{*,\sigma,d}$ such that

$$\eta L(\Phi) + (1-\eta)V L'(\Psi) = Vh$$

in the distribution sense. We recall the quantity $Z_d(x) = \partial_d V(x)(\tilde{\eta}(4r_1) + \tilde{\eta}(4r_{-1}))$ defined in subsection 2.3 and we have defined the projection

$$\Pi_d^\perp(\Phi) = \Phi - \langle \Phi, Z_d \rangle \frac{Z_d}{\|Z_d\|_{L^2(\mathbb{R}^2)}^2}.$$

We want to show that $(c, d) \mapsto \Phi_{c,d}$ is of class C^1 from values of c, d such that $0 < c < c_0(\sigma)$ and $\frac{1}{2d} < c < \frac{2}{d}$ to $\mathcal{E}_{*,\sigma,d}$. The first obstacle is that $\mathcal{E}_{*,\sigma,d}$ depends on d (through \tilde{r}), both in the weights in $\|\cdot\|_{*,\sigma,d}$ and in the

orthogonality required: $\langle \Phi, Z_d \rangle = 0$. To be able to use the implicit function theorem, we first need to write an equation on Φ in a space that does not depend on d . The norm $\|\cdot\|_{*,\sigma,d}$ depends on d (through \tilde{r}):

$$\begin{aligned} \|\Psi\|_{*,\sigma,d} &= \|V\Psi\|_{C^2(\{\tilde{r}\leq 3\})} + \|\tilde{r}^{1+\sigma}\Psi_1\|_{L^\infty(\{\tilde{r}\geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla\Psi_1\|_{L^\infty(\{\tilde{r}\geq 2\})} \\ &+ \|\tilde{r}^\sigma\Psi_2\|_{L^\infty(\{\tilde{r}\geq 2\})} + \|\tilde{r}^{1+\sigma}\nabla\Psi_2\|_{L^\infty(\{\tilde{r}\geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla^2\Psi\|_{L^\infty(\{\tilde{r}\geq 2\})}. \end{aligned}$$

For $d_\otimes \in \mathbb{R}, d_\otimes \geq 10$ and $d \in \mathbb{R}$ such that $|d - d_\otimes| < \delta$ for some small $\delta > 0$ (that we will fix later on), we define

$$\begin{aligned} \|\Phi\|_{\otimes,\sigma,d_\otimes} &:= \|\Phi\|_{C^2(\{\tilde{r}_\otimes\leq 3\})} + \left\| \tilde{r}_\otimes^{1+\sigma}\Re\left(\frac{\Phi}{V_\otimes}\right) \right\|_{L^\infty(\{\tilde{r}_\otimes\geq 2\})} + \left\| \tilde{r}_\otimes^{2+\sigma}\nabla\Re\left(\frac{\Phi}{V_\otimes}\right) \right\|_{L^\infty(\{\tilde{r}_\otimes\geq 2\})} \\ &+ \left\| \tilde{r}_\otimes^\sigma\Im\left(\frac{\Phi}{V_\otimes}\right) \right\|_{L^\infty(\{\tilde{r}_\otimes\geq 2\})} + \left\| \tilde{r}_\otimes^{1+\sigma}\nabla\Im\left(\frac{\Phi}{V_\otimes}\right) \right\|_{L^\infty(\{\tilde{r}_\otimes\geq 2\})} \\ &+ \left\| \tilde{r}_\otimes^{2+\sigma}\nabla^2\left(\frac{\Phi}{V_\otimes}\right) \right\|_{L^\infty(\{\tilde{r}_\otimes\geq 2\})}, \end{aligned}$$

where $V_\otimes = V_1(x - d_\otimes\vec{e}_1)V_{-1}(x + d_\otimes\vec{e}_1)$ and $\tilde{r}_\otimes = \min(r_{1,\otimes}, r_{-1,\otimes})$ with $r_{1,\otimes} = |x - d_\otimes\vec{e}_1|, r_{-1,\otimes} = |x + d_\otimes\vec{e}_1|$. Then, for $\Phi = V\Psi \in \mathcal{E}_{*,\sigma,d}$ (V taken in d),

$$K_1\|\Psi\|_{*,\sigma,d} \leq \|\Phi\|_{\otimes,\sigma,d_\otimes} \leq K_2\|\Psi\|_{*,\sigma,d} \quad (3.1)$$

where $K_{1,2} > 0$ are absolute when $|d - d_\otimes| < \delta$. Indeed, we check with simple geometric arguments that if $\tilde{r}_\otimes \geq 1$, V taken in d , then $\tilde{r} \geq 1/2$ and we have

$$\left| \frac{V}{V_\otimes} - 1 \right| \leq \frac{K}{(1+\tilde{r})} \quad \text{and} \quad \left| \nabla\left(\frac{V}{V_\otimes}\right) \right| \leq \frac{K}{(1+\tilde{r})^2} \quad (3.2)$$

for a universal constant $K > 0$. Moreover, we have, for instance, if $\tilde{r}_\otimes \geq 2$ (hence $\tilde{r}_\otimes \leq 2\tilde{r}$),

$$\begin{aligned} \left| \tilde{r}_\otimes^{1+\sigma}\Re\left(\frac{\Phi}{V_\otimes}\right) \right| &\leq \left| \tilde{r}_\otimes^{1+\sigma}\Re\left(\frac{\Phi}{V}\right) \right| + \left| \tilde{r}_\otimes^{1+\sigma}\Re\left(\frac{\Phi}{V}\left(\frac{V}{V_\otimes} - 1\right)\right) \right| \\ &\leq K\|\Psi\|_{*,\sigma,d} + K\left| r_\otimes^\sigma \frac{\Phi}{V} \right| \leq K\|\Psi\|_{*,\sigma,d}. \end{aligned}$$

Using (3.2), we can estimate similarly all the terms in (3.1).

We define similarly, for $g = V_\otimes(g_1 + ig_2) \in C^1(\mathbb{R}^2)$, $\sigma' > 0$

$$\|g\|_{\otimes\otimes,\sigma',d_\otimes} := \|g\|_{C^1(\{\tilde{r}_\otimes\leq 3\})} + \|\tilde{r}_\otimes^{1+\sigma'}g_1\|_{L^\infty(\{\tilde{r}_\otimes\geq 2\})} + \|\tilde{r}_\otimes^{2+\sigma'}g_2\|_{L^\infty(\{\tilde{r}_\otimes\geq 2\})} + \|\tilde{r}_\otimes^{2+\sigma'}\nabla g\|_{L^\infty(\{\tilde{r}_\otimes\geq 2\})}.$$

We have that there exist $C_1, C_2 > 0$ universal constants such that, for $0 < \sigma' < 1$ and any $d, d_\otimes \geq 10$ with $|d - d_\otimes| < \delta$, for any $Vh \in \mathcal{E}_{**,\sigma',d}$, $g = Vh$,

$$C_1\|h\|_{**,\sigma',d} \leq \|g\|_{\otimes\otimes,\sigma',d_\otimes} \leq C_2\|h\|_{**,\sigma',d}.$$

We define the spaces, for $\sigma, \sigma' > 0$,

$$\mathcal{E}_{\otimes,\sigma,d_\otimes} :=$$

$$\{\Phi \in C^2(\mathbb{R}^2, \mathbb{C}), \|\Phi\|_{\otimes,\sigma,d_\otimes} < +\infty, \langle \Phi, Z_{d_\otimes} \rangle = 0, \forall x \in \mathbb{R}^2, \Phi(x_1, x_2) = \overline{\Phi(x_1, -x_2)} = \Phi(-x_1, x_2)\}$$

and

$$\mathcal{E}_{\otimes\otimes,\sigma',d_\otimes} := \{g \in C^1(\mathbb{R}^2, \mathbb{C}), \|g\|_{\otimes\otimes,\sigma',d_\otimes} < +\infty\}.$$

We infer that, from Proposition 2.17, that the operator

$$(\eta L(\cdot) + (1-\eta)V L'(\cdot/V))^{-1} o\Pi_d^\perp$$

goes from $\mathcal{E}_{\otimes\otimes,\sigma',d_\otimes}$ to $\mathcal{E}_{\otimes,\sigma,d_\otimes}$, and that (for $0 < \sigma < \sigma' < 1$)

$$\|(\eta L(\cdot) + (1-\eta)V L'(\cdot/V))^{-1} o\Pi_d^\perp\|_{\mathcal{E}_{\otimes\otimes,\sigma',d_\otimes} \rightarrow \mathcal{E}_{\otimes,\sigma,d_\otimes}}$$

is bounded independently of c, d and d_{\otimes} if $|d - d_{\otimes}| < \delta$. Indeed, the norms $\|\cdot\|_{*,\sigma,d}$ and $\|\cdot\|_{\otimes,\sigma,d_{\otimes}}$ are equivalent, as well as the norms $\|\cdot\|_{**,\sigma',d}$ and $\|\cdot\|_{\otimes\otimes,\sigma',d_{\otimes}}$ for any $\sigma, \sigma' > 0$. About the orthogonality, we replaced $\langle \Phi, Z_d \rangle = 0$ by $\langle \Phi, Z_{d_{\otimes}} \rangle = 0$. This does not change the proof of Proposition 2.17, since when we argue by contradiction, if for a universal constant $|\lambda| \leq \delta$ we took the orthogonality $\langle \Phi, Z_{d+\lambda} \rangle = 0$ instead of $\langle \Phi, Z_d \rangle = 0$, the proof does not change, given that δ is small enough (independently of d). To be specific, we have to take δ small enough such that $\langle \partial_{x_1} V_1, \partial_{x_1} V_1(\cdot + \lambda) \rangle > 0$ for all $\lambda \in]-\delta, \delta[$.

Therefore, we take a sequence $\mathcal{D}^{(n)} > 0$ going to infinity such that $|\mathcal{D}^{(n+1)} - \mathcal{D}^{(n)}| < \delta/2$, and for any given d large enough, there exists $k(d)$ such that $d \in]D^{(k(d))} - \delta/2, D^{(k(d))} + \delta/2[$, and the proof of Proposition 2.17 holds with the orthogonality $\langle \Phi, Z_{D^{(k(d))}} \rangle = 0$ for any value of d in $]D^{(k(d))} - \delta/2, D^{(k(d))} + \delta/2[$. We denote $D^{(k(d))} = d_{\otimes}$. The inversion of the linearized operator then holds for $d \in]D^{(n)} - \delta/2, D^{(n)} + \delta/2[$ with $D^{(n)} = d_{\otimes}$, for all $n \in \mathbb{N}$ large enough.

Furthermore, the contraction arguments given in the proof of Proposition 2.21 still hold (because the norms are equivalent), hence we can define $\Phi_{c,d}$ by a fixed point argument if $\frac{1}{2d} \leq c \leq \frac{2}{d}$ and $|d - d_{\otimes}| < \delta$ in the space $\mathcal{E}_{\otimes,\sigma,d_{\otimes}}$ that does not depend on d .

We want to emphasize the fact that we change a little the definition of the spaces compared to section 2. In particular, for $\Phi = V\Psi$, the norm $\|\cdot\|_{\otimes,\sigma,d_{\otimes}}$ is on the function Φ , and before, for $\|\cdot\|_{*,\sigma,d}$, it was on Ψ . This is because V depends on d , and we want to remove any dependence on d . The same remark holds for $\|\cdot\|_{\otimes\otimes,\sigma',d_{\otimes}}$ and $\|\cdot\|_{\otimes\otimes,\sigma',d}$ (with $g = Vh$).

We continue, and we define

$$H(\Phi, c, d) := (\eta L(\cdot) + (1 - \eta)VL'(\cdot/V))^{-1}(-\Pi_d^\perp(F(\Phi/V))) + \Phi.$$

The function $\Phi_{c,d} \in \mathcal{E}_{\otimes,\sigma,d_{\otimes}}$ is defined, for $\frac{1}{2d} < c < \frac{2}{d}$ and $|d - d_{\otimes}| < \delta$, by being the only solution in a ball of $\mathcal{E}_{\otimes,\sigma,d_{\otimes}}$ (with a radius depending on $\sigma, \sigma', d_{\otimes}$ and c but not d) to the implicit equation on Φ : $H(\Phi, c, d) = 0$. This means that we shall be able to use the implicit function theorem in the space $\mathcal{E}_{\otimes,\sigma,d_{\otimes}}$ on the equation $H(\Phi, c, d) = 0$ to show that $\Phi_{c,d}$ is a C^1 function of d in $\mathcal{E}_{\otimes,\sigma,d_{\otimes}}$ (for values of d such that $\frac{1}{2d} < c < \frac{2}{d}$ and $|d - d_{\otimes}| < \delta$). We want to differentiate this equation with respect to Φ at a fixed c and d , and show that we can invert the operator obtained when we take Φ close to $\Phi_{c,d}$. Since $(\eta L(\cdot) + (1 - \eta)VL'(\cdot/V))^{-1}$ and Π_d^\perp are linear operators that do not depend on Φ , it is easy to check that $H(\Phi, c, d)$ is differentiable with respect to Φ , and we compute

$$d_\Phi H(\Phi, c, d)(\varphi) = (\eta L(\cdot) + (1 - \eta)VL'(\cdot/V))^{-1}(\Pi_d^\perp(-d_\Psi F(\varphi/V))) + \varphi.$$

To show that $d_\Phi H(\Phi, c, d) : \mathcal{E}_{\otimes,\sigma,d_{\otimes}} \rightarrow \mathcal{E}_{\otimes,\sigma,d_{\otimes}}$ and that it is invertible, it is enough to check that

$$\|(\eta L(\cdot) + (1 - \eta)VL'(\cdot/V))^{-1}(\Pi_d^\perp(d_\Psi F(\cdot/V)))\|_{\mathcal{E}_{\otimes,\sigma,d_{\otimes}} \rightarrow \mathcal{E}_{\otimes,\sigma,d_{\otimes}}} = o_{c \rightarrow 0}^\sigma(1), \quad (3.3)$$

which implies that $d_\Phi H(\Phi, c, d)$ is a small perturbation of Id for small values of c (at fixed σ). From Proposition 2.17, we have that $\|(\eta L(\cdot) + (1 - \eta)VL'(\cdot/V))^{-1} \circ \Pi_d^\perp\|_{\mathcal{E}_{\otimes\otimes,\sigma',d_{\otimes}} \rightarrow \mathcal{E}_{\otimes,\sigma,d_{\otimes}}}$ is bounded independently of d and d_{\otimes} if $|d - d_{\otimes}| < \delta$, thus it is enough to check that, for some $\sigma' > \sigma$ (we will take $\sigma' = \frac{1+\sigma}{2} > \sigma$),

$$\|d_\Psi F(\cdot)\|_{\mathcal{E}_{\otimes,\sigma,d_{\otimes}} \rightarrow \mathcal{E}_{\otimes\otimes,\sigma',d_{\otimes}}} = o_{c \rightarrow 0}^{\sigma,\sigma'}(1).$$

This is a consequence of the following lemma (for functions $\Phi = V\Psi$ such that $\|\Psi\|_{*,\sigma,d} = o_{c \rightarrow 0}^\sigma(1)$, which is the case if Φ is near $\Phi_{c,d}$ since $\|\Psi_{c,d}\|_{*,\sigma,d} \leq K(\sigma, \sigma')c^{1-\sigma'}$), where we do the computations with the $*$ -norms since they are equivalent, with uniform constants, to the \otimes -norms. We define

$$\gamma(\sigma) := \frac{1 + \sigma}{2} > \sigma.$$

Lemma 3.1 *There exists $C > 0$ such that, for $0 < \sigma < 1$ and functions $\Phi = V\Psi$, $\varphi = V\psi \in \mathcal{E}_{*,\sigma,d}$, if $\frac{1}{2d} < c < \frac{2}{d}$ and $\|\Psi\|_{*,\sigma,d} \leq 1$, then*

$$\|d_\Psi F(\psi)\|_{**, \gamma(\sigma), d} \leq C \|\Psi\|_{*,\sigma,d} \|\psi\|_{*,\sigma,d}.$$

3.1.2 Proof of the differentiability of $\Phi_{c,d}$ with respect of c and d

We shall now show that $c \mapsto \Phi_{c,d}$ is C^1 and compute estimates on $\partial_c \Psi_{c,d}$ at fixed d , and then show that $d \mapsto \Phi_{c,d}$ is C^1 at fixed c and estimate $\partial_d \Phi_{c,d}$. These estimates will be useful in subsection 4.6. For $d \mapsto \Phi_{c,d}$, we will use the implicit function theorem (see Lemma 3.3), but we start here with the derivation with respect to c .

Lemma 3.2 *For $0 < \sigma < 1$, there exists $c_0(\sigma) > 0$ such that, at fixed $d > \frac{1}{2c_0(\sigma)}$,*

$$c \mapsto \Phi_{c,d} \in C^1 \left(\left[\frac{1}{2d}, \frac{2}{d} \right] \cap]0, c_0(\sigma)[, \mathcal{E}_{*,\sigma,d} \right).$$

Remark that, at fixed d , $\partial_c \Phi_{c,d} = V \partial_c \Psi_{c,d}$.

Proof In this proof, we consider a fixed $d > \frac{1}{2c_0(\sigma)}$. We define, for $c \in \mathbb{R}$ such that $\frac{1}{2d} < c < \frac{2}{d}$ and $0 < c < c_0(\sigma)$, the operator

$$\mathbb{H}_c : \Phi \mapsto (\eta L(\cdot) + (1 - \eta)VL'(\cdot/V))^{-1}(\Pi_d^\perp(F(\Phi/V)))$$

from $\mathcal{E}_{\otimes,\sigma,d_\otimes}$ to $\mathcal{E}_{\otimes,\sigma,d_\otimes}$. The dependency on c is coming from both F and $(\eta L(\cdot) + (1 - \eta)VL'(\cdot/V))^{-1}$, and in this proof, we will add a subscript on these functions giving the value of c where it is taken. Take $c' \in \mathbb{R}$ such that $\frac{1}{2d} < c' < \frac{2}{d}$ and $0 < c' < c_0(\sigma)$, and let us show that

$$\|\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) - \mathbb{H}_c(\Phi_{c',d})\|_{\otimes,\sigma,d_\otimes} = o_{\varepsilon \rightarrow 0}^{\sigma,c}(1).$$

In particular, remark that we look for a convergence uniform in c' . By definition of the operator $(\eta L(\cdot) + (1 - \eta)VL'(\cdot/V))^{-1}$, the function $\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})$ (in $\mathcal{E}_{\otimes,\sigma,d_\otimes}$) is such that, in the distribution sense,

$$\left(\eta L(\cdot) + (1 - \eta)VL' \left(\frac{\cdot}{V} \right) \right)_{c+\varepsilon} (\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})) = \Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c',d}/V)).$$

Since $\Phi_{c',d} \in C^\infty(\mathbb{R}^2)$, we have that $\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) \in C^\infty(\mathbb{R}^2)$ and the equation is satisfied in the strong sense. Furthermore, since $\Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c',d}/V)) \in \mathcal{E}_{\otimes\otimes,\frac{2+\sigma}{3},d_\otimes}$ by Lemmas 2.22 to 2.24 with $\|\Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c',d}/V))\|_{\otimes\otimes,\frac{2+\sigma}{3},d_\otimes} \leq K(\sigma)$ (since $\Phi_{c',d} \in \mathcal{E}_{\otimes,\frac{2+\sigma}{3},d_\otimes}$ with $\|\Phi_{c',d}\|_{\otimes,\frac{2+\sigma}{3},d_\otimes} \leq K(\sigma)$), we have, by Lemma 2.18, that $\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) \in \mathcal{E}_{\otimes,\gamma(\sigma),d_\otimes}$ (since $\gamma(\sigma) < \frac{2+\sigma}{3}$) with, from Proposition 2.17, $\|\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})\|_{\otimes,\gamma(\sigma),d_\otimes} \leq K(\sigma)$. We check similarly that

$$\left(\eta L(\cdot) + (1 - \eta)VL' \left(\frac{\cdot}{V} \right) \right)_c (\mathbb{H}_c(\Phi_{c',d})) = \Pi_d^\perp(F_c(\Phi_{c',d}/V)).$$

Now, from the definitions of L and L' from Lemma 2.7, we have

$$\begin{aligned} \left(\eta L(\cdot) + (1 - \eta)VL' \left(\frac{\cdot}{V} \right) \right)_{c+\varepsilon} (\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})) &= \left(\eta L(\cdot) + (1 - \eta)VL' \left(\frac{\cdot}{V} \right) \right)_c (\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})) \\ &\quad - i\varepsilon \eta \partial_{x_2} \mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) \\ &\quad - i\varepsilon (1 - \eta) V \partial_{x_2} \left(\frac{\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})}{V} \right), \end{aligned}$$

and therefore

$$\begin{aligned} &\left(\eta L(\cdot) + (1 - \eta)VL' \left(\frac{\cdot}{V} \right) \right)_c (\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) - \mathbb{H}_c(\Phi_{c',d})) \\ &= -(\Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c',d}/V) - F_c(\Phi_{c',d}/V))) \\ &\quad - i\varepsilon \left(\eta \partial_{x_2} \mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) + (1 - \eta) V \partial_{x_2} \left(\frac{\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})}{V} \right) \right). \end{aligned}$$

We check, using $\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) \in \mathcal{E}_{\otimes,\gamma(\sigma),d_\otimes}$, $\|\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})\|_{\otimes,\gamma(\sigma),d_\otimes} \leq K(\sigma)$ that

$$i\varepsilon \left(\eta \partial_{x_2} \mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) + (1 - \eta) V \partial_{x_2} \left(\frac{\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})}{V} \right) \right) \in \mathcal{E}_{\otimes\otimes,\gamma(\sigma),d_\otimes},$$

with

$$\left\| i\varepsilon \left(\eta \partial_{x_2} \mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) + (1-\eta)V \partial_{x_2} \left(\frac{\mathbb{H}_{c+\varepsilon}(\Phi_{c',d})}{V} \right) \right) \right\|_{\otimes \otimes, \gamma(\sigma), d_\otimes} \leq K(\sigma)\varepsilon.$$

In particular, by Proposition 2.17 (from $\mathcal{E}_{\otimes \otimes, \gamma(\sigma), d_\otimes}$ to $\mathcal{E}_{\otimes, \sigma, d_\otimes}$), we have

$$\begin{aligned} & \|\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) - \mathbb{H}_c(\Phi_{c',d})\|_{\otimes, \sigma, d_\otimes} \\ & \leq K(\sigma) \|\Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c',d}/V) - F_c(\Phi_{c',d}/V))\|_{\otimes \otimes, \gamma(\sigma), d_\otimes} \\ & + K(\sigma)\varepsilon. \end{aligned}$$

We recall that

$$F_c(\Psi) = E - ic\partial_{x_2}V + V(1-\eta)(-\nabla\Psi \cdot \nabla\Psi + |V|^2S(\Psi)) + R_c(\Psi),$$

therefore

$$F_{c+\varepsilon}(\Phi_{c',d}/V) - F_c(\Phi_{c',d}/V) = -i\varepsilon\partial_{x_2}V + R_{c+\varepsilon}(\Phi_{c',d}/V) - R_c(\Phi_{c',d}/V).$$

By Lemma 2.5 (for $i\partial_{x_2}V$) and the definition of R_c , we check that, for any $0 < \sigma < 1$, since $\|\Psi_{c',d}\|_{*,\sigma,d} \leq K(\sigma)c_0(\sigma)^{1-\gamma(\sigma)} \leq K(\sigma)$,

$$\|\Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c',d}/V) - F_c(\Phi_{c',d}/V))\|_{\otimes \otimes, \sigma, d_\otimes} \leq K(\sigma)\frac{\varepsilon}{c}.$$

We conclude that

$$\|\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) - \mathbb{H}_c(\Phi_{c',d})\|_{\otimes, \sigma, d_\otimes} = o_{\varepsilon \rightarrow 0}^{\sigma, c}(1),$$

thus $\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) \rightarrow \mathbb{H}_c(\Phi_{c',d})$ when $\varepsilon \rightarrow 0$ in $\mathcal{E}_{\otimes, \sigma, d_\otimes}$ uniformly in c' . We remark that it is also uniform in d in any compact set of $]0, c_0(\sigma)[$.

The next step is to show that $c \mapsto \Phi_{c,d}$ is a continuous function in $\mathcal{E}_{*,\sigma,d}$. Take ε_n a sequence such that $\varepsilon_n \rightarrow 0$ when $n \rightarrow \infty$, then $\|\Phi_{c+\varepsilon_n,d}\|_{*,\sigma,d} \leq K_0(\sigma, \sigma')(c + \varepsilon_n)^{1-\sigma'}$ (for $K_0(\sigma, \sigma')$ the constant in Proposition 2.21), and (in the strong sense)

$$\left(\eta L(\cdot) + (1-\eta)VL' \left(\frac{\cdot}{V} \right) \right)_{c+\varepsilon_n}(\Phi_{c+\varepsilon_n,d}) + \Pi_d^\perp(F_{c+\varepsilon_n}(\Phi_{c+\varepsilon_n,d}/V)) = 0.$$

With the same arguments as in step 1 of the proof of Proposition 2.17, we check that, up to a subsequence, $\Phi_{c+\varepsilon_n} \rightarrow \Phi$ locally uniformly in \mathbb{R}^2 for some function $\Phi \in \mathcal{E}_{*,\sigma,d}$ such that $\|\Phi\|_{*,\sigma,d} \leq K_0(\sigma, \sigma')c^{1-\sigma'}$. Then, since

$$\mathbb{H}_{c+\varepsilon_n}(\Phi_{c+\varepsilon_n,d}) + \Phi_{c+\varepsilon_n,d} = 0,$$

by taking the limit when $n \rightarrow \infty$, up to a subsequence, since $\mathbb{H}_{c+\varepsilon}(\Phi_{c',d}) \rightarrow \mathbb{H}_c(\Phi_{c',d})$ when $\varepsilon \rightarrow 0$ in $\mathcal{E}_{*,\sigma,d}$ (the norm is equivalent to the one of $\mathcal{E}_{\otimes, \sigma, d_\otimes}$) uniformly in c' , we have

$$\mathbb{H}_c(\Phi) + \Phi = 0.$$

But then, $\Phi \in \mathcal{E}_{*,\sigma,d}$, $\|\Phi\|_{*,\sigma,d} \leq K_0(\sigma, \sigma')c^{1-\sigma'}$ and $\mathbb{H}_c(\Phi) + \Phi = H(\Phi, c, d) = 0$. By Proposition 2.21, this implies that $\Phi = \Phi_{c,d}$, therefore $\Phi_{c,d}$ is an accumulation point of $\Phi_{c+\varepsilon_n,d}$. It is the only accumulation point, since any other will also satisfy $\Phi \in \mathcal{E}_{*,\sigma,d}$, $\|\Phi\|_{*,\sigma,d} \leq K_0(\sigma, \sigma')c^{1-\sigma'}$ and $H(\Phi, c, d) = 0$. Therefore, $\Phi_{c+\varepsilon_n,d} \rightarrow \Phi_{c,d}$ in $\mathcal{E}_{*,\sigma,d}$, hence $c \mapsto \Phi_{c,d}$ is a continuous function in $\mathcal{E}_{*,\sigma,d}$.

Now, let us show that it is a C^1 function in $\mathcal{E}_{*,\sigma,d}$. Since $\mathbb{H}_c(\Phi_{c,d}) + \Phi_{c,d} = 0$, we have

$$\begin{aligned} & \left(\eta L(\cdot) + (1-\eta)VL' \left(\frac{\cdot}{V} \right) \right)_c(\Phi_{c+\varepsilon,d} - \Phi_{c,d}) \\ & = -(\Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c+\varepsilon,d}/V) - F_c(\Phi_{c,d}/V))) \\ & - i\varepsilon \left(\eta \partial_{x_2} \Phi_{c+\varepsilon,d} + (1-\eta)V \partial_{x_2} \left(\frac{\Phi_{c+\varepsilon,d}}{V} \right) \right). \end{aligned}$$

Furthermore, from $\|\Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c',d}/V) - F_c(\Phi_{c',d}/V))\|_{\otimes \otimes, \sigma, d_\otimes} \leq K(\sigma, c)\varepsilon$ and

$$\left\| i\varepsilon \left(\eta \partial_{x_2} \Phi_{c+\varepsilon,d} + (1-\eta)V \partial_{x_2} \left(\frac{\Phi_{c+\varepsilon,d}}{V} \right) \right) \right\|_{\otimes \otimes, \sigma, d_\otimes} \leq K(\sigma, c)\varepsilon,$$

we deduce that $\|\Phi_{c+\varepsilon,d} - \Phi_{c,d}\|_{\otimes, \sigma, d_\otimes} \leq K(\sigma, c)\varepsilon$.

From the definition of F , we infer that

$$\begin{aligned}
F_{c+\varepsilon}(\Phi_{c+\varepsilon,d}/V) - F_c(\Phi_{c,d}/V) &= -i\varepsilon\partial_{x_2}V \\
&+ V(1-\eta)(-\nabla\Psi_{c+\varepsilon,d}\cdot\nabla\Psi_{c+\varepsilon,d} + \nabla\Psi_{c,d}\cdot\nabla\Psi_{c,d}) \\
&+ V(1-\eta)|V|^2(S(\Psi_{c+\varepsilon,d}) - S(\Psi_{c,d})) \\
&+ R_{c+\varepsilon}(\Psi_{c+\varepsilon,d}) - R_c(\Psi_{c,d}).
\end{aligned}$$

Now, regrouping the terms of $\Pi_d^\perp(d_\Psi F_c((\Phi_{c+\varepsilon,d} - \Phi_{c,d})/V))$ and using $\|\Phi_{c+\varepsilon,d} - \Phi_{c,d}\|_{\otimes,\sigma,d_\otimes} \leq K(\sigma,c)\varepsilon$ for the remaining nonlinear terms (which will be at least quadratic in $\Phi_{c+\varepsilon,d} - \Phi_{c,d}$, since F is C^∞ with respect to Ψ), as well as the fact that $c \mapsto R_c \in C^\infty(]0, c_0(\sigma)[, C^1(\mathbb{R}^2))$, for any $0 < \sigma < 1$,

$$\begin{aligned}
\Pi_d^\perp(F_{c+\varepsilon}(\Phi_{c+\varepsilon,d}/V) - F_c(\Phi_{c,d}/V)) &= \Pi_d^\perp(d_\Psi F_c((\Phi_{c+\varepsilon,d} - \Phi_{c,d})/V)) \\
&+ \varepsilon\Pi_d^\perp(-i\partial_{x_2}V) \\
&+ O_{\|\cdot\|_{**,\sigma,d}}^{\sigma,c}(\varepsilon^2),
\end{aligned}$$

where $O_{\|\cdot\|_{**,\sigma,d}}^{\sigma,c}(\varepsilon^2)$ is a quantity going to 0 as ε^2 when $\varepsilon \rightarrow 0$ in the norm $\|\cdot\|_{**,\sigma,d}$ at fixed σ, c . We deduce that

$$\begin{aligned}
&\left(\text{Id} + (\eta L(\cdot) + (1-\eta)V L'(\frac{\cdot}{V}))_c^{-1}(\Pi_d^\perp(d_\Psi F_c(\cdot/V)))\right)((\Phi_{c+\varepsilon,d} - \Phi_{c,d})) \\
&= (\eta L(\cdot) + (1-\eta)V L'(\frac{\cdot}{V}))_c^{-1} \left(-\varepsilon\Pi_d^\perp(-i\partial_{x_2}V) - i\varepsilon \left(\eta\partial_{x_2}\Phi_{c+\varepsilon,d} + (1-\eta)V\partial_{x_2}\left(\frac{\Phi_{c+\varepsilon,d}}{V}\right) \right) \right) \\
&+ (\eta L(\cdot) + (1-\eta)V L'(\frac{\cdot}{V}))_c^{-1} (O_{\|\cdot\|_{**,\sigma,d}}^{\sigma,c}(\varepsilon^2)),
\end{aligned}$$

and we have shown that $\left(\text{Id} + (\eta L(\cdot) + (1-\eta)V L'(\frac{\cdot}{V}))_c^{-1}(\Pi_d^\perp(\frac{1}{V}d_\Phi F_c(\cdot/V)))\right)$ is invertible from $\mathcal{E}_{\otimes,\sigma,d_\otimes}$ to $\mathcal{E}_{\otimes,\sigma,d_\otimes}$ (with an operator norm equal to $1 + o_{c \rightarrow 0}^\sigma(1)$ if taken in $\Phi = \Phi_{c,d}$, see Lemma 3.1). Furthermore, $\Phi_{c,d}$ is continuous with respect to c in $\mathcal{E}_{\otimes,\gamma(\sigma),d_\otimes}$ (with the same computations as previously, replacing σ by $\gamma(\sigma)$), therefore

$$\eta\partial_{x_2}\Phi_{c+\varepsilon,d} + (1-\eta)V\partial_{x_2}\left(\frac{\Phi_{c+\varepsilon,d}}{V}\right) \rightarrow \eta\partial_{x_2}\Phi_{c,d} + (1-\eta)V\partial_{x_2}\left(\frac{\Phi_{c,d}}{V}\right)$$

in $\mathcal{E}_{\otimes\otimes,\gamma(\sigma),d_\otimes}$ when $\varepsilon \rightarrow 0$. We deduce that $c \mapsto \Phi_{c,d}$ is C^1 in $\mathcal{E}_{\otimes,\sigma,d_\otimes}$ (and therefore in $\mathcal{E}_{*,\sigma,d}$). \square

Now, we infer, with similar arguments, the differentiability of $\Phi_{c,d}$ with respect to d .

Lemma 3.3 *For $0 < \sigma < 1$, there exists $c_0(\sigma) > 0$ such that, for $0 < c < c_0(\sigma)$,*

$$d \mapsto \Phi_{c,d} \in C^1 \left(\left[\frac{1}{2c}, \frac{2}{c} \right] \cap \left[d_\otimes - \frac{\delta}{2}, d_\otimes + \frac{\delta}{2} \right], \mathcal{E}_{\otimes,\sigma,d_\otimes} \right).$$

We recall that $\delta > 0$ is defined at the beginning of this subsection.

We check easily by standard elliptic regularity arguments that $\partial_c\Phi_{c,d} \in C^\infty(\mathbb{R}^2, \mathbb{C})$. Furthermore, $c \mapsto \Phi_{c,d}$ is C^1 with values in $\mathcal{E}_{*,\sigma,d}$, therefore $\partial_c\nabla\Phi_{c,d}$ is well defined (in $C^0(\mathbb{R}^2, \mathbb{C})$). Let us show that it is equal to $\nabla\partial_c\Phi_{c,d}$. For $\varphi \in C_c^\infty(\mathbb{R}^2, \mathbb{C})$, we have, by derivation under an integral, that

$$\begin{aligned}
\int_{\mathbb{R}^2} \partial_c\nabla\Phi_{c,d}\varphi &= \partial_c \int_{\mathbb{R}^2} \nabla\Phi_{c,d}\varphi \\
&= -\partial_c \int_{\mathbb{R}^2} \Phi_{c,d}\nabla\varphi \\
&= -\int_{\mathbb{R}^2} \partial_c\Phi_{c,d}\nabla\varphi \\
&= \int_{\mathbb{R}^2} \nabla\partial_c\Phi_{c,d}\varphi.
\end{aligned}$$

Therefore $\partial_c\nabla\Phi_{c,d} = \nabla\partial_c\Phi_{c,d}$ in the distribution sense, and thus in the strong sense. Furthermore, thanks to the equation $\eta L(\Phi_{c,d}) + (1-\eta)V L'(\Psi_{c,d}) + F(\Psi_{c,d}) = \lambda(c,d)Z_d$, we can isolate $\Delta\Phi_{c,d}$ as in (2.11), and show in particular

that it is a C^1 function of c . By similar arguments as for the gradient, we can show that $\partial_c \Delta \Phi_{c,d} = \Delta \partial_c \Phi_{c,d}$. Furthermore, the same proof holds if we differentiate $\Phi_{c,d}$ with respect to d . We can therefore inverse derivatives in position and derivatives with respect to c or d on $\Phi_{c,d}$.

Let us also show that $(c, d) \mapsto \partial_c \Phi_{c,d}$ is a continuous function from $\Omega := \{(c, d) \in \mathbb{R}^2, 0 < c < c_0(\sigma), \frac{1}{2c} < d < \frac{2}{c}\}$ to $\mathcal{E}_{*,\sigma,d}$. With the same compactness argument used in the proof of the continuity of $c \mapsto \Phi_{c,d}$, we can show that $(c, d) \mapsto \partial_c \Phi_{c,d}$ is continuous from Ω to $\mathcal{E}_{*,\sigma,d}$. From the proof of Lemma 3.2, we have that

$$\begin{aligned} & \left(\text{Id} + \left(\eta L(\cdot) + (1 - \eta) V L' \left(\frac{\cdot}{V} \right) \right)^{-1} (\Pi_d^\perp (d_\Psi F(\cdot/V))) \right) (\partial_c \Phi_c) \\ &= \Pi_d^\perp (\partial_c F(\Phi_{c,d}/V)) - i\eta \partial_{x_2} \Phi_{c,d} + (1 - \eta) V \partial_{x_2} \left(\frac{\Phi_{c,d}}{V} \right). \end{aligned}$$

Since $(c, d) \mapsto \Phi_{c,d}$ is continuous from Ω to $\mathcal{E}_{*,\sigma,d}$, and that the dependence on (c, d) of the other terms of the right-hand side is explicit, we check that $\Pi_d^\perp (\partial_c F(\Phi_{c,d}/V)) - i\eta \partial_{x_2} \Phi_{c,d} + (1 - \eta) V \partial_{x_2} \left(\frac{\Phi_{c,d}}{V} \right)$ is continuous from Ω to $\mathcal{E}_{**, \gamma(\sigma), d}$. We check also that $(c, d) \mapsto \left(\text{Id} + \left(\eta L(\cdot) + (1 - \eta) V L' \left(\frac{\cdot}{V} \right) \right)^{-1} (\Pi_d^\perp (d_\Psi F(\cdot/V))) \right)$ is continuous from Ω to $\mathcal{E}_{**, \gamma(\sigma), d} \rightarrow \mathcal{E}_{*,\sigma,d}$, and thus $(c, d) \mapsto \partial_c \Phi_{c,d}$ is a continuous function from Ω to $\mathcal{E}_{*,\sigma,d}$. The same proof holds for $(c, d) \mapsto \partial_d \Phi_{c,d}$.

We end this subsection with the symmetries of $\partial_d \Phi_{c,d}$.

Lemma 3.4 *The function $\partial_d \Phi_{c,d}$ satisfies the symmetries: for $x = (x_1, x_2) \in \mathbb{R}^2$,*

$$\partial_d \Phi_{c,d}(x_1, x_2) = \partial_d \Phi_{c,d_c}(-x_1, x_2) = \overline{\partial_d \Phi_{c,d}(x_1, -x_2)}.$$

Proof From subsection 2.3,

$$\forall x = (x_1, x_2) \in \mathbb{R}^2, \Psi_{c,d}(x_1, x_2) = \overline{\Psi_{c,d}(x_1, -x_2)} = \Psi_{c,d}(-x_1, x_2)$$

and V enjoys the same symmetries, therefore for all $d \in \mathbb{R}$ such that $\frac{1}{2c} < d < \frac{2}{c}$,

$$\Phi_{c,d}(x_1, x_2) = \Phi_{c,d}(-x_1, x_2) = \overline{\Phi_{c,d}(x_1, -x_2)}.$$

Since

$$\partial_d \Phi_{c,d} = \lim_{\varepsilon \rightarrow 0} \frac{\Phi_{c,d+\varepsilon} - \Phi_{c,d}}{\varepsilon},$$

these symmetries also hold for $\partial_d \Phi_{c,d}$. □

3.2 End of the construction and properties of Q_c

A consequence of equation (2.23) and Proposition 2.26 is that, for $0 < \sigma < 1$, there exists $c_0(\sigma) > 0$ such that, for $0 < c < c_0(\sigma)$,

$$\eta L(\Phi_{c,d}) + (1 - \eta) V L'(\Psi_{c,d}) + F(\Psi_{c,d}) = \lambda(c, d) Z_d$$

with

$$\lambda(c, d) \int_{\mathbb{R}^2} |\partial_d V|^2 \eta = \pi \left(\frac{1}{d} - c \right) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Following the proof of Proposition 2.26, with Lemmas 3.2 and 3.3, we can check that the $O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$ is continuous with respect of c and d . Therefore, by the intermediate value theorem, there exists $d_c > 0$ such that $\lambda(c, d_c) = 0$, with

$$d_c = \frac{1}{c} + O_{c \rightarrow 0}^\sigma(c^{-\sigma}),$$

for $c > 0$ small enough. Then, for the function $\Phi_{c,d_c} = V \Psi_{c,d_c}$ with $\|\Psi_{c,d_c}\|_{*,\sigma,d_c} \leq K(\sigma, \sigma') c^{1-\sigma'}$, we have

$$\eta L(\Phi_{c,d_c}) + (1 - \eta) V L'(\Psi_{c,d_c}) + F(\Psi_{c,d_c}) = 0,$$

meaning that if we define

$$Q_c := \eta V(1 + \Psi_{c,d_c}) + (1 - \eta) V e^{\Psi_{c,d_c}},$$

then Q_c solves (TW_c).

3.2.1 Behaviour at infinity and energy estimation

Lemma 3.5 *The function Q_c satisfies $Q_c(x) \rightarrow 1$ when $|x| \rightarrow \infty$.*

Proof From $\|\Psi_{c,d_c}\|_{*,\sigma,d_c} \leq K(\sigma,\sigma')c^{1-\sigma'}$ we have $\Psi_{c,d_c}(x) \rightarrow 0$ when $|x| \rightarrow \infty$. Furthermore $|1-V|^2 \leq \frac{C(d_c)}{1+\tilde{r}^2}$ by Lemma 2.3 and $Q_c = Ve^{\Psi_{c,d_c}}$ for large values of $|x|$, hence $Q_c(x) \rightarrow 1$ when $|x| \rightarrow \infty$. \square

In the statement of Theorem 1.1, we have set $Q_c = V + \Gamma_{c,d_c}$, we therefore define

$$\Gamma_{c,d_c} := \eta V \Psi_{c,d_c} + (1-\eta)V(e^{\Psi_{c,d_c}} - 1). \quad (3.4)$$

We compute that

$$\left\| \frac{\Gamma_{c,d_c}}{V} \right\|_{*,\sigma,d_c} \leq K \|\Psi_{c,d_c}\|_{*,\sigma,d_c} + \|(1-\eta)(e^{\Psi_{c,d_c}} - 1 - \Psi_{c,d_c})\|_{*,\sigma,d_c},$$

and since $\|\Psi_{c,d_c}\|_{*,\sigma,d_c} \leq 1$ for c small enough (depending on σ), we have

$$\|(1-\eta)(e^{\Psi_{c,d_c}} - 1 - \Psi_{c,d_c})\|_{*,\sigma,d_c} \leq K \left\| (1-\eta)\Psi_{c,d_c}^2 \sum_{n=2}^{+\infty} \frac{\Psi_{c,d_c}^{n-2}}{n!} \right\|_{*,\sigma,d_c}.$$

Now, for $0 < \sigma < \sigma' < 1$, we have $\frac{1+\sigma'}{2} > \frac{1+\sigma}{2}$, hence

$$|\Psi_{c,d_c}| \leq K(\sigma,\sigma') \frac{c^{1-\frac{1+\sigma'}{2}}}{(1+\tilde{r})^{\frac{1+\sigma}{2}}} \quad \text{and} \quad |\nabla \Psi_{c,d_c}| \leq K(\sigma,\sigma') \frac{c^{1-\frac{1+\sigma'}{2}}}{(1+\tilde{r})^{1+\frac{1+\sigma}{2}}},$$

therefore

$$|\Psi_{c,d_c}|^2 \leq K(\sigma,\sigma') \frac{c^{1-\sigma'}}{(1+\tilde{r})^{1+\sigma}} \quad \text{and} \quad |\nabla \Psi_{c,d_c}|^2 \leq K(\sigma,\sigma') \frac{c^{1-\sigma'}}{(1+\tilde{r})^{2+\sigma}}.$$

Thus, with $|\nabla^2 \Psi_{c,d_c}| \leq K(\sigma,\sigma') \frac{c^{1-\sigma'}}{(1+\tilde{r})^{1+\sigma}}$, we check that, for any $0 < \sigma < \sigma' < 1$,

$$\left\| (1-\eta)\Psi_{c,d_c}^2 \sum_{n=2}^{+\infty} \frac{\Psi_{c,d_c}^{n-2}}{n!} \right\|_{*,\sigma,d_c} \leq K(\sigma,\sigma')c^{1-\sigma'}.$$

Combining this result with $\|\Psi_{c,d_c}\|_{*,\sigma,d_c} \leq K(\sigma,\sigma')c^{1-\sigma'}$, we deduce that

$$\left\| \frac{\Gamma_{c,d_c}}{V} \right\|_{*,\sigma,d_c} \leq K(\sigma,\sigma')c^{1-\sigma'}. \quad (3.5)$$

In particular, we have, for any $0 < \sigma < \sigma' < 1$, $0 < c < c_0(\sigma,\sigma')$, that

$$|\Gamma_{c,d_c}| \leq \frac{K(\sigma,\sigma')c^{1-\sigma'}}{(1+\tilde{r})^\sigma}, \quad (3.6)$$

$$\left| \Re e \left(\frac{\Gamma_{c,d_c}}{V} \right) \right| \leq \frac{K(\sigma,\sigma')c^{1-\sigma'}}{(1+\tilde{r})^{1+\sigma}}, \quad (3.7)$$

and, if $\tilde{r} \geq 2$,

$$|\nabla \Gamma_{c,d_c}| \leq \left| \nabla \left(\frac{\Gamma_{c,d_c}}{V} \right) \right| + \left| \frac{\nabla V}{V} \right| \times \left| \frac{\Gamma_{c,d_c}}{V} \right|,$$

therefore, using $|\nabla V| \leq \frac{K}{(1+\tilde{r})}$ from Lemma 2.1, we have

$$|\nabla \Gamma_{c,d_c}| \leq \frac{K(\sigma,\sigma')c^{1-\sigma'}}{(1+\tilde{r})^{1+\sigma}}. \quad (3.8)$$

Estimate (3.8) remains true in $\{\tilde{r} \leq 2\}$ since $\|\Gamma_{c,d_c}\|_{C^1(\{\tilde{r} \leq 2\})} \leq \left\| \frac{\Gamma_{c,d_c}}{V} \right\|_{*,\sigma,d_c} \leq K(\sigma,\sigma')c^{1-\sigma'}$. We now show the estimates on Γ_{c,d_c} of Theorem 1.1.

Lemma 3.6 For $+\infty \geq p > 2$, there exists $c_0(p) > 0$ such that if $0 < c < c_0(p)$, we have $\Gamma_{c,d_c} \in L^p(\mathbb{R}^2)$, $\nabla\Gamma_{c,d_c} \in L^{p-1}(\mathbb{R}^2)$ and

$$\|\Gamma_{c,d_c}\|_{L^p(\mathbb{R}^2)} + \|\nabla\Gamma_{c,d_c}\|_{L^{p-1}(\mathbb{R}^2)} = o_{c \rightarrow 0}(1).$$

Proof If $p = +\infty$, using (3.6) and (3.8), we infer

$$\begin{aligned} \|\Gamma_{c,d_c}\|_{L^\infty(\mathbb{R}^2)} &\leq K(\sigma)c^{1-\sigma}, \\ \|\nabla\Gamma_{c,d_c}\|_{L^\infty(\mathbb{R}^2)} &\leq K(\sigma)c^{1-\sigma}, \end{aligned}$$

hence the result holds. If $2 < p < +\infty$ then, by (3.6),

$$\int_{\mathbb{R}^2} |\Gamma_{c,d_c}|^p \leq \int_{\mathbb{R}^2} \frac{\|\Gamma_{c,d_c}\|_{*,\sigma,d_c}^p}{(1+\tilde{r})^{p\sigma}} dx \leq \int_{\mathbb{R}^2} \frac{K(\sigma,\sigma')c^{(1-\sigma')p}}{(1+\tilde{r})^{p\sigma}} dx.$$

Taking $0 < \sigma < \sigma' < 1$ such that $p\sigma > 2$ then gives the result. Furthermore, by (3.8),

$$\int_{\mathbb{R}^2} |\nabla\Gamma_{c,d_c}|^p \leq \int_{\mathbb{R}^2} \frac{K(\sigma,\sigma')c^{(1-\sigma')p}}{(1+\tilde{r})^{p(\sigma+1)}} dx,$$

so for $p > 1$ we can take $0 < \sigma < \sigma' < 1$ such that $p(\sigma+1) > 2$ and we have the result. \square

Remark that we can have better estimates on Γ_{c,d_c} , in particular if we look at real and imaginary parts of $\frac{\Gamma_{c,d_c}}{V}$. For instance it is possible to show that

$$\left\| \Re\left(\frac{\Gamma_{c,d_c}}{V}\right) \right\|_{L^p(\{\tilde{r} \geq 1\})} = o_{c \rightarrow 0}(1)$$

for $p > 1$ instead of $p > 2$. This estimate does not hold for \tilde{r} small since it is not clear that Ψ_{c,d_c} is bounded there (but Φ_{c,d_c} is). This is due to the fact that the zeros of Q_c are not exactly those of V .

Lemma 3.7 The travelling wave Q_c has finite energy, that is:

$$E(Q_c) = \frac{1}{2} \int_{\mathbb{R}^2} |\nabla Q_c|^2 + \frac{1}{4} \int_{\mathbb{R}^2} (1 - |Q_c|^2)^2 < +\infty.$$

Proof Far from the vortices, $\nabla Q_c = \nabla(V_1 V_{-1})e^{\Psi_{c,d_c}} + \nabla\Psi_{c,d_c} V_1 V_{-1} e^{\Psi_{c,d_c}}$. We know that, for $\tilde{r} \geq 1$,

$$|\nabla\Psi_{c,d_c}| \leq \frac{K(\sigma)}{\tilde{r}^{1+\sigma}}$$

and (by Lemma 2.3)

$$|\nabla(V_1 V_{-1})| \leq \frac{K(c)}{\tilde{r}^2},$$

hence

$$|\nabla Q_c|^2 \leq \frac{K(c,\sigma)}{\tilde{r}^{2+2\sigma}}$$

and is therefore integrable. On the other hand,

$$|1 - |Q_c|^2| = \left| 1 - |V_1 V_{-1}|^2 e^{2\Re(\Psi_{c,d_c})} \right| \leq K(1 - |V_1 V_{-1}|^2 + |V_1 V_{-1}|^2 |\Re(\Psi_{c,d_c})|),$$

and we have

$$1 - |V_1 V_{-1}|^2 = O\left(\frac{1}{\tilde{r}^2}\right) \quad \text{and} \quad \Re(\Psi_{c,d_c}) = O\left(\frac{1}{\tilde{r}^{1+\sigma}}\right),$$

therefore

$$(1 - |Q_c|^2)^2 = O\left(\frac{1}{\tilde{r}^{2+2\sigma}}\right)$$

and is integrable. \square

At this point, we have finished the proof of the construction of Q_c . In the next two subsection, we add some estimates on Q_c that will be useful for the differentiability of the branch, and others that are interesting in themselves.

3.2.2 A set of estimations on Q_c

The next Lemma gives additional estimates on Q_c which are more precise but more technical than the ones in Theorem 1.1.

Lemma 3.8 *For any $0 < \sigma < \sigma' < 1$, there exists $c_0(\sigma, \sigma'), K(\sigma, \sigma') > 0$ such that for $0 < c < c_0(\sigma, \sigma')$ we have*

$$\|\Psi_{c,d_c}\|_{*,\sigma,d_c} \leq K(\sigma, \sigma')c^{1-\sigma'}. \quad (3.9)$$

Furthermore, for any $0 < \sigma < 1$, there exist $c_0(\sigma), K(\sigma) > 0$ such that for $0 < c < c_0(\sigma)$ we have

$$\begin{aligned} & \|V\Psi_{c,d_c}\|_{C^1(\tilde{r} \leq 3)} + \|\tilde{r}^\sigma \Im(\Psi_{c,d_c})\|_{L^\infty(\tilde{r} \geq 2)} + \|\tilde{r}^{1+\sigma} \Re(\Psi_{c,d_c})\|_{L^\infty(\tilde{r} \geq 2)} \\ + & \|\tilde{r}^{1+\sigma} \Im(\nabla \Psi_{c,d_c})\|_{L^\infty(\tilde{r} \geq 2)} + \|\tilde{r}^{2+\sigma} \Re(\nabla \Psi_{c,d_c})\|_{L^\infty(\tilde{r} \geq 2)} \\ \leq & K(\sigma)c^{1-\sigma}, \end{aligned} \quad (3.10)$$

$$|1 - |Q_c|| \leq \frac{K(\sigma)}{(1 + \tilde{r})^{1+\sigma}}, \quad (3.11)$$

$$|Q_c - V| \leq \frac{K(\sigma)c^{1-\sigma}}{(1 + \tilde{r})^\sigma}, \quad (3.12)$$

$$||Q_c|^2 - |V|^2| \leq \frac{K(\sigma)c^{1-\sigma}}{(1 + \tilde{r})^{1+\sigma}}, \quad (3.13)$$

$$|\Re(\nabla Q_c \overline{Q_c})| \leq \frac{K(\sigma)}{(1 + \tilde{r})^{1+\sigma}}, \quad (3.14)$$

$$|\Im(\nabla Q_c \overline{Q_c})| \leq \frac{K}{1 + \tilde{r}} \quad (3.15)$$

Equation (3.10) is a slight improvements of (3.9). It is, except for the second derivatives, the estimate in the case $\sigma' = \sigma$.

Proof The first estimate (3.9) comes from the construction of the solution.

We now take χ a cutoff function with value 1 in $\{\tilde{r} \geq 2\}$ and 0 in $\{\tilde{r} \leq 1\}$, we write $\tilde{\Psi} = \chi\Psi_{c,d_c}$ and $\tilde{h} = \chi h$, where h contains the nonlinear and source terms. The functions $\tilde{\Psi} = \tilde{\Psi}_1 + i\tilde{\Psi}_2$ and $\tilde{h} = \tilde{h}_1 + i\tilde{h}_2$ satisfy the system

$$\begin{cases} \Delta \tilde{\Psi}_1 - 2\tilde{\Psi}_1 = -\tilde{h}_1 - 2\Re\left(\frac{\nabla V}{V} \cdot \nabla \tilde{\Psi}\right) - 2(1 - |V|^2)\tilde{\Psi}_1 + c\partial_{x_2}\tilde{\Psi}_2 + \text{Loc}_1(\Psi) \\ \Delta \tilde{\Psi}_2 = -\tilde{h}_2 - 2\Im\left(\frac{\nabla V}{V} \cdot \nabla \tilde{\Psi}\right) + \text{Loc}_2(\Psi) - c\partial_{x_2}\tilde{\Psi}_1, \end{cases}$$

where $\text{Loc}_1(\Psi), \text{Loc}_2(\Psi)$ are localized terms. From Lemmas 2.22 to 2.24, we check that for any $0 < \sigma < 1$,

$$\|\tilde{h}\|_{**,\sigma,d} \leq K(\sigma)c^{1-\sigma}.$$

Furthermore, as in the proof of Proposition 2.17, we check that (using $\|\tilde{\Psi}\|_{*,\sigma/2,d} \leq K(\sigma)c^{1-\sigma}$)

$$\left\| \frac{\nabla V}{V} \cdot \nabla \tilde{\Psi} - 2(1 - |V|^2)\Re(\tilde{\Psi}) + \text{Loc}(\Psi) \right\|_{**,\sigma,d} \leq K(\sigma)c^{1-\sigma}.$$

Finally, with (3.9), for $\sigma' = \frac{1+\sigma}{2} > \sigma$,

$$\|c\partial_{x_2}\tilde{\Psi}\|_{**,\sigma,d} \leq K(\sigma)c\|\tilde{\Psi}\|_{*,\sigma,d} \leq K(\sigma)c^{1+1-\frac{1+\sigma}{2}} \leq K(\sigma)c^{1-\sigma}.$$

With Lemma 2.10 for $\alpha = 1 + \sigma > 0$, we deduce from the first equation of the system that

$$\begin{aligned} & \|(1 + \tilde{r})^{1+\sigma}\tilde{\Psi}_1\|_{L^\infty(\mathbb{R}^2)} \\ \leq & K(\sigma) \left\| (1 + \tilde{r})^{1+\sigma} \left(-\tilde{h}_1 - 2\Re\left(\frac{\nabla V}{V} \cdot \nabla \tilde{\Psi}\right) - 2(1 - |V|^2)\tilde{\Psi}_1 + c\partial_{x_2}\tilde{\Psi}_2 + \text{Loc}_1(\Psi) \right) \right\|_{L^\infty(\mathbb{R}^2)} \\ \leq & K(\sigma)c^{1-\sigma}, \end{aligned}$$

and, by differentiating the equation, by Lemma 2.10 for $\alpha = 2 + \sigma > 0$

$$\begin{aligned} & \|(1 + \tilde{r})^{2+\sigma} \nabla \tilde{\Psi}_1\|_{L^\infty(\mathbb{R}^2)} \\ & \leq K(\sigma) \left\| (1 + \tilde{r})^{2+\sigma} \nabla \left(-\tilde{h}_1 - 2\Re \left(\frac{\nabla V}{V} \cdot \nabla \tilde{\Psi} \right) - 2(1 - |V|^2) \tilde{\Psi}_1 + c \partial_{x_2} \tilde{\Psi}_2 + \text{Loc}_1(\Psi) \right) \right\|_{L^\infty(\mathbb{R}^2)} \\ & \leq K(\sigma) c^{1-\sigma}. \end{aligned}$$

Now, using Lemma 2.8 and $\|(1 + \tilde{r})^{2+\sigma} \nabla \tilde{\Psi}_1\|_{L^\infty(\mathbb{R}^2)} \leq K(\sigma) c^{1-\sigma}$, we infer that

$$\begin{aligned} & \|(1 + \tilde{r})^\sigma \tilde{\Psi}_2\|_{L^\infty(\mathbb{R}^2)} + \|(1 + \tilde{r})^{1+\sigma} \nabla \tilde{\Psi}_2\|_{L^\infty(\mathbb{R}^2)} \\ & \leq K(\sigma) \left\| (1 + \tilde{r})^{2+\sigma} \left(-\tilde{h}_2 - 2\Im \left(\frac{\nabla V}{V} \cdot \nabla \tilde{\Psi} \right) + \text{Loc}_2(\Psi) - c \partial_{x_2} \tilde{\Psi}_1 \right) \right\|_{L^\infty(\mathbb{R}^2)} \\ & \leq K(\sigma) c^{1-\sigma}, \end{aligned}$$

which concludes the proof of (3.10). The other estimates are consequences of it. \square

3.2.3 Estimations on derivatives of $\Phi_{c,d}$ with respect to c and d at $d = d_c$.

We cannot easily compute $\partial_d \Psi_{c,d|d=d_c}$ because of issues locally around the vortices (due to the fact that $\Psi_{c,d}$ is unbounded near $\tilde{r} = 0$, and changing d change the position of the vortices). We infer instead an estimate on $\partial_d \Phi_{c,d|d=d_c}$, as well as an estimate on $\partial_c \Psi_{c,d|d=d_c}$.

Lemma 3.9 *For any $0 < \sigma < \sigma' < 1$, $c \in \mathbb{R}$ such that $\frac{1}{2d} < c < \frac{2}{d}$ and $0 < c < c_0(\sigma, \sigma')$, we have*

$$\|\partial_c \Psi_{c,d|d=d_c}\|_{*,\sigma,d} \leq K(\sigma, \sigma') c^{-\sigma'}$$

and

$$\left\| \frac{\partial_d \Phi_{c,d}}{V} \Big|_{d=d_c} \right\|_{*,\sigma,d_c} \leq K(\sigma, \sigma') c^{1-\sigma'},$$

with $K(\sigma, \sigma') > 0$ depending only on σ, σ' .

4 Differentiability of the branch $c \mapsto Q_c$

The goal of this section is to prove that the constructed branch is C^1 , and to give the leading order term of $\partial_c Q_c$ as $c \rightarrow 0$. The result is the following one.

Proposition 4.1 *For any $+\infty \geq p > 2$, there exists $c_0(p) > 0$ such that*

$$c \mapsto Q_c - 1 \in C^1(]0, c_0(p)[, X_p),$$

with the estimate

$$\left\| \partial_c Q_c + \left(\frac{1 + o_{c \rightarrow 0}(1)}{c^2} \right) \partial_d (V_1(\cdot - d \vec{e}_1^\top) V_{-1}(\cdot + d \vec{e}_1^\top)) \Big|_{d=d_c} \right\|_{X_p} = o_{c \rightarrow 0} \left(\frac{1}{c^2} \right).$$

Proposition 4.1, together with subsection 3.2, ends the proof of Theorem 1.1. Subsections 4.1 to 4.7 are devoted to the proof of Proposition 4.1.

In this section, to make the dependances on c and d clear, we use the following notations. We denote $\Phi_{c,d}$, $\Psi_{c,d}$ and $\Gamma_{c,d}$ in order to emphasize the dependence of Φ, Ψ and Γ in Proposition 2.21 on c and d . A value of d that makes $\lambda(c, d) = 0$ in Proposition 2.26 is written d_c . We will show later on that there exist one and only one value of d_c satisfying this in $] \frac{c}{2}, 2c[$. With these notations, $Q_c = V_1(\cdot - d_c \vec{e}_1^\top) V_{-1}(\cdot + d_c \vec{e}_1^\top) + \Gamma_{c,d_c}$ is the solution of (TW_c) we constructed in section 3.

In subsection 3.1 we showed that $\Phi_{c,d}$ is a C^1 function of both c and d . We also have computed estimates for the derivatives of $\Phi_{c,d}$ with respect to c and d in Lemma 3.9, that will be useful here.

The goal is to show that d_c is a C^1 function of c . We will do this by the implicit function theorem, but this requires a lot of computations. In particular, in Proposition 2.26, d_c was chosen so that

$$\langle L(\Phi_{c,d}) - (1 - \eta)(E - ic\partial_{x_2}V)\Psi_{c,d} + F(\Psi_{c,d}), \partial_d V \rangle = 0,$$

but we may equivalently define it by the implicit equation

$$\int_{B(d\vec{e}_1, d^{\varepsilon'}) \cup B(-d\vec{e}_1, d^{\varepsilon'})} \Re((L(\Phi_{c,d}) - (1 - \eta)E\Psi_{c,d} + F(\Psi_{c,d}))\overline{\partial_d V}) = 0.$$

This is the same equation but the scalar product is not taken on the whole space but only on $B(d\vec{e}_1, d^{\varepsilon'}) \cup B(-d\vec{e}_1, d^{\varepsilon'})$ for some $0 < \varepsilon' < 1$ (we will take $\varepsilon' = 13/24$ but this value is purely technical, other values are possible). The only reason why we take it in the whole space in Lemma 2.26 was because of the boundary terms that will appear in the integration by parts when we write

$$\langle L(\Phi), \partial_d V \rangle = \langle \Phi, L(\partial_d V) \rangle.$$

With the boundary terms on the boundary of $B(\pm d\vec{e}_1, d^{\varepsilon'})$, $\varepsilon' > 0$, we are far enough from the vortices to make them small enough for our estimations. Thanks to this we can separate what happens near the vortex V_1 from what happens near the vortex V_{-1} because now the integrals are in two well separated domain, one around each vortex. We use this in subsection 4.1. We need to differentiate the equation with respect to d . If we write $Q_{c,d} = V + \Gamma_{c,d}$, then $\partial_d Q_c = \partial_d V + \partial_d(\Gamma_{c,d})$. The term $\partial_d V$ is easy to compute and to understand: we just move both vortices in opposite direction. But $\partial_d \Gamma_{c,d}$ is very difficult to understand, and our estimations on Γ_{c,d_c} are not enough to compute easily what happens with sufficient precision to control its contribution. We would rather write $Q_{c,d}$ under the form

$$Q_{c,d}(x) = (V_1(x - d\vec{e}_1) + \tilde{\Gamma}_1(x - d\vec{e}_1)) + (V_{-1}(x + d\vec{e}_1) + \tilde{\Gamma}_{-1}(x + d\vec{e}_1)) + \text{Err}$$

where $\tilde{\Gamma}_1(x - d\vec{e}_1)$ is centered near V_1 , is small and is here because of the existence of V_{-1} far away. Then the term we understand is

$$\partial_{x_1+d}(V_1(x - d\vec{e}_1) + \tilde{\Gamma}_1(x - d\vec{e}_1))$$

which is what changes near the center of V_1 when we move only the other vortex. This can be computed more easily and that is what we do in subsection 4.3. This term is easy to compute only near the vortex V_1 , and that is one of the reasons we work only on $B(d\vec{e}_1, d^{\varepsilon'})$. The main contribution to the variation of the position of V_{-1} is as expected from the source term $E - ic\partial_{x_2}V$. This is the computation of subsection 4.4.

Furthermore, most estimations boils down to what happen near each vortex, see for instance the contribution of E in step 5 of the proof of Proposition 2.26, where we separate the contribution far from both vortices and close to them. By integrating only on $B(d\vec{e}_1, d^{\varepsilon'})$ we reduce the number of estimations we need to do. Moreover, in such a ball the contribution of the vortex V_{-1} and its derivatives are easy to compute, see subsection 4.2.

Subsection 4.5 gathers all the estimations needed to show that only the contribution from the source term is of leading order. Subsection 4.6 and 4.7 are easy computations using previous subsections to compute the first order term of $\partial_c Q_c$.

The main and most difficult part is subsection 4.3. We want to show that $\partial_{x_1+d}(\tilde{\Gamma}_1(x - d\vec{e}_1))$ is much smaller than $\tilde{\Gamma}_1(x - d\vec{e}_1)$, i.e. that the derivative with respect to $x_1 + d$ gives us additional smallness in c . For this we do a proof by contradiction which follows closely what was done in the proof of Proposition 2.17.

We define the following differential operators:

$$\partial_{y_1} := \partial_{x_1} - \partial_d,$$

$$\partial_{z_1} := \partial_{x_1} + \partial_d.$$

These notations follow the definitions of $y_1 = x_1 - d$ and $z_1 = x_1 + d$ from (2.1). The derivative in d is taken at fixed c . The function $\partial_d \Phi_{c,d}$ is the derivative of Φ with respect to d at fixed c and we shall use the notation

$$\partial_d \Phi_{c,d_c} := \partial_d \Phi_{c,d|d=d_c},$$

and similarly for $\partial_d \Gamma_{c,d_c}$ and $\partial_d \Psi_{c,d_c}$. The derivatives ∂_{y_1} and ∂_{z_1} behave naturally on function depending on x and d only through y or z , as shown in the following lemma.

Lemma 4.2 For any $\mathfrak{F} \in C^1(\mathbb{R}^2, \mathbb{C})$, we have

$$\partial_{y_1}(\mathfrak{F}(z)) = \partial_{z_1}(\mathfrak{F}(y)) = 0$$

and

$$\begin{aligned}\partial_{y_1}(\mathfrak{F}(y)) &= 2\partial_{x_1}\mathfrak{F}(y), \\ \partial_{z_1}(\mathfrak{F}(z)) &= 2\partial_{x_1}\mathfrak{F}(z).\end{aligned}$$

We have an estimate on $\partial_d\Phi_{c,d|d=d_c}$, but it is not enough to show that d_c is a C^1 function of c . The main idea of the proof is to compute an estimate on $\partial_{z_1}\Phi_{c,d_c} = \partial_{x_1}\Phi_{c,d_c} + \partial_d\Phi_{c,d_c}$ near the vortex V_1 which is better than the ones on $\partial_{x_1}\Phi_{c,d_c}$ and $\partial_d\Phi_{c,d_c}$. In particular we will have $\partial_{z_1}\Phi_{c,d_c} = o_{c \rightarrow 0}(c^{1+\lambda})$ for some $\lambda > 0$ instead of $o_{c \rightarrow 0}(c^{1-\sigma})$ for $\sigma > 0$. This estimate is done in Proposition 4.5. First, we compute a first rough estimate on $\partial_{z_1}\Psi_{c,d}$, which is a corollary of Lemma 3.3.

Corollary 4.3 For χ a smooth cutoff function with value 1 in $\{r_{-1} \geq 3\}$ and 0 in $\{r_{-1} \leq 2\}$, for $0 < \sigma < \sigma' < 1$, there exist $c_0(\sigma, \sigma') > 0$ such that, for $0 < c < c_0(\sigma, \sigma')$, we have

$$\begin{aligned}& \|V\chi\partial_{z_1}\Psi_{c,d|d=d_c}\|_{C^1(\{\bar{r} \leq 3\})} \\ & + \|\tilde{r}^{1+\sigma}\Re(\partial_{z_1}\Psi_{c,d|d=d_c})\|_{L^\infty(\{\bar{r} \geq 2\})} + \|\tilde{r}^{2+\sigma}\nabla\Re(\partial_{z_1}\Psi_{c,d|d=d_c})\|_{L^\infty(\{\bar{r} \geq 2\})} \\ & + \|\tilde{r}^\sigma\Im(\partial_{z_1}\Psi_{c,d|d=d_c})\|_{L^\infty(\{\bar{r} \geq 2\})} + \|\tilde{r}^{1+\sigma}\nabla\Im(\partial_{z_1}\Psi_{c,d|d=d_c})\|_{L^\infty(\{\bar{r} \geq 2\})} \\ & \leq K(\sigma, \sigma')c^{1-\sigma'}.\end{aligned}$$

4.1 Recasting the implicit equation defining d_c

At this point, we do not know if d_c is uniquely defined for $c > 0$. We denote by d_c a value defined by the implicit equation on d :

$$\langle \text{TW}_c(Q_{c,d}), \partial_d V \rangle = 0,$$

where

$$Q_{c,d} := V + \Gamma_{c,d},$$

with $\Gamma_{c,d} = \eta V \Psi_{c,d} + (1 - \eta)V(e^{\Psi_{c,d}} - 1)$, which is a C^1 function of d and c in $\mathcal{E}_{*,\sigma,d}$ thanks to subsection 3.1. Remark that d_c is also defined by the implicit equation for $0 < \varepsilon' < 1$:

$$\int_{B(d\bar{e}_1^\dagger, d\varepsilon') \cup B(-d\bar{e}_1^\dagger, d\varepsilon')} \Re(\overline{\partial_d V} \text{TW}_c(Q_{c,d})) = 0,$$

that we will use instead because of the reasons explained at the begining of section 4. We can check easily that $\partial_d Q_{c,d}, \partial_c Q_{c,d} \in C^\infty(\mathbb{R}^2)$ (by looking at the equations they satisfy in the distribution sense and using standard elliptic regularity arguments), and furthermore, that $d \mapsto \partial_d Q_{c,d}$ and $c \mapsto \partial_c Q_{c,d}$ are continuous functions (on their domain of definition in $C_{\text{loc}}^\infty(\mathbb{R}^2)$ for instance). From now on, we take any $0 < \varepsilon' < 1$, but we will fix its value later on. We want to differentiate this quantity with respect to d and take the result at a value d_c such that $\text{TW}_c(Q_{c,d_c}) = 0$ in \mathbb{R}^2 . In particular, we have

$$\begin{aligned}\partial_d \int_{B(d\bar{e}_1^\dagger, d\varepsilon') \cup B(-d\bar{e}_1^\dagger, d\varepsilon')} \Re(\overline{\partial_d V} \text{TW}_c(Q_{c,d}))|_{d=d_c} &= \\ \int_{B(d\bar{e}_1^\dagger, d\varepsilon') \cup B(-d\bar{e}_1^\dagger, d\varepsilon')} \Re(\overline{\partial_d V} \partial_d(\text{TW}_c(Q_{c,d})))|_{d=d_c} &\cdot\end{aligned}$$

Now, by symmetry, we remark that

$$\int_{B(d\bar{e}_1^\dagger, d\varepsilon') \cup B(-d\bar{e}_1^\dagger, d\varepsilon')} \Re(\overline{\partial_d V} \partial_d(\text{TW}_c(Q_{c,d})) = 2 \int_{B(d\bar{e}_1^\dagger, d\varepsilon')} \Re(\overline{\partial_d V} \partial_d(\text{TW}_c(Q_{c,d})).$$

We will use the two operators we have already defined:

$$\partial_{y_1} = \partial_{x_1} - \partial_d \quad \text{and} \quad \partial_{z_1} = \partial_{x_1} + \partial_d.$$

Since $\text{TW}_c(Q_{c,d_c}) = 0$ everywhere in \mathbb{R}^2 , we therefore have $\partial_{x_1}(\text{TW}_c(Q_{c,d_c})) = 0$, hence, at $d = d_c$,

$$\partial_d(\text{TW}_c(Q_{c,d})) = \partial_{z_1}(\text{TW}_c(Q_{c,d})).$$

We write

$$\text{TW}_c(Q_{c,d}) = \text{TW}_c(V) + L(\Gamma_{c,d}) + \text{NL}_V(\Gamma_{c,d}),$$

with

$$L(\Gamma_{c,d}) = -\Delta\Gamma_{c,d} - ic\partial_{x_2}\Gamma_{c,d} - (1 - |V|^2)\Gamma_{c,d} + 2\Re(\bar{V}\Gamma_{c,d})V$$

and

$$\text{NL}_V(\Gamma_{c,d}) := 2\Re(\bar{V}\Gamma_{c,d})\Gamma_{c,d} + |\Gamma_{c,d}|^2(V + \Gamma_{c,d}).$$

We compute

$$\partial_{z_1}(\text{TW}_c(Q_{c,d})) = \partial_{z_1}(\text{TW}_c(V)) + L(\partial_{z_1}\Gamma_{c,d}) + (\partial_{z_1}L)(\Gamma_{c,d}) + \partial_{z_1}(\text{NL}_V(\Gamma_{c,d})),$$

therefore, at $d = d_c$,

$$\begin{aligned} \partial_d \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\bar{\partial}_d V \text{TW}_c(Q_{c,d})) &= \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\bar{\partial}_d V \partial_{z_1}(\text{TW}_c(V))) \\ &+ \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\bar{\partial}_d V L(\partial_{z_1}\Gamma_{c,d})) + \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\bar{\partial}_d V (\partial_{z_1}L)(\Gamma_{c,d})) \\ &+ \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\bar{\partial}_d V \partial_{z_1}(\text{NL}_V(\Gamma_{c,d}))) \end{aligned} \quad (4.1)$$

since the boundary term is 0 (when the differentiation is on the d in $B(d\vec{e}_1, d^{\varepsilon'})$) because $\text{TW}_c(Q_{c,d_c}) = 0$. We need to estimate those four terms at $d = d_c$, and that is the goal of the next subsections. Subsections 4.2 and 4.3 yield estimates on the derivatives of V_{-1} and $\partial_{z_1}\Psi_{c,d}$ respectively in $B'_d := B(d\vec{e}_1, d^{\varepsilon'})$. Subsection 4.4 is about the estimation of

$$\int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\bar{\partial}_d V \partial_{z_1}(\text{TW}_c(V)))$$

which will be the leading order term, and subsection 4.5 shows that all the other terms are smaller for d_c large enough.

4.2 Estimates on the derivatives of V_{-1} in $B(d\vec{e}_1, d^\varepsilon)$

We infer the following estimations.

Lemma 4.4 *For $0 < \varepsilon < 1$, in $B(d\vec{e}_1, d^\varepsilon)$, with the $O(\cdot)$ being always real valued, we have*

$$\begin{aligned} \partial_{x_1}V_{-1} &= \left(O_{d \rightarrow \infty} \left(\frac{1}{d^3} \right) + i O_{d \rightarrow \infty} \left(\frac{1}{d^{2-\varepsilon}} \right) \right) V_{-1}, \\ \partial_{x_2}V_{-1} &= \left(O_{d \rightarrow \infty} \left(\frac{1}{d^{4-\varepsilon}} \right) + i O_{d \rightarrow \infty} \left(\frac{1}{d} \right) \right) V_{-1}, \\ \partial_{x_1 x_1}V_{-1} &= \left(O_{d \rightarrow \infty} \left(\frac{1}{d^{4-2\varepsilon}} \right) + i O_{d \rightarrow \infty} \left(\frac{1}{d^{3-\varepsilon}} \right) \right) V_{-1}, \\ \partial_{x_1 x_2}V_{-1} &= \left(O_{d \rightarrow \infty} \left(\frac{1}{d^{3-\varepsilon}} \right) + \frac{i}{4d^2} \left(1 + O_{d \rightarrow \infty} \left(\frac{1}{d^{1-\varepsilon}} \right) \right) \right) V_{-1}. \end{aligned}$$

4.3 Estimate on $\partial_{z_1} \Psi_{c,d}$ in $B(d\vec{e}_1, d^{\varepsilon'})$

We define the following norms for $\Psi = \Psi_1 + i\Psi_2$ and $h = h_1 + ih_2$, $0 < \alpha < 1, 0 < \varepsilon' < \varepsilon < 1$:

$$\begin{aligned} \|\Psi\|_{*,B'_d} &:= \|V\Psi\|_{C^1(\{r_1 \leq 2\})} \\ &+ \|r_1^{1-\alpha}\Psi_1\|_{L^\infty(\{d^{\varepsilon'} \geq r_1 \geq 2\})} + \|r_1^{1-\alpha}\nabla\Psi_1\|_{L^\infty(\{d^{\varepsilon'} \geq r_1 \geq 2\})} \\ &+ \|r_1^{-\alpha}\Psi_2\|_{L^\infty(\{d^{\varepsilon'} \geq r_1 \geq 2\})} + \|r_1^{1-\alpha}\nabla\Psi_2\|_{L^\infty(\{d^{\varepsilon'} \geq r_1 \geq 2\})} \end{aligned}$$

and

$$\begin{aligned} \|h\|_{**,B_d} &:= \|Vh\|_{C^0(\{r_1 \leq 3\})} \\ &+ \|r_1^{1-\alpha}h_1\|_{L^\infty(\{d^\varepsilon \geq r_1 \geq 2\})} + \|r_1^{2-\alpha}h_2\|_{L^\infty(\{d^\varepsilon \geq r_1 \geq 2\})}. \end{aligned}$$

They are the norms $\|\cdot\|_{*,-\alpha,d}$ and $\|\cdot\|_{**, -\alpha,d}$ of subsection 2.3, but without the second derivatives, less decay on the gradient of the real part for $\|\cdot\|_{*,B'_d}$, and only on $B'_d = B(d\vec{e}_1, d^{\varepsilon'})$ for $\|\cdot\|_{*,B'_d}$ and on $B_d := B(d\vec{e}_1, d^\varepsilon)$ for $\|\cdot\|_{**,B_d}$. The other main difference with the previous norms is that we require less decay (we take $-\alpha < 0$ instead of $\sigma > 0$ in the decay) in space, which here, since the norms are only in $\{r_1 \leq d^\varepsilon\}$, can be compensated by some smallness in c .

From Corollary 4.3, we have that $\|\partial_{z_1} \Psi_{c,d_c}\|_{*,B'_{d_c}} < +\infty$. We want to show the following proposition.

Proposition 4.5 *For $0 < \alpha < 1, 0 < \varepsilon' < \varepsilon < 1, 0 < \lambda < 1$, if*

$$\lambda < (1 + \alpha)\varepsilon',$$

$$\lambda + (1 - \alpha)\varepsilon' < 2\varepsilon - \varepsilon'$$

and

$$\lambda < 2 - \varepsilon(2 - \alpha),$$

we have

$$\|\partial_{z_1} \Psi_{c,d|d=d_c}\|_{*,B'_{d_c}} = o_{c \rightarrow 0}(c^{1+\lambda}).$$

Such a choice of parameters $(\lambda, \alpha, \varepsilon, \varepsilon')$ exists, we can take for instance $\alpha = 1/2, \lambda = 3/4, \varepsilon = 19/24$ and $\varepsilon' = 13/24$. Furthermore, with this particular choice of parameters, we also have

$$\lambda + (1 - \alpha)\varepsilon' > 1, \tag{4.2}$$

which will be useful later on. These conditions are bounds on how much additional smallness we can have on $\partial_{z_1} \Psi_{c,d}$ near $d_c\vec{e}_1$.

The main goal of this proposition is to have a decay in c better than $O_{c \rightarrow 0}(c)$, which is not obvious from the estimates we have done until now. The estimate on $\partial_{z_1} \Psi_{c,d|d=d_c}$ from Corollary 4.3 will not be enough in the computation of $\partial_c d_c$ for the nonlinear terms. The proof of Proposition 4.5 follows closely the proof of the invertibility of the linearized operator in Proposition 2.17. We want to invert the same linearized operator, but with a different norm, which is better locally around the vortex V_1 .

The reason why we take B_d a little bigger than B'_d is to make the elliptic estimates of step 2 in Proposition 2.17 work here too. The main idea of this proposition is to show that if we move V_{-1} a little, then locally around V_1 the change is very small. We now start the proof of Proposition 4.5.

Proof First, we remark that in B_d , since $\varepsilon < 1, \tilde{r} = r_1$.

Step 1. Computation of the equation on $\partial_{z_1} \Psi_{c,d}$.

Recall that $\Phi_{c,d}$ solves the equation (with $\Phi_{c,d} = V\Psi_{c,d}$)

$$\eta L(\Phi_{c,d}) + (1 - \eta)VL'(\Psi_{c,d}) + F(\Psi_{c,d}) = \lambda(c, d)Z_d,$$

and we recall that $\lambda(c, d) = \frac{\langle F(\Psi_{c,d}), Z_d \rangle}{\|Z_d\|_{L^2(\mathbb{R}^2)}^2}$, and we check easily, with Lemma 3.3, that it is a C^1 function of d . The equation on $\Phi_{c,d}$ holds for any $x \in \mathbb{R}^2$ and any $d \in \mathbb{R}$, $\frac{1}{2d} < c < \frac{2}{d}$, hence

$$\partial_{z_1}(\eta L(\Phi_{c,d}) + (1 - \eta)V L'(\Psi_{c,d}) + \Pi_d^\perp(F(\Psi_{c,d})) - \lambda(c, d)Z_d) = 0.$$

We compute

$$\begin{aligned} \partial_{z_1}(\lambda(c, d)Z_d) &= (\partial_{x_1} + \partial_d)(\lambda(c, d)Z_d) \\ &= \partial_d \lambda(c, d)Z_d + \lambda(c, d)\partial_{z_1}Z_d, \end{aligned}$$

and we recall, from the proof of Proposition 2.26 that

$$\lambda(c, d) \int_{\mathbb{R}^2} |\partial_d V|^2 \eta^2 = \pi \left(\frac{1}{d} - c \right) + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

With Lemma 3.3 and Corollary 4.3, as well as Lemma 2.6, we infer that the terms contributing to the $O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$ are such that, when differentiated with respect to d , their contributions are still a $O_{c \rightarrow 0}^\sigma(c^{2-\sigma})$. Indeed, if the derivative with respect to d fall on a $\Psi_{c,d}$, then by Lemma 3.3 and Corollary 4.3, the same estimates used in the proof of Proposition 2.26 still hold. If the derivative fall on a term depending on V , by Lemma 2.6, we gain some decay in the integrals. We deduce that, since $\lambda(c, d_c) = 0$,

$$\partial_d \lambda(c, d)|_{d=d_c} = \frac{-\pi}{d_c^2} + O_{c \rightarrow 0}^\sigma(c^{2-\sigma}) = O_{c \rightarrow 0}^\sigma(c^{2-\sigma}).$$

Here, we see why the fact that d is differentiable with respect to c is not obvious. The main contribution is at this point not enough to beat the error terms. Therefore, showing that $\partial_d \lambda(c, d) \neq 0$ is not simple here. This is why we need improved estimations on $\partial_{z_1} \Psi_{c,d_c}$, that will give us the fact that the error terms are a $O_{c \rightarrow 0}^\varepsilon(c^{2+\varepsilon})$ for some $\varepsilon > 0$.

Now, writing

$$\text{TW}_c(Q_{c,d}) = \eta L(\Phi_{c,d}) + (1 - \eta)V L'(\Psi_{c,d}) + F(\Psi_{c,d}),$$

(with the notations of Lemma 2.7), we have (since $\lambda(c, d_c) = 0$)

$$(\partial_{z_1}(\text{TW}_c(Q_{c,d})) - \partial_d \lambda(c, d)Z_d)|_{d=d_c} = 0.$$

We recall that

$$F(\Psi_{c,d}) = E - ic\partial_{x_2}V + V(1 - \eta)(-\nabla \Psi_{c,d} \cdot \nabla \Psi_{c,d} + |V|^2 S(\Psi_{c,d})) + R(\Psi_{c,d}),$$

where $R(\Psi_{c,d})$ is a sum of terms at least quadratic in $\Psi_{c,d}$ or $\Phi_{c,d}$ localized in the area where $\eta \neq 0$.

We compute

$$\begin{aligned} \partial_{z_1}(\text{TW}_c(Q_{c,d})) &= \eta L(V\partial_{z_1}\Psi_{c,d}) + (1 - \eta)V L'(\partial_{z_1}\Psi_{c,d}) \\ &+ \eta \partial_{z_1}L(\Phi_{c,d}) + (1 - \eta)V \partial_{z_1}L'(\Psi_{c,d}) + \partial_{z_1}(E - ic\partial_{x_2}V) \\ &+ \eta L(\partial_{z_1}V\Psi_{c,d}) + (1 - \eta)\partial_{z_1}V L'(\Psi_{c,d}) \\ &+ \partial_{z_1}\eta(L(\Phi_{c,d}) - V L'(\Psi_{c,d}) - ic\partial_{x_2}\Phi_{c,d}) \\ &- \partial_{z_1}\eta V(-ic\partial_{x_2}\Psi_{c,d} - \nabla \Psi_{c,d} \cdot \nabla \Psi_{c,d} + |V|^2 S(\Psi_{c,d})) \\ &+ \partial_{z_1}(R(\Psi_{c,d})) \\ &+ \partial_{z_1}V(1 - \eta)(-ic\partial_{x_2}\Psi_{c,d} - \nabla \Psi_{c,d} \cdot \nabla \Psi_{c,d} + |V|^2 S(\Psi_{c,d})) \\ &+ V(1 - \eta)\partial_{z_1}(-ic\partial_{x_2}\Psi_{c,d} - \nabla \Psi_{c,d} \cdot \nabla \Psi_{c,d} + |V|^2 S(\Psi_{c,d})). \end{aligned}$$

We regroup the terms in the following way. We define

$$\mathcal{L}(\partial_{z_1}\Psi_{c,d}) := \eta L(V\partial_{z_1}\Psi_{c,d}) + (1 - \eta)V L'(\partial_{z_1}\Psi_{c,d}),$$

which is the same linearized operator we have inverted in Proposition 2.17 (taken in $\partial_{z_1}\Psi_{c,d}$), and we define the operator

$$\mathcal{L}_{\partial_{z_1}}(\Psi_{c,d}) := \eta\partial_{z_1}L(\Phi_{c,d}) + (1-\eta)V\partial_{z_1}L'(\Psi_{c,d}) + \eta L(\partial_{z_1}V\Psi_{c,d}) + (1-\eta)\partial_{z_1}VL'(\Psi_{c,d}).$$

We already have shown that $\text{TW}_c(V) = E - ic\partial_{x_2}V$, therefore

$$\partial_{z_1}(\text{TW}_c(V)) = \partial_{z_1}(E - ic\partial_{x_2}V).$$

We define the local error

$$\text{Err}_{\text{loc}} := \partial_{z_1}(R(\Psi_{c,d})) - \partial_d\lambda(c,d)Z_d,$$

the far away error

$$\text{Err}_{\text{far}} := \partial_{z_1}V(1-\eta)(-\nabla\Psi_{c,d}\cdot\nabla\Psi_{c,d} + |V|^2S(\Psi))$$

and the nonlinear terms

$$\text{NL}_{\partial_{z_1}}(\Psi_{c,d}) := V(1-\eta)\partial_{z_1}(-\nabla\Psi_{c,d}\cdot\nabla\Psi_{c,d} + |V|^2S(\Psi_{c,d})).$$

Finally, we write the cutoff error

$$\text{Err}_{\text{cut}} := \partial_{z_1}\eta(L(\Phi_{c,d}) - VL'(\Psi_{c,d}) + ic\partial_{x_2}\Psi_{c,d} + \nabla\Psi_{c,d}\cdot\nabla\Psi_{c,d} - |V|^2S(\Psi_{c,d}))$$

which is supported in the area $\{2 \leq r_{-1} \leq 3\}$, and in particular is zero in $B(d_c\vec{e}_1^\lambda, d_c^\varepsilon)$. With these definitions, we have, at $d = d_c$,

$$\begin{aligned} & (\partial_{z_1}(\eta L(\Phi_{c,d}) + (1-\eta)VL'(\Psi_{c,d}) + F(\Psi_{c,d})) - \partial_d\lambda(c,d)Z_d)|_{d=d_c} \\ &= \mathcal{L}(\partial_{z_1}\Psi_{c,d})|_{d=d_c} \\ &+ (\partial_{z_1}(\text{TW}_c(V)) + \mathcal{L}_{\partial_{z_1}}(\Psi_{c,d}) + \text{NL}_{\partial_{z_1}}(\Psi_{c,d}))|_{d=d_c} \\ &+ (\text{Err}_{\text{loc}} + \text{Err}_{\text{far}} + \text{Err}_{\text{cut}})|_{d=d_c}. \end{aligned}$$

The equation satisfied by $\partial_{z_1}\Psi_{c,d}$ at $d = d_c$ is therefore

$$(\mathcal{L}(\partial_{z_1}\Psi_{c,d}) + \partial_{z_1}(\text{TW}_c(V)) + \mathcal{L}_{\partial_{z_1}}(\Psi_{c,d}) + \text{NL}_{\partial_{z_1}}(\Psi_{c,d}) + \text{Err}_{\text{loc}} + \text{Err}_{\text{far}} + \text{Err}_{\text{cut}})|_{d=d_c} = 0.$$

Step 2. Beginning of the contradiction argument.

Now, suppose that the result of Proposition 4.5 is false. The scheme of this proof is the same as in Proposition 2.17. Then, there exist an absolute constant $\delta > 0$ and sequences $\partial_{z_1}\Psi_n$, $c_n \rightarrow 0$, $d_n \rightarrow \infty$ such that

$$d_n^{1+\lambda}\|\partial_{z_1}\Psi_n|_{d=d_n}\|_{*,B'_{d_n}} \geq \delta,$$

where we write $d_n = d_{c_n}$ (a value such that $\lambda(c_n, d_n) = 0$ in Proposition 2.26). We have just shown that Ψ_n (where we omit the subscripts in d_n, c_n) satisfies

$$\mathcal{L}(\partial_{z_1}\Psi_n) + \partial_{z_1}(\text{TW}_{c_n}(V)) + \mathcal{L}_{\partial_{z_1}}(\Psi_n) + \text{NL}_{\partial_{z_1}}(\Psi_n) + \text{Err}_{\text{loc}} + \text{Err}_{\text{far}} + \text{Err}_{\text{cut}} = 0.$$

The function

$$\frac{(V\partial_{z_1}\Psi_n)(\cdot - d_n\vec{e}_1^\lambda)}{\|\partial_{z_1}\Psi_n\|_{*,B'_{d_n}}}$$

converges locally uniformly up to a subsequence to a limit \mathfrak{G} , since it is bounded in $\|\cdot\|_{*,B'_\lambda}$ for any $\lambda > 0$ (for the same reasons that $\Psi_n \rightarrow \Psi$ locally uniformly in the beginning of the proof of Proposition 2.17).

The equation on $\partial_{z_1}\Psi_n$ is

$$\mathcal{L}(\partial_{z_1}\Psi_n) + Vh_n = 0, \tag{4.3}$$

with

$$Vh_n := \partial_{z_1}(\text{TW}_{c_n}(V)) + \mathcal{L}_{\partial_{z_1}}(\Psi_n) + \text{NL}_{\partial_{z_1}}(\Psi_n) + \text{Err}_{\text{loc}} + \text{Err}_{\text{far}} + \text{Err}_{\text{cut}}.$$

The goal of Proposition 2.17 was to estimate $\|\Psi\|_{*,\sigma,d}$ with $\|h\|_{**,\sigma',d}$ for the equation $\mathcal{L}(\Psi) = h$ if d is large enough (given an orthogonality condition on Ψ). Here we do the same thing, but localized in space, and with a very particular h_n that we will estimate. To continue as in the proof of Proposition 2.17, we want to show that

$$\frac{h_n(\cdot - d_n \vec{e}_1)}{\|\partial_{z_1} \Psi_n\|_{*,B'_{d_n}}} \rightarrow 0$$

in C_{loc}^0 so that we get at the limit (following the +1 vortex) in (4.3)

$$L_{V_1}(\mathfrak{G}) = 0,$$

using the same techniques as in the proof of Proposition 2.17. It will be enough for that to show that

$$\left\| \frac{h_n}{\|\partial_{z_1} \Psi_n\|_{*,B'_{d_n}}} \right\|_{**,B_{d_n}} \rightarrow 0 \quad (4.4)$$

and we will also use this estimate later on. Remark that here, the problem is no longer symmetric in x_1 , in particular, we cannot use the same argument near the -1 vortex, but it is not needed.

Step 3. Proof of (4.4).

Recall the definition of $\|\cdot\|_{**,B_{d_n}}$:

$$\|h\|_{**,B_{d_n}} = \|Vh\|_{C^0(\{r_1 \leq 3\})} + \|r_1^{1-\alpha} h_1\|_{L^\infty(\{d_n^\varepsilon \geq r_1 \geq 2\})} + \|r_1^{2-\alpha} h_2\|_{L^\infty(\{d_n^\varepsilon \geq r_1 \geq 2\})}.$$

Since

$$d_n^{1+\lambda} \|\partial_{z_1} \Psi_n|_{d=d_n}\|_{*,B'_{d_n}} \geq \delta,$$

we have

$$\frac{1}{\|\partial_{z_1} \Psi_n\|_{*,B'_{d_n}}} \leq \frac{1}{\delta c_n^{1+\lambda}},$$

therefore it is enough to show that

$$\|h_n\|_{**,B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}) \quad (4.5)$$

to have (4.4). We recall that

$$Vh_n = \partial_{z_1}(\text{TW}_{c_n}(V)) + \mathcal{L}_{\partial_{z_1}}(\Psi_n) + \text{NL}_{\partial_{z_1}}(\Psi_n) + \text{Err}_{\text{loc}} + \text{Err}_{\text{far}} + \text{Err}_{\text{cut}}.$$

The contribution of $\partial_{z_1}(\text{TW}_{c_n}(V))$ will be established in step 3.1, $\mathcal{L}_{\partial_{z_1}}(\Psi_n)$ in step 3.2, $\text{NL}_{\partial_{z_1}}(\Psi_n)$ in step 3.3, and finally, $\text{Err}_{\text{loc}} + \text{Err}_{\text{far}} + \text{Err}_{\text{cut}}$ in step 3.4.

Step 3.1. Proof of $\left\| \frac{\partial_{z_1} \text{TW}_{c_n}(V)}{V} \right\|_{**,B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda})$.

Recall from (2.2) that

$$\text{TW}_c(V) = E - ic\partial_{x_2}V = -2\nabla V_1 \cdot \nabla V_{-1} + (1 - |V_1|^2)(1 - |V_{-1}|^2)V_1V_{-1} - ic\partial_{x_2}V,$$

therefore, with Lemma 4.2, we have

$$\partial_{z_1}(\text{TW}_c(V)) = -4\nabla V_1 \cdot \nabla \partial_{x_1}V_{-1} + 2(1 - |V_1|^2)V_1\partial_{x_1}((1 - |V_{-1}|^2)V_{-1}) - 2ic\partial_{x_2}(V_1\partial_{x_1}V_{-1}).$$

We now estimate this quantity at $d = d_n$. We have, in $\{r_1 \leq d_n^\varepsilon\}$,

$$|(1 - |V_1|^2)V_1\partial_{x_1}((1 - |V_{-1}|^2)V_{-1})| \leq \frac{K}{1 + r_1^2} \times \frac{1}{d_n^3},$$

and using $\lambda < 1$, $\alpha > 0$, we deduce

$$\left\| \frac{(1 - |V_1|^2)V_1\partial_{x_1}((1 - |V_{-1}|^2)V_{-1})}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

We compute with Lemmas 2.2 and 4.4 that

$$\Re \left(\frac{4\nabla V_1 \cdot \nabla \partial_{x_1} V_{-1}}{V} \right) = 4\Re \left(\frac{\nabla V_1}{V_1} \right) \cdot \Re \left(\frac{\nabla \partial_{x_1} V_{-1}}{V_{-1}} \right) - 4\Im \left(\frac{\nabla V_1}{V_1} \right) \cdot \Im \left(\frac{\nabla \partial_{x_1} V_{-1}}{V_{-1}} \right),$$

leading to

$$|V| \left| \Re \left(\frac{4\nabla V_1 \cdot \nabla \partial_{x_1} V_{-1}}{V} \right) \right| \leq \frac{K}{(1 + r_1^3)d_n^{3-\varepsilon}} + \frac{K}{(1 + r_1)d_n^2}$$

in $\{r_1 \leq d_n^\varepsilon\}$ for a universal constant K . Since $\lambda < 1$ and $\alpha > 0$, we have

$$\left\| \Re \left(\frac{4\nabla V_1 \cdot \nabla \partial_{x_1} V_{-1}}{V} \right) \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

Similarly, we have in $\{r_1 \leq d_n^\varepsilon\}$ that

$$|V| \left| \Im \left(\frac{4\nabla V_1 \cdot \nabla \partial_{x_1} V_{-1}}{V} \right) \right| \leq \frac{K}{(1 + r_1^3)d_n^2} + \frac{K}{(1 + r_1)d_n^{3-\varepsilon}}.$$

Therefore, using

$$\frac{1}{d_n} \leq \frac{K}{(1 + r_1)^{1/\varepsilon}},$$

since we are in $B_{d_n} = B(d_n \vec{e}_1, d_n^\varepsilon)$, and

$$\lambda < 2 - \varepsilon(2 - \alpha),$$

which is one of the hypothesis of the lemma, we have

$$\left\| i\Im \left(\frac{4\nabla V_1 \cdot \nabla \partial_{x_1} V_{-1}}{V} \right) \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

Now, for $2ic_n\partial_{x_2}(V_1\partial_{x_1}V_{-1}) = 2ic_n\partial_{x_2}V_1\partial_{x_1}V_{-1} + 2ic_n\partial_{x_1x_2}V_{-1}V_1$, we estimate similarly (still using Lemma 2.2 and 4.4)

$$|V| \left| \Re \left(\frac{ic_n\partial_{x_2}V_1\partial_{x_1}V_{-1}}{V} \right) \right| \leq \frac{K}{(1 + r_1^3)d_n^{3-\varepsilon}} + \frac{K}{(1 + r_1)d_n^4},$$

$$|V| \left| \Im \left(\frac{ic_n\partial_{x_2}V_1\partial_{x_1}V_{-1}}{V} \right) \right| \leq \frac{K}{(1 + r_1^3)d_n^4} + \frac{K}{(1 + r_1)d_n^{3-\varepsilon}},$$

therefore, using $\frac{1}{d_n} \leq \frac{K}{(1+r_1)^{1/\varepsilon}}$, we have, under the condition

$$\lambda < 2 - \varepsilon(2 - \alpha)$$

for the imaginary part (as for the previous term) and with no condition for the real part (since $\alpha > 0, \lambda < 1$), that

$$\left\| \frac{2ic_n\partial_{x_2}V_1\partial_{x_1}V_{-1}}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

We then compute (still using Lemma 2.2 and 4.4)

$$\left| \Re \left(\frac{ic_n\partial_{x_1x_2}V_{-1}V_1}{V} \right) \right| \leq \frac{K}{d_n^3},$$

$$\left| \Im \left(\frac{ic_n \partial_{x_1 x_2} V_{-1} V_1}{V} \right) \right| \leq \frac{K}{d_n^{4-\varepsilon}},$$

therefore, using $\frac{1}{d_n} \leq \frac{K}{(1+r_1)^{1/\varepsilon}}$, we have, under the conditions

$$\lambda < 2 - \varepsilon(1 - \alpha) \quad \text{and} \quad \lambda < 3 - \varepsilon(3 - \alpha),$$

which are met since

$$\lambda < 2 - \varepsilon(2 - \alpha) = 2 - \varepsilon(1 - \alpha) - \varepsilon < 2 - \varepsilon(1 - \alpha),$$

and $\lambda < 2 - \varepsilon(2 - \alpha) = 3 - \varepsilon(3 - \alpha) - 1 + \varepsilon < 3 - \varepsilon(3 - \alpha)$, that

$$\left\| \frac{ic \partial_{x_2} (V_1 \partial_{x_1 x_2} V_{-1})}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

This concludes the proof of step 3.1.

Step 3.2. Proof of $\left\| \frac{\mathcal{L}_{\partial_{z_1}}(\Psi_n)}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda})$.

We have defined

$$\mathcal{L}_{\partial_{z_1}}(\Psi_n) = \eta(\partial_{z_1} L)(\Phi_n) + (1 - \eta)V(\partial_{z_1} L')(\Psi_n) + \eta L((\partial_{z_1} V)\Psi_n) + (1 - \eta)\partial_{z_1} V L'(\Psi_n).$$

We recall from Lemma 2.7 that

$$L'(\Psi_n) = -\Delta \Psi_n - 2 \frac{\nabla V}{V} \cdot \nabla \Psi_n + 2|V|^2 \Re(\Psi_n) - ic_n \partial_{x_2} \Psi_n,$$

$$L(\Phi_n) = -\Delta \Phi_n - (1 - |V|^2)\Phi_n + 2\Re(\bar{V}\Phi_n)V - ic_n \partial_{x_2} \Phi_n,$$

hence

$$(\partial_{z_1} L)(\Phi_n) = 4\Re(\bar{V}_{-1} \partial_{x_1} V_{-1})\Phi_n + 4\Re(\bar{\partial_{x_1} V_{-1} V_1} \Phi_n)V + 4\Re(\bar{V}\Phi_n)V_1 \partial_{x_1} V_{-1}.$$

We shall now estimate all these terms one by one.

Since $\eta \partial_{z_1} L(\Phi_n)$ is compactly supported in $\{\tilde{r} \leq 2\}$ and $\|\cdot\|_{**, B_{d_n}}$ looks at the function only on $\{r_1 \leq d^\varepsilon\}$, using Lemma 4.4 ($\nabla V_{-1} = O_{c \rightarrow 0}(c)$) and $\|\Psi_n\|_{*, \frac{1-\lambda}{4}, d_n} \leq K(\lambda)c^{\frac{1+\lambda}{2}}$, we check that

$$\left\| \frac{\eta \partial_{z_1} L(\Phi_n)}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

With the same arguments, we also check that

$$\left\| \frac{\eta L(\partial_{z_1} V \Psi_n)}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

Now, with $\|\Psi_n\|_{*, \sigma, d_n} \leq K(\sigma, \sigma')c_n^{1-\sigma'}$, we check that for any $0 < \sigma < \sigma' < 1$,

$$|L'(\Psi_n)| \leq \frac{K(\sigma, \sigma')}{(1+r_1)^{1+\sigma} d_n^{1-\sigma'}},$$

therefore, with Lemma 4.4, we have

$$|(1 - \eta)\partial_{z_1} V L'(\Psi_n)| \leq \frac{K(\sigma, \sigma')}{(1+r_1)^{1+\sigma} d_n^{3-\varepsilon-\sigma'}}.$$

In particular, we check that if

$$\lambda < 2 - \varepsilon(2 - \alpha),$$

we can take σ, σ' such that $0 < \sigma < \sigma' < \frac{2-\varepsilon(2-\alpha)-\lambda}{1-\varepsilon}$, hence

$$\left\| \frac{(1-\eta)\partial_{z_1} V L'(\Psi_n)}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

Finally, we estimate

$$|\partial_{z_1} L'(\Psi_n)| \leq K \left| \partial_{x_1} \frac{\nabla V_{-1}}{V_{-1}} \cdot \nabla \Psi_n \right| + K |\Re(\partial_{x_1} V_{-1} \overline{V_{-1}}) \Re(\Psi_n)|.$$

With Lemma 4.4 and $\|\Psi_n\|_{*, \sigma, d_n} \leq K(\sigma, \sigma') c_n^{1-\sigma}$ (from (3.9)), we check that

$$|(1-\eta)V\partial_{z_1} L'(\Psi_n)| \leq \frac{K(\sigma, \sigma')(1-\eta)}{r_1^{1+\sigma} d_n^{4-\varepsilon-\sigma'}},$$

therefore, with the same condition as for the previous term, namely

$$\lambda < 2 - (2 - \alpha)\varepsilon,$$

we infer, taking $\sigma < \sigma'$ small enough,

$$\left\| \frac{(1-\eta)V\partial_{z_1} L'(\Psi_n)}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

This concludes the proof of step 3.2.

Step 3.3. Proof of $\left\| \frac{\text{NL}_{\partial_{z_1}}(\Psi_n)}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda})$.

We recall

$$\text{NL}_{\partial_{z_1}}(\Psi_n) = V(1-\eta)\partial_{z_1}(-\nabla \Psi_n \cdot \nabla \Psi_n + |V|^2 S(\Psi_n)),$$

with $S(\Psi_n) = e^{2\Re(\Psi_n)} - 1 - 2\Re(\Psi_n)$. We compute

$$\begin{aligned} \partial_{z_1}(-\nabla \Psi_n \cdot \nabla \Psi_n + |V|^2 S(\Psi_n)) &= -2\nabla \partial_{z_1} \Psi_n \cdot \nabla \Psi_n \\ &\quad + 4\Re(\partial_{x_1} V_{-1} \overline{V_{-1}}) S(\Psi_n) \\ &\quad + |V|^2 \partial_{z_1} S(\Psi_n). \end{aligned}$$

Now, with Corollary 4.3 and (3.9), we check that, for any $0 < \sigma < \sigma' < 1$, $r_1 \geq 2$,

$$|\nabla \partial_{z_1} \Psi_n \cdot \nabla \Psi_n| \leq \frac{K(\sigma, \sigma')}{r_1^{2+2\sigma} d_n^{2-2\sigma'}},$$

$$|4\Re(\partial_{x_1} V_{-1} \overline{V_{-1}}) S(\Psi_n) + |V|^2 \partial_{z_1} S(\Psi_n)| \leq \frac{K(\sigma, \sigma')}{r_1^{2+2\sigma} d_n^{2-2\sigma'}},$$

therefore, taking $\sigma < \sigma' < \frac{1-\lambda}{2}$, we check that

$$\|(1-\eta)(-2\nabla \partial_{z_1} \Psi_n \cdot \nabla \Psi_n + 4\Re(\partial_{x_1} V_{-1} \overline{V_{-1}}) S(\Psi_n) + |V|^2 \partial_{z_1} S(\Psi_n))\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

The proof of step 3.3 is complete.

Step 3.4. Proof of $\left\| \frac{\text{Err}_{\text{loc}} + \text{Err}_{\text{far}} + \text{Err}_{\text{cut}}}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda})$.

We recall

$$\text{Err}_{\text{cut}} = \partial_{z_1} \eta(L(\Phi_n) - V L'(\Psi_n) + ic \partial_{x_2} \Psi_n + \nabla \Psi_n \cdot \nabla \Psi_n - |V|^2 S(\Psi_n)),$$

$$\text{Err}_{\text{loc}} = \partial_{z_1}(R(\Psi_n)) - \partial_d \lambda(c_n, d_n) Z_{d_n},$$

$$\text{Err}_{\text{far}} = \partial_{z_1} V(1-\eta)(-\nabla \Psi_n \cdot \nabla \Psi_n + |V|^2 S(\Psi_n)).$$

Err_{cut} is compactly supported in $\{r_{-1} \leq 2\}$, therefore $\text{Err}_{\text{cut}} = 0$ in B_{d_n} , hence

$$\left\| \frac{\text{Err}_{\text{cut}}}{V} \right\|_{**, B_{d_n}} = 0.$$

Now, Err_{loc} is supported in $\{r_1 \leq 2\}$, and from Lemma 2.7, we know that $R(\Psi_n)$ is a sum of terms at least quadratic in Ψ_n or Φ_n localized in the area where $\eta \neq 0$. Therefore, from Corollary 4.3 and (3.10), we check that

$$|\partial_{z_1}(R(\Psi_n))| \leq \frac{K(\sigma)}{d_n^{2-2\sigma}},$$

and we have check in step 1 that $|\partial_d \lambda(c_n, d_n)| = O_{c_n \rightarrow 0}^\sigma(c_n^{2-\sigma})$. Thus, taking $\sigma < \frac{1-\lambda}{2}$,

$$\left\| \frac{\text{Err}_{\text{loc}}}{V} \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

From (3.9), we check that, for any $1 > \sigma' > \sigma > 0$, in $\{r_1 \leq d_n^\varepsilon\}$,

$$|-\nabla \Psi_n \cdot \nabla \Psi_n + |V|^2 S(\Psi_n)| \leq \frac{K(\sigma, \sigma')}{(1+r_1)^{2+2\sigma} d_n^{2-2\sigma'}},$$

and from Lemma 4.4, we have there

$$|\partial_{z_1} V| \leq \frac{K}{d_n^{2-\varepsilon}},$$

therefore, choosing $\sigma < \sigma'$ small enough, we have

$$\left\| \frac{\partial_{z_1} V}{V} (1-\eta)(-\nabla \Psi_n \cdot \nabla \Psi_n + |V|^2 S(\Psi_n)) \right\|_{**, B_{d_n}} = o_{c_n \rightarrow 0}(c_n^{1+\lambda}).$$

This ends the proof of step 3.4 and hence of (4.4).

Step 4. Three additional estimates on h_n .

We infer that we can complete estimates done in the previous step to show following three estimates:

$$\|V h_n\|_{L^\infty(\{\tilde{r} \leq 3\})} + \|\tilde{r}^{1+\sigma} \Re(h_n)\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma} \Im(h_n)\|_{L^\infty(\{\tilde{r} \geq 2\})} \leq K(\sigma, \sigma') c_n^{1-\sigma'}. \quad (4.6)$$

In the right half-plane,

$$|h_n| \leq \frac{K(\sigma) c_n^{1+\sigma}}{(1+r_1)}, \quad (4.7)$$

and, in the left half-plane,

$$|h_n| \leq \frac{K(\sigma) c_n^{1-\sigma}}{(1+r_{-1})^2}. \quad (4.8)$$

Observe that h_n is not symmetrical with respect to x_1 because of the cutoff. Recall that

$$V h_n = \partial_{z_1}(\text{TW}_{c_n}(V)) + \mathcal{L}_{\partial_{z_1}}(\Psi_n) + \text{NL}_{\partial_{z_1}}(\Psi_n) + \text{Err}_{\text{loc}} + \text{Err}_{\text{far}} + \text{Err}_{\text{cut}}.$$

Step 5. Inner estimates.

By the estimation we have just proved, we have in particular

$$\frac{h_n(\cdot - d_n \vec{e}_1)}{\|\partial_{z_1} \Psi_n\|_{**, B'_{d_n}}} \rightarrow 0$$

in C_{loc}^0 (which corresponds to follow the +1 vortex). Therefore, at the limit, in the distribution sense,

$$L_{V_1}(\mathfrak{G}) = 0$$

in all \mathbb{R}^2 . If we show that $\langle \mathfrak{G}, \chi \partial_{x_1} V_1 \rangle = 0$ for χ a cutoff near 0, we can then use Theorem 2.16 to show, similarly as in the proof of Proposition 2.17, that $\mathfrak{G} = 0$ since

$$\left\| \frac{(V \partial_{z_1} \Psi_n)(\cdot - d_n \vec{e}_1)}{\|\partial_{z_1} \Psi_n\|_{*, B'_{d_n}}} \right\|_{*, B_{d_n}} = 1,$$

hence $\|\mathfrak{G}\|_{H_{V_1}} < +\infty$. We recall that, by construction, we have $\langle \Phi_{c,d}, Z_d \rangle = 0$. By symmetry, this implies that $\langle \Phi_{c,d}, \eta(y) \partial_d V \rangle = 0$. Both $\Phi_{c,d}$ and $\eta(y) \partial_d V$ are C^1 with respect to d , and therefore

$$0 = \partial_d \langle \Phi_{c,d}, \eta(y) \partial_d V \rangle = \langle \partial_d \Phi_{c,d}, \eta(y) \partial_d V \rangle + \langle \Phi_{c,d}, \partial_d (\eta(y) \partial_d V) \rangle.$$

Furthermore, $\langle \partial_{x_1} \Phi_{c,d}, \eta(y) \partial_d V \rangle = -\langle \Phi_{c,d}, \partial_{x_1} \eta(y) \partial_d V \rangle$, thus

$$\langle \partial_{z_1} \Phi_{c,d}, \eta(y) \partial_d V \rangle = -\langle \Phi_{c,d}, \eta(y) \partial_{z_1} \partial_d V \rangle,$$

and we check easily that $|\eta(y) \partial_{z_1} \partial_d V| \leq K c \eta(y)$, therefore, since $\|\Psi_{c,d}\|_{*, \sigma, d} \leq K(\sigma, \sigma') c^{1-\sigma'}$, we have $|\langle \partial_{z_1} \Phi_{c,d}, \eta(y) \partial_d V \rangle| \leq K(\sigma, \sigma') c^{2-\sigma'}$, and thus, taking $0 < \sigma' < 1 - \lambda$, for c_n and d_n , $n \rightarrow \infty$, we infer that $\langle \mathfrak{G}, \eta \partial_{x_1} V_1 \rangle = 0$.

We continue as in the proof of Proposition 2.17. The fact that $\mathfrak{G} = 0$ gives us that for any $R > 0$, we have

$$\frac{\|V \partial_{z_1} \Psi_n\|_{L^\infty(\{r_1 \leq R\})} + \|\nabla(V \partial_{z_1} \Psi_n)\|_{L^\infty(\{r_1 \leq R\})}}{\|\partial_{z_1} \Psi_n\|_{*, B'_{d_n}}} \rightarrow 0.$$

Step 6. Outer computations.

We have the same outer computations as in step 2 of the proof of Proposition 2.17, but with $\mathcal{Y}_n = \frac{\partial_{z_1} \Psi_n}{\|\partial_{z_1} \Psi_n\|_{*, B_{d_n}}}$ playing the role of Ψ_n and $\mathcal{H}_n = \frac{h_n}{\|\partial_{z_1} \Psi_n\|_{*, B_{d_n}}}$ playing the role of h_n , since they satisfy the same equation. We showed in (4.4) that

$$\|\mathcal{H}_n\|_{**, B_{d_n}} = o_{n \rightarrow \infty}(1),$$

and the system of equation is, with $\mathcal{Y}_n = \mathcal{Y}_1 + i\mathcal{Y}_2$ and $\mathcal{H}_n = \mathcal{H}_1 + i\mathcal{H}_2$,

$$\begin{cases} \Delta \mathcal{Y}_1 - 2|V|^2 \mathcal{Y}_1 = -\mathcal{H}_1 - 2\Re\left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n\right) + c \partial_{x_2} \mathcal{Y}_2 \\ \Delta \mathcal{Y}_2 + c \partial_{x_2} \mathcal{Y}_1 = -\mathcal{H}_2 - 2\Im\left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n\right). \end{cases}$$

Recall the two balls $B_{d_n} = B(d_n \vec{e}_1, d_n^\varepsilon)$ and $B'_{d_n} = B(d_n \vec{e}_1, d_n^{\varepsilon'})$. We have, as in the proof of Proposition 2.17, outside $\{r_1 \leq R\}$ but in B'_{d_n} , that $\|\mathcal{Y}_n\|_{*, B'_{d_n}} = 1$ and $\|\mathcal{H}_n\|_{**, B_{d_n}} = o_{n \rightarrow \infty}(1)$, therefore

$$|\Delta \mathcal{Y}_1 - 2\mathcal{Y}_1| \leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1+r_1)^{1-\alpha}} \quad (4.9)$$

and

$$|\Delta \mathcal{Y}_2 + c \partial_{x_2} \mathcal{Y}_1| \leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1+r_1)^{2-\alpha}}. \quad (4.10)$$

We want to extend these estimates in $B_{d_n} = B(d_n \vec{e}_1, d_n^\varepsilon)$ and not only on $B'_{d_n} = B(d_n \vec{e}_1, d_n^{\varepsilon'})$. Since $\|\mathcal{H}_n\|_{**, B_{d_n}} = o_{n \rightarrow \infty}(1)$ from (4.4), the estimates on \mathcal{H}_1 and \mathcal{H}_2 are already on B_{d_n} , leaving $c \partial_{x_2} \mathcal{Y}_2$ and the real and imaginary parts of $\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n$ to estimate.

First, we check that, in $B_{d_n} \setminus B'_{d_n}$,

$$|c_n \partial_{x_2} \mathcal{Y}_2| \leq \frac{d_n^{1+\lambda} c_n^{2-\sigma}}{(1+r_1)^{1+\sigma}} = \frac{o_{n \rightarrow \infty}(1)}{(1+r_1)^{1-\sigma}}$$

taking $\sigma > 0$ small enough. We use $\mathcal{Y}_n = \frac{\partial_{z_1} \Psi_n}{\|\partial_{z_1} \Psi_n\|_{*, B'_{d_n}}}, \frac{1}{\|\partial_{z_1} \Psi_n\|_{*, B'_{d_n}}} \leq K d_n^{1+\lambda}$ and Corollary 4.3 to compute, for any $1 > \sigma > 0$,

$$\left| \Re \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) \right| \leq \left| \frac{\nabla V}{V} \right| \times |\nabla \mathcal{Y}_n| \leq \frac{K(\sigma) d_n^{1+\lambda}}{r_1^{2+\sigma} d_n^{1-\sigma}} \leq \frac{K(\sigma)}{r_1^{2+\sigma} d_n^{-\sigma-\lambda}}.$$

In $B_{d_n} \setminus B'_{d_n}$, we have $r_1 \geq d_n^{\varepsilon'}$, therefore

$$\left| \Re \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) \right| \leq \frac{K(\sigma)}{r_1^{1-\alpha} d_n^{-\sigma-\lambda+(1+\alpha+\sigma)\varepsilon'}}.$$

Since we assume

$$\lambda < (1 + \alpha)\varepsilon',$$

then we can choose $\sigma > 0$ small such that $-\sigma - \lambda + (1 + \alpha + \sigma)\varepsilon' > 0$ and deduce, in $B_{d_n} \setminus B'_{d_n}$, that

$$\left| \Re \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) \right| \leq \frac{o_{n \rightarrow \infty}(1)}{r_1^{1-\alpha}}.$$

This result shows that (4.9) holds on B_{d_n} . Now, we compute

$$\left| \Im \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) \right| \leq \left| \Re \left(\frac{\nabla V}{V} \right) \cdot \Im(\nabla \mathcal{Y}_n) \right| + \left| \Re(\nabla \mathcal{Y}_n) \cdot \Im \left(\frac{\nabla V}{V} \right) \right|,$$

and with Corollary 4.3, Lemma 2.2 and 4.4, we estimate

$$\left| \Re \left(\frac{\nabla V}{V} \right) \cdot \Im(\nabla \mathcal{Y}_n) \right| \leq K(\sigma) \left(\frac{1}{d_n^3} + \frac{1}{r_1^3} \right) \frac{d_n^{1+\lambda}}{r_1^{1+\sigma} d_n^{1-\sigma}}$$

and

$$\left| \Re(\nabla \mathcal{Y}_n) \cdot \Im \left(\frac{\nabla V}{V} \right) \right| \leq K(\sigma) \frac{d_n^{1+\lambda}}{r_1^{2+\sigma} d_n^{1-\sigma}} \left(\frac{1}{d_n^{2-\varepsilon}} + \frac{1}{r_1} \right).$$

In $B_{d_n} \setminus B'_{d_n}$, we have $d_n^{\varepsilon} \geq r_1 \geq d_n^{\varepsilon'}$, and with similar estimates as for the previous term, we check that, since $\lambda < (1 + \alpha)\varepsilon'$, we have

$$\lambda < (2 + \alpha)\varepsilon',$$

for the first term, and

$$\lambda < (1 + \alpha)\varepsilon'$$

for the second one. We can find $\sigma > 0$ such that

$$\left| \Im \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) \right| \leq \frac{o_{n \rightarrow \infty}(1)}{(1 + r_1)^{2-\alpha}}$$

in $B_{d_n} \setminus B'_{d_n}$. We deduce that (4.10) holds on B_{d_n} . Additionally, we will use (from Lemma 3.3) for $0 < \sigma < \sigma' < 1$,

$$\begin{aligned} & \|V \chi \mathcal{Y}_n\|_{C^1(\{\tilde{r} \leq 3\})} \\ & + \|\tilde{r}^{1+\sigma} \Re(\mathcal{Y}_n)\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{1+\sigma} \nabla \Re(\mathcal{Y}_n)\|_{L^\infty(\{\tilde{r} \geq 2\})} \\ & + \|\tilde{r}^\sigma \Im(\mathcal{Y}_n)\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{1+\sigma} \nabla \Im(\mathcal{Y}_n)\|_{L^\infty(\{\tilde{r} \geq 2\})} \\ & \leq K(\sigma, \sigma') c_n^{1-\sigma'} d_n^{1+\lambda} \\ & \leq K(\sigma, \sigma') d_n^{\lambda+\sigma''} \end{aligned} \tag{4.11}$$

and from (4.6),

$$\|V \mathcal{H}_n\|_{L^\infty(\{\tilde{r} \leq 3\})} + \|\tilde{r}^{1+\sigma} \Re(\mathcal{H}_n)\|_{L^\infty(\{\tilde{r} \geq 2\})} + \|\tilde{r}^{2+\sigma} \Im(\mathcal{H}_n)\|_{L^\infty(\{\tilde{r} \geq 2\})} \leq K(\sigma, \sigma') d_n^{\lambda+\sigma''} \tag{4.12}$$

to do estimates outside of B_{d_n} . These estimates are not optimal (in particular in the smallness in c_n) but we will only use them on parts far away from the center of V_1 . Thanks to (4.7), we have a slightly better estimate in the right half-plane, that is, for $0 < \sigma < 1$,

$$|\mathcal{H}_n| \leq K|h_n|d_n^{1+\lambda} \leq \frac{K(\sigma)d_n^{\lambda-\sigma}}{(1+r_1)}. \quad (4.13)$$

Step 7. Elliptic estimates.

At this point, we have on \mathcal{Y}_n that $\|\mathcal{Y}_n\|_{*,B'_{d_n}} = 1$, $\|V\mathcal{Y}_n\|_{L^\infty(\{r_1 \leq R\})} + \|\nabla(V\mathcal{Y}_n)\|_{L^\infty(\{r_1 \leq R\})} \rightarrow 0$ as $n \rightarrow \infty$ for any $R > 1$, and with $\mathcal{Y}_n = \mathcal{Y}_1 + i\mathcal{Y}_2$,

$$\begin{aligned} |\Delta\mathcal{Y}_2 + c\partial_{x_2}\mathcal{Y}_1| &\leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1+r_1)^{2-\alpha}}, \\ |\Delta\mathcal{Y}_1 - 2|V|^2\mathcal{Y}_1| &\leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1+r_1)^{1-\alpha}}. \end{aligned}$$

We want to show that $\|\mathcal{Y}_n\|_{*,B'_{d_n}} = o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)$. We want to use similar elliptic estimates as in the proof of Proposition 2.17, but we have to show that they still work if we only have the estimate in $B_{d_n} = B(d_n \vec{e}_1^\varepsilon, d_n^\varepsilon)$ and we want the final estimates in $B'_{d_n} = B(d_n \vec{e}_1^{\varepsilon'}, d_n^{\varepsilon'})$, with $\varepsilon' < \varepsilon$.

Step 7.1. Elliptic estimate for \mathcal{Y}_2 .

We start by solving the following problem in \mathbb{R}^2 :

$$\Delta\zeta = f,$$

with

$$f := -\mathcal{H}_2 - 2\mathfrak{I}m\left(\frac{\nabla V}{V} \cdot \nabla\mathcal{Y}_n\right),$$

which is odd in x_2 (the derivation with respect to z_1 breaks the symmetry on x_1 , but not on x_2) and satisfies

$$|f| \leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1+r_1)^{2-\alpha}}$$

in $B_{d_n} = B(d_n \vec{e}_1^\varepsilon, d_n^\varepsilon)$, and, from (4.11) and (4.12),

$$|f| \leq \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma'}}{(1+\tilde{r})^{2+\sigma}} \quad (4.14)$$

in \mathbb{R}^2 , for any $1 > \sigma' > \sigma > 0$. We write, for $x \in B(d_n \vec{e}_1^{\varepsilon'}, d_n^{\varepsilon'})$,

$$\nabla\zeta(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{x-Y}{|x-Y|^2} f(Y) dY. \quad (4.15)$$

With the assumption

$$\lambda + \varepsilon'(1-\alpha) < 2\varepsilon - \varepsilon',$$

we check that, for $x \in B(d_n \vec{e}_1^{\varepsilon'}, d_n^{\varepsilon'})$,

$$|\nabla\zeta(x)| \leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1+|x-d_n \vec{e}_1^{\varepsilon'}|)^{1-\alpha}}. \quad (4.16)$$

Using Lemma 2.8 and (4.14), we also have, in all \mathbb{R}^2 this time, that

$$|\nabla\zeta(x)| \leq \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma'}}{(1+\tilde{r})^{1+\sigma}}. \quad (4.17)$$

Here, we cannot integrate from infinity (since the estimate is only on a ball) to get an estimation on ζ , but this will be dealt with later on.

Now, we define $\mathcal{Y}'_2 := \mathcal{Y}_2 - \zeta$, and we have, for $j \in \{1, 2\}$,

$$\partial_{x_j} \mathcal{Y}'_2 = K_j * f',$$

where

$$f' := -\mathcal{H}_1 - 2\mathfrak{Re} \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) - (1 - |V|^2) \mathcal{Y}_1 - c \partial_{x_2} \zeta.$$

We first estimate the convolution in $B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon)$. With $\|\mathcal{Y}_n\|_{*, B'_{d_n}} = 1$, we check that, with some margin in $B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon)$,

$$\left| 2\mathfrak{Re} \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) - (1 - |V|^2) \mathcal{Y}_1 \right| \leq \frac{o_{R \rightarrow \infty}(1)}{(1 + r_1)^{3/2 - \alpha}}.$$

Now, we have shown in step 6 that

$$\left| \mathfrak{Re} \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) \right| \leq \frac{o_{n \rightarrow \infty}(1)}{r_1^{1 - \alpha + \sigma''}}$$

for some $\sigma'' > 0$. In $B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon) \setminus B(d_n \vec{e}_1^\rightarrow, d_n^{\varepsilon'})$, we have

$$|(1 - |V|^2) \mathcal{Y}_1| \leq \frac{d_n^{\lambda + \sigma'}}{r_1^{3 + \sigma}} \leq \frac{d_n^{\lambda + \sigma' - (2 + \alpha - \sigma'')\varepsilon'}}{r_1^{1 - \alpha + \sigma''}} = \frac{o_{n \rightarrow \infty}(1)}{r_1^{1 - \alpha + \sigma''}}$$

given that σ' and σ'' are small enough since $\lambda - (2 + \alpha)\varepsilon' < 0$. Therefore, we check with the same computations (since we have some margin $\sigma'' > 0$ on the decay) that

$$\left| \int_{B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon)} K_j(x - Y) \left(2\mathfrak{Re} \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) - (1 - |V|^2) \mathcal{Y}_1 \right) (Y) dY \right| \leq \frac{o_{R \rightarrow \infty}(1)}{(1 + r_1)^{1 - \alpha}}.$$

Now, using (4.16), we check that

$$\left| \int_{B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon)} K_j(x - Y) (c \partial_{x_2} \zeta)(Y) dY \right| \leq \frac{c(o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1))}{(1 + r_1)^{1 - \alpha - 1/10}}.$$

And, since $x \in B(d_n \vec{e}_1^\rightarrow, d_n^{\varepsilon'})$, $c(1 + r_1)^{1/10} \leq K$, therefore

$$\left| \int_{B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon)} K_j(x - Y) (c \partial_{x_2} \zeta)(Y) dY \right| \leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1 + r_1)^{1 - \alpha}}.$$

For the last remaining term, we use (4.7) with $\sigma = \frac{\lambda + 1}{2}$ to estimate

$$|\mathcal{H}_1| \leq \frac{o_{n \rightarrow 0}^R(1)}{(1 + r_1)},$$

and then, from Lemma 2.13 (only changing the integral from \mathbb{R}^2 to $B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon)$ in the proof), we infer

$$\left| \int_{B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon)} K_j(x - Y) \mathcal{H}_1(Y) dY \right| \leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1 + r_1)^{1 - \alpha}}.$$

Combining these estimates, we have shown that

$$\left| \int_{B(d_n \vec{e}_1^\rightarrow, d_n^\varepsilon)} K_j(x - Y) f'(Y) dY \right| \leq \frac{o_{R \rightarrow \infty}(1) + o_{n \rightarrow \infty}^R(1)}{(1 + r_1)^{1 - \alpha}}.$$

Now, we focus on the left half-plane. From (4.8), we have

$$|\mathcal{H}_1| \leq \frac{K(\sigma)c_n^{1-\sigma}d_n^{1+\lambda}}{(1+r_{-1})^2}.$$

Furthermore, we check, using (4.11) and (4.17) that, in the left half-plane,

$$\left| -2\Re \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) - (1 - |V|^2)\mathcal{Y}_1 \right| \leq \frac{K(\sigma, \sigma')d_n^{1+\lambda}c_n^{1-\sigma'}}{(1+r_{-1})^{2+\sigma}}$$

and

$$|c_n \partial_{x_2} \zeta| \leq \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma'}c_n}{(1+r_{-1})^{1+\sigma}}.$$

We have by Theorem 2.12 (since $x \in B(d_n \vec{e}_1, d_n^{\varepsilon'})$) that $|K_j(x - Y)| \leq \frac{K}{d_n^\beta (1+\tilde{r}(Y))^{2-\beta}}$ for Y in the left half-plane, for any $0 \leq \beta \leq 2$. Therefore, taking $\beta = 2 - \sigma$, we have

$$\left| \int_{\{y_1 \leq 0\}} K_j(x - Y) \mathcal{H}_1(Y) dY \right| \leq \int_{\mathbb{R}^2} \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma+\sigma'-2}}{(1+\tilde{r})^{2+\sigma}} \leq \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma+\sigma'-2+(1-\alpha)\varepsilon'}}{(1+|x - d_n \vec{e}_1|)^{1-\alpha}}.$$

Taking $\beta = 2$, we have

$$\begin{aligned} & \left| \int_{\{Y_1 \leq 0\}} K_j(x - Y) \left(-2\Re \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) - (1 - |V|^2)\mathcal{Y}_1 \right) (Y) dY \right| \\ & \leq \int_{\mathbb{R}^2} \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma'-2}}{(1+\tilde{r})^{2+\sigma}} \\ & \leq \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma'-2+(1-\alpha)\varepsilon'}}{(1+|x - d_n \vec{e}_1|)^{1-\alpha}}, \end{aligned}$$

and finally, taking $\beta = 1$, we estimate

$$\left| \int_{\{Y_1 \leq 0\}} K_j(x - Y) c_n \partial_{x_2} \zeta(Y) dY \right| \leq \int_{\mathbb{R}^2} \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma'-2}}{(1+\tilde{r})^{2+\sigma}} \leq \frac{K(\sigma, \sigma')d_n^{\lambda+\sigma'-2+(1-\alpha)\varepsilon'}}{(1+|x - d_n \vec{e}_1|)^{1-\alpha}}.$$

Thus, taking $\sigma' > \sigma > 0$ small enough, since $\lambda - 2 + (1 - \alpha)\varepsilon' < 0$, we have

$$\left| \int_{\{Y_1 \leq 0\}} K_j(x - Y) f'(Y) dY \right| \leq \frac{o_{n \rightarrow 0}(1)}{(1+|x - d_n \vec{e}_1|)^{1-\alpha}}.$$

We are left with the estimation in $\Omega := \{Y_1 \geq 0\} \setminus B(d_n \vec{e}_1, d_n^\varepsilon)$. We infer that, in Ω , we have, for $0 < \sigma < \sigma' < 1$

$$|f'| \leq \frac{K(\sigma')d_n^{\lambda-\sigma'}}{(1+r_1)} + \frac{K(\sigma)d_n^{\lambda+\sigma}}{(1+r_1)^{2+\sigma}}.$$

Indeed, from equation (4.13) and (4.17), we have $|\mathcal{H}_1 - c \partial_{x_2} \zeta| \leq \frac{K(\sigma)d_n^{\lambda+\sigma}}{(1+r_1)}$, and using (4.11), we check that

$$\left| 2\Re \left(\frac{\nabla V}{V} \cdot \nabla \mathcal{Y}_n \right) - (1 - |V|^2)\mathcal{Y}_1 \right| \leq \frac{K(\sigma)d_n^{\lambda+\sigma}}{(1+\tilde{r})^{2+\sigma}}.$$

Now, for $y \in \Omega$, $x \in B(d_n \vec{e}_1, d_n^{\varepsilon'})$, we have from Theorem 2.12 that

$$|K_j(x - Y)| \leq \frac{K}{d_n^{2\varepsilon}}$$

and

$$|K_j(x - Y)| \leq \frac{K}{(1 + \tilde{r}(Y))^{3/2} d_n^{\varepsilon/2}}.$$

We deduce that, for $x \in B(d_n \vec{e}_1, d_n^{\varepsilon'})$,

$$\begin{aligned} \int_{\Omega} |K_j(x - Y)| \frac{K(\sigma') d_n^{\lambda - \sigma'}}{(1 + r_1(Y))} dY &\leq K(\sigma') d_n^{\lambda - \sigma' - \varepsilon/2} \int_{\mathbb{R}^2} \frac{K}{(1 + \tilde{r}(Y))^{5/2}} dY \\ &\leq \frac{K(\sigma') d_n^{\lambda - \sigma' + (1 - \alpha)\varepsilon' - \varepsilon/2}}{(1 + |x - d_n \vec{e}_1|)^{1 - \alpha}} = \frac{o_{n \rightarrow 0}(1)}{(1 + |x - d_n \vec{e}_1|)^{1 - \alpha}} \end{aligned}$$

taking $\sigma' < 1$ large enough (since $\lambda + (1 - \alpha)\varepsilon' - 1 - \varepsilon/2 < 0$), and

$$\begin{aligned} \int_{\Omega} |K_j(x - Y)| \frac{K(\sigma) d_n^{\lambda + \sigma}}{(1 + \tilde{r}(Y))^{2 + \sigma}} dY &\leq K(\sigma) d_n^{\lambda + \sigma - 2\varepsilon} \int_{\mathbb{R}^2} \frac{1}{(1 + \tilde{r}(Y))^{2 + \sigma}} dY \\ &\leq \frac{K(\sigma) d_n^{\lambda + \sigma + (1 - \alpha)\varepsilon' - 2\varepsilon}}{(1 + |x - d_n \vec{e}_1|)^{1 - \alpha}} = \frac{o_{n \rightarrow 0}(1)}{(1 + |x - d_n \vec{e}_1|)^{1 - \alpha}} \end{aligned}$$

taking $\sigma > 0$ small enough (since $\lambda + (1 - \alpha)\varepsilon' - 2\varepsilon < 0$). We deduce that, for $x \in B(d_n \vec{e}_1, d_n^{\varepsilon'})$,

$$|\partial_{x_j} \mathcal{Y}'_2| = |K_j * f'| \leq \frac{o_{n \rightarrow 0}(1) + o_{R \rightarrow \infty}(1)}{(1 + |x - d_n \vec{e}_1|)^{1 - \alpha}}.$$

With (4.16), we have shown that

$$|\partial_{x_j} \mathcal{Y}_2| \leq \frac{o_{n \rightarrow 0}(1) + o_{R \rightarrow \infty}(1)}{(1 + |x - d_n \vec{e}_1|)^{1 - \alpha}}.$$

Now, since $|\mathcal{Y}_2| + |\nabla \mathcal{Y}_2| = o_{R \rightarrow \infty}(1)$ in $B(d_n \vec{e}_1, 10)$, by integration from $d_n \vec{e}_1$, we check that, since $\alpha > 0$,

$$|\mathcal{Y}_2| \leq \frac{o_{n \rightarrow 0}(1) + o_{R \rightarrow \infty}(1)}{(1 + |x - d_n \vec{e}_1|)^{-\alpha}}.$$

Step 7.2. Elliptic estimate for \mathcal{Y}_1 .

For \mathcal{Y}_1 we also use the function K_0 and we have

$$\mathcal{Y}_1 = \frac{1}{2\pi} K_0 \left(\sqrt{2} |\cdot| \right) * (-\Delta \mathcal{Y}_1 + 2\mathcal{Y}_1),$$

therefore

$$\begin{aligned} |\mathcal{Y}_1|(x) &\leq \int_{\bar{B}_{d_n}(x)} \frac{1}{2\pi} K_0 \left(\sqrt{2} |x - Y| \right) |(-\Delta \mathcal{Y}_1 + 2\mathcal{Y}_1)(Y)| dY \\ &+ \int_{\mathbb{R} \setminus \bar{B}_{d_n}(x)} \frac{1}{2\pi} K_0 \left(\sqrt{2} |x - Y| \right) |(-\Delta \mathcal{Y}_1 + 2\mathcal{Y}_1)(Y)| dY, \end{aligned}$$

where $B_{d_n}(x) = B(x - d_n \vec{e}_1, d_n^{\varepsilon})$. The first term can be computed as for Lemma 2.10, and for the second term, in $\mathbb{R} \setminus B_{d_n}$, we have

$$K_0 \left(\sqrt{2} |x| \right) \leq K e^{-d_n^{\varepsilon/2}} e^{-|x|^{1/2}}$$

from Lemma 2.9, which, with (4.11) and (4.12), make the term integrable and a $o_{d_n \rightarrow \infty}(e^{-d_n^{\varepsilon/4}})$, which is enough to show that

$$|\nabla \mathcal{Y}_1| + |\mathcal{Y}_1| \leq \frac{o_{n \rightarrow \infty}(1) + o_{R \rightarrow \infty}(1)}{(1 + r_1)^{1 - \alpha}}.$$

Step 8. Conclusion.

We conclude that there is a contradiction, as in the end of the proof of Proposition 2.17. This ends the proof of Proposition 4.5. \square

In the rest of this chapter, we take $\alpha, \varepsilon, \varepsilon', \lambda$ such that they satisfy the conditions of Proposition 4.5, and

$$\lambda + (1 - \alpha)\varepsilon' > 1.$$

4.4 Proof of $\int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \partial_{z_1}(\text{TW}_c(V)))|_{d=d_c} = \frac{-\pi}{d_c^2} + o_{d_c \rightarrow \infty} \left(\frac{1}{d_c^2} \right)$

From (2.2), the equation on V is

$$\text{TW}_c(V) = E - ic\partial_{x_2} V = -2\nabla V_1 \cdot \nabla V_{-1} + (1 - |V_1|^2)(1 - |V_{-1}|^2)V_1 V_{-1} - ic\partial_{x_2}(V_1 V_{-1}).$$

We use Lemma 4.2 to compute

$$\partial_{z_1} V = \partial_{x_1} V_1 V_{-1} + \partial_{x_1} V_{-1} V_1 - (-\partial_{x_1} V_1 V_{-1} + \partial_{x_1} V_{-1} V_1) = 2\partial_{x_1} V_1 V_{-1}.$$

Therefore

$$\partial_{z_1} \text{TW}_c(V) = -4\nabla V_1 \cdot \nabla \partial_{x_1} V_{-1} + 2(1 - |V_1|^2)V_1 \partial_{x_1}((1 - |V_{-1}|^2)V_{-1}) - 2ic\partial_{x_2}(V_1 \partial_{x_1} V_{-1}),$$

and then

$$\begin{aligned} \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \partial_{z_1}(\text{TW}_c(V))) &= -4 \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \nabla V_1 \cdot \nabla \partial_{x_1} V_{-1}) \\ &+ 2 \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} (1 - |V_1|^2) V_1 \partial_{x_1}((1 - |V_{-1}|^2) V_{-1})) \\ &- 2 \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} ic\partial_{x_2}(V_1 \partial_{x_1} V_{-1})). \end{aligned}$$

We want to compute this quantity at $d = d_c$. We omit the subscript and use only d in this proof. In fact, it works for any d such that $\frac{1}{2d} \leq c \leq \frac{2}{d}$. We infer, by Lemma 4.4, that

$$\int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} (1 - |V_1|^2) V_1 \partial_{x_1}((1 - |V_{-1}|^2) V_{-1})) = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right)$$

and

$$\int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} ic\partial_{x_2}(V_1 \partial_{x_1} V_{-1})) = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

Let us show that $\int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} (-4\nabla V_1 \cdot \nabla \partial_{x_1} V_{-1})) = -\frac{\pi}{d^2} + o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right)$. We have

$$-4\nabla V_1 \cdot \nabla \partial_{x_1} V_{-1} = -4\partial_{x_1} V_1 \partial_{x_1 x_1} V_{-1} - 4\partial_{x_2} V_1 \partial_{x_1 x_2} V_{-1}.$$

Remark that using $|\partial_d V| \leq \frac{K}{(1+r_1)}$ and Lemma 4.4 once again,

$$\left| \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \partial_{x_1} V_1 \partial_{x_1 x_1} V_{-1}) \right| \leq \int_{B(d\vec{e}_1, d^{\varepsilon'})} \frac{K}{(1+r_1^2)d^{3-\varepsilon'}} = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

Moreover,

$$\begin{aligned} &-4 \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \partial_{x_2} V_1 \partial_{x_1 x_2} V_{-1}) = \\ &4 \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_{x_1} V_1 V_{-1}} \partial_{x_2} V_1 \partial_{x_1 x_2} V_{-1}) - 4 \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_{x_1} V_{-1} V_1} \partial_{x_2} V_1 \partial_{x_1 x_2} V_{-1}). \end{aligned} \quad (4.18)$$

For the first integral in (4.18), we write

$$\begin{aligned} &4 \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_{x_1} V_1 V_{-1}} \partial_{x_2} V_1 \partial_{x_1 x_2} V_{-1}) = \\ &4 \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_{x_1} V_1} \partial_{x_2} V_1) \Re(\overline{V_{-1}} \partial_{x_1 x_2} V_{-1}) - \Im(\overline{\partial_{x_1} V_1} \partial_{x_2} V_1) \Im(\overline{V_{-1}} \partial_{x_1 x_2} V_{-1}). \end{aligned}$$

For the first contribution, we have

$$\left| \int_{B(d\bar{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_{x_1} V_1} \partial_{x_2} V_1) \Re(\overline{V_{-1}} \partial_{x_1 x_2} V_{-1}) \right| \leq \int_{B(d\bar{e}_1, d^{\varepsilon'})} \frac{K}{(1+r_1^2)d^{3-\varepsilon'}} = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

For the second contribution, recall from Lemma 2.2 that

$$\partial_{x_1} V_1 = \left(\cos(\theta_1)u - \frac{i}{r_1} \sin(\theta_1) \right) V_1 \quad \text{and} \quad \partial_{x_2} V_1 = \left(\sin(\theta_1)u + \frac{i}{r_1} \cos(\theta_1) \right) V_1,$$

therefore

$$\Im(\overline{\partial_{x_1} V_1} \partial_{x_2} V_1) = \frac{u}{r_1} |V_1|^2,$$

and then, by Lemma 4.4,

$$-4 \int_{B(d\bar{e}_1, d^{\varepsilon'})} \Im(\overline{\partial_{x_1} V_1} \partial_{x_2} V_1) \Im(\overline{V_{-1}} \partial_{x_1 x_2} V_{-1}) = -4 \int_{B(d\bar{e}_1, d^{\varepsilon'})} \frac{u}{r_1} \frac{1}{4d^2} |V_1|^2 dr_1 + o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right)$$

since

$$\int_{B(d\bar{e}_1, d^{\varepsilon'})} \frac{u}{r_1} \frac{1}{4d^{2+1/4}} |V_1|^2 dr_1 = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

We compute, using $|V_1|^2 = \rho_1^2$, $u = \frac{\rho_1'}{\rho_1}$ and Lemma 2.1,

$$-4 \int_{B(d\bar{e}_1, d^{\varepsilon'})} \frac{u}{r_1} \frac{|V_1|^2}{4d^2} dr_1 = \frac{-2\pi}{d^2} \int_0^{d^{\varepsilon'}} \rho_1'(r_1) \rho_1(r_1) dr_1 = \frac{-\pi}{d^2} [\rho_1^2]_0^{d^{\varepsilon'}} = \frac{-\pi}{d^2} + o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

We obtain the estimate for the first integral in (4.18):

$$4 \int_{B(d\bar{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_{x_1} V_1} \overline{V_{-1}} \partial_{x_2} V_1 \partial_{x_1 x_2} V_{-1}) = \frac{-\pi}{d^2} + o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

For the second integral in (4.18), we estimate

$$\left| \int_{B(d\bar{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_{x_1} V_{-1}} \overline{V_1} \partial_{x_2} V_1 \partial_{x_1 x_2} V_{-1}) \right| \leq \int_{B(d\bar{e}_1, d^{\varepsilon'})} \frac{K}{(1+r_1)d^{4-\varepsilon'}} = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

This ends the proof of this subsection.

4.5 Proof of $\partial_d \int_{B(d\bar{e}_1, d^{\varepsilon'}) \cup B(-d\bar{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \text{TW}_c(Q_{c,d}))|_{d=d_c} = \frac{-2\pi}{d_c^2} + o_{d_c \rightarrow \infty} \left(\frac{1}{d_c^2} \right)$

In order to prove the result of this subsection, by using (4.1) and the result of subsection 4.4 we just have to show that at $d = d_c$,

$$\begin{aligned} & \int_{B(d\bar{e}_1, \varepsilon')} \Re(\overline{\partial_d V} L(\partial_{z_1} \Gamma_{c,d_c})) + \int_{B(d\bar{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V}(\partial_{z_1} L)(\Gamma_{c,d_c})) \\ & + \int_{B(d\bar{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \partial_{z_1}(\text{NL}_V(\Gamma_{c,d_c}))) = o_{d_c \rightarrow \infty} \left(\frac{1}{d_c^2} \right). \end{aligned}$$

Similarly to subsection 4.4, we omit the subscript on d_c in the proof.

Step 1. Proof of $\int_{B(d\bar{e}_1, d^{3/4})} \Re(\overline{\partial_d V} L(\partial_{z_1} \Gamma_{c,d})) = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right)$.

For this term, we want to do integration by parts and use that $L(\partial_d V)$ is very small, but since the integral is not on the whole space, there are the two boundary terms:

$$\begin{aligned} & \left| \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} L(\partial_{z_1} \Gamma_{c,d})) \right| \leq \left| \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(L(\partial_d V) \overline{\partial_{z_1} \Gamma_{c,d}}) \right| \\ & + \left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \nabla \partial_{z_1} \Gamma_{c,d}) \right| + \left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\nabla \overline{\partial_d V} \partial_{z_1} \Gamma_{c,d}) \right|, \end{aligned}$$

where $\partial B(d\vec{e}_1, d^{\varepsilon'})$ is the boundary of $B(d\vec{e}_1, d^{\varepsilon'})$. On $\partial B(d\vec{e}_1, d^{\varepsilon'})$, we have

$$\Gamma_{c,d} = V(e^{\Psi_{c,d}} - 1),$$

hence

$$\partial_{z_1} \Gamma_{c,d} = 2V_1 \partial_{x_1} V_{-1}(e^{\Psi_{c,d}} - 1) + V \partial_{z_1} \Psi_{c,d} e^{\Psi_{c,d}} \quad (4.19)$$

and

$$\begin{aligned} \nabla \partial_{z_1} \Gamma_{c,d} &= 2\nabla V_1 \partial_{x_1} V_{-1}(e^{\Psi_{c,d}} - 1) + 2V_1 \nabla \partial_{x_1} V_{-1}(e^{\Psi_{c,d}} - 1) + 2V_1 \partial_{x_1} V_{-1} \nabla \Psi_{c,d} e^{\Psi_{c,d}} \\ &+ \nabla V \partial_{z_1} \Psi_{c,d} e^{\Psi_{c,d}} + V \nabla \partial_{z_1} \Psi_{c,d} e^{\Psi_{c,d}} + V \partial_{z_1} \Psi_{c,d} \nabla \Psi_{c,d} e^{\Psi_{c,d}}. \end{aligned} \quad (4.20)$$

By Lemmas 2.2 and 4.4, Proposition 4.5 and (3.10), we infer on $\partial B(d\vec{e}_1, d^{\varepsilon'})$ that, for any $1 > \sigma > 0$,

$$|\partial_{z_1} \Gamma_{c,d}| \leq \frac{K(\sigma)}{d^{2-\varepsilon'} d^{1-\sigma}} + \frac{K}{d^{1+\lambda-\alpha\varepsilon'}}. \quad (4.21)$$

Thus, still on $\partial B(d\vec{e}_1, d^{\varepsilon'})$, from Lemma 2.6 we compute

$$\left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\nabla \overline{\partial_d V} \partial_{z_1} \Gamma_{c,d}) \right| \leq \frac{K}{d^{\varepsilon'}} \left(\frac{K(\sigma)}{d^{2-\varepsilon'} d^{1-\sigma}} + \frac{K}{d^{1+\lambda-\alpha\varepsilon'}} \right) \leq \frac{K(\sigma)}{d^{3-\sigma}} + \frac{K}{d^{1+\lambda+(1-\alpha)\varepsilon'}}.$$

Since $3 - \sigma > 2$ and $\lambda + (1 - \alpha)\varepsilon' > 1$ by (4.2), we have

$$\left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\nabla \overline{\partial_d V} \partial_{z_1} \Gamma_{c,d}) \right| = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

For (4.20), we estimate on $\partial B(d\vec{e}_1, d^{\varepsilon'})$, still using Lemmas 2.2 and 4.4, Proposition 4.5 and (3.10), for any $1 > \sigma > 0$,

$$|2\nabla V_1 \partial_{x_1} V_{-1}(e^{\Psi_{c,d}} - 1) + 2V_1 \nabla \partial_{x_1} V_{-1}(e^{\Psi_{c,d}} - 1) + 2V_1 \partial_{x_1} V_{-1} \nabla \Psi_{c,d} e^{\Psi_{c,d}}| \leq \frac{K(\sigma)}{d^{3-\sigma}},$$

and

$$|\nabla V \partial_{z_1} \Psi_{c,d} e^{\Psi_{c,d}} + V \nabla \partial_{z_1} \Psi_{c,d} e^{\Psi_{c,d}} + V \partial_{z_1} \Psi_{c,d} \nabla \Psi_{c,d} e^{\Psi_{c,d}}| \leq \frac{K}{d^{1+\lambda+(1-\alpha)\varepsilon'}} + \frac{K(\sigma)}{e^{2+\lambda+(1-\alpha)\varepsilon'-\sigma}}.$$

In particular, from (4.20), we can find $1 > \sigma > 0$ such that, on $\partial B(d\vec{e}_1, d^{\varepsilon'})$,

$$|\nabla \partial_{z_1} \Gamma_{c,d}| = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right),$$

thus

$$\left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \nabla \partial_{z_1} \Gamma_{c,d}) \right| = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

From (2.27), we know that

$$|L(\partial_d V)| \leq \frac{K}{(1 + \tilde{r}^2)d}.$$

Moreover, by Proposition 4.5, we have $|\partial_{z_1}\Gamma_{c,d}| \leq \frac{K}{d^{1+\lambda-\alpha\varepsilon'}}$ in $B(d\vec{e}_1^\lambda, d^{\varepsilon'})$, which is enough to show that

$$\int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \Re(L(\overline{\partial_d V})\partial_{z_1}\Gamma_{c,d}) = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

Step 2. Proof of $\int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \Re(\overline{\partial_d V}(\partial_{z_1}L)(\Gamma_{c,d})) = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right)$.

We have

$$(\partial_{z_1}L)(\Gamma_{c,d}) = 4\Re(\overline{V_{-1}}\partial_{x_1}V_{-1})\Gamma_{c,d} + 4\Re(\overline{\partial_{x_1}V_{-1}V_1}\Gamma_{c,d})V + 4\Re(\overline{V}\Gamma_{c,d})V_1\partial_{x_1}V_{-1},$$

thus

$$\begin{aligned} \int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \Re(\overline{\partial_d V}(\partial_{z_1}L)(\Gamma_{c,d})) &= 4 \int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \Re(\overline{\partial_d V}\Gamma_{c,d})\Re(\overline{V_{-1}}\partial_{x_1}V_{-1}) \\ &+ 4 \int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \Re(\overline{\partial_{x_1}V_{-1}V_1}\Gamma_{c,d})\Re(\overline{\partial_d V}V) \\ &+ 4 \int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \Re(\overline{\partial_d V}V_1\partial_{x_1}V_{-1})\Re(\overline{V}\Gamma_{c,d}). \end{aligned} \quad (4.22)$$

Using $|\partial_d V| \leq \frac{K}{1+r_1}$,

$$\Re(\overline{V_{-1}}\partial_{x_1}V_{-1}) = O_{d \rightarrow \infty} \left(\frac{1}{d^3} \right) \quad \text{and} \quad |\Gamma_{c,d}| \leq \frac{K}{(1+r_1)^{1/2}d^{1/2}}$$

from Lemma 2.6, Lemma 2.2 and (3.5) respectively, we may bound

$$\left| \int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \Re(\overline{\partial_d V}\Gamma_{c,d})\Re(\overline{V_{-1}}\partial_{x_1}V_{-1}) \right| \leq \int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \frac{K}{(1+r_1)^{1+1/2}d^{3+1/2}} = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

The second term of (4.22) is

$$4 \int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} \Re(\overline{\partial_{x_1}V_{-1}V_1}\Gamma_{c,d})\Re(\overline{\partial_d V}V).$$

We compute that

$$|\Re(\overline{\partial_{x_1}V_{-1}V_1}\Gamma_{c,d})| \leq \frac{K}{(1+r_1)^{1/8}d^{17/8}} \quad \text{and} \quad |\Re(\overline{\partial_d V}V)| \leq \frac{K}{(1+r_1)^3}$$

in $B(d\vec{e}_1^\lambda, d^{\varepsilon'})$ using

$$|\Gamma_{c,d}| \leq \frac{K}{(1+r_1)^{1/8}d^{7/8}}$$

by (3.10) and the definition of $\Gamma_{c,d}$. Therefore, since $17/8 > 2$,

$$\int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} 4\Re(\overline{\partial_{x_1}V_{-1}V_1}\Gamma_{c,d})\Re(\overline{\partial_d V}V) = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

The last term of (4.22) is

$$\int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} 4\Re(\overline{V}\Gamma_{c,d})\Re(V_1\partial_{x_1}V_{-1}\overline{\partial_d V}).$$

Recalling that

$$|\Re(\overline{V}\Gamma_{c,d})| \leq K|\Re(\Psi)| \leq \frac{K}{(1+r_1)^{1+1/8}d^{7/8}}$$

and

$$|\Re(V_1\partial_{x_1}V_{-1}\overline{\partial_d V})| \leq \frac{K}{d^{5/4}(1+r_1)},$$

we deduce

$$\int_{B(d\vec{e}_1^\lambda, d^{\varepsilon'})} 4\Re(\overline{V}\Gamma_{c,d})\Re(V_1\partial_{x_1}V_{-1}\overline{\partial_d V}) = o_{d \rightarrow \infty} \left(\frac{1}{d^2} \right).$$

Step 3. Proof of $\int_{B(d\vec{e}_1, d\varepsilon')} \Re(\overline{\partial_d V} \partial_{z_1}(\text{NL}_V(\Gamma_{c,d}))) = o_{d \rightarrow \infty} \left(\frac{1}{d^2}\right)$.

Recall that

$$\begin{aligned} \partial_{z_1} \text{NL}_V(\Gamma_{c,d}) &= 4\Re(\overline{\partial_{x_1} V_{-1} V_1} \Gamma_{c,d}) \Gamma_{c,d} + 2\Re(\overline{V} \partial_{z_1} \Gamma_{c,d}) \Gamma_{c,d} + 2\Re(\overline{V} \Gamma_{c,d}) \partial_{z_1} \Gamma_{c,d} \\ &\quad + 2\Re(\overline{\Gamma_{c,d}} \partial_{z_1} \Gamma_{c,d})(V + \Gamma_{c,d}) + |\Gamma_{c,d}|^2 (2\partial_{x_1} V_{-1} V_1 + \partial_{z_1} \Gamma_{c,d}). \end{aligned}$$

We write

$$\int_{B(d\vec{e}_1, d\varepsilon')} \Re(\overline{\partial_d V} \partial_{z_1}(\text{NL}_V(\Gamma_{c,d}))) = I_1 + I_2 + I_3 + I_4 + I_5,$$

with

$$\begin{aligned} I_1 &= \int_{B(d\vec{e}_1, d\varepsilon')} 4\Re(\overline{\partial_d V} \Gamma_{c,d}) \Re(\overline{\partial_{x_1} V_{-1} V_1} \Gamma_{c,d}), \\ I_2 &= \int_{B(d\vec{e}_1, d\varepsilon')} 2\Re(\overline{\partial_d V} \Gamma_{c,d}) \Re(\overline{V} \partial_{z_1} \Gamma_{c,d}), \\ I_3 &= \int_{B(d\vec{e}_1, d\varepsilon')} 2\Re(\overline{V} \Gamma_{c,d}) \Re(\overline{\partial_d V} \partial_{z_1} \Gamma_{c,d}), \\ I_4 &= \int_{B(d\vec{e}_1, d\varepsilon')} 2\Re(\overline{\Gamma_{c,d}} \partial_{z_1} \Gamma_{c,d}) \Re(\overline{\partial_d V} V) + 2\Re(\overline{\Gamma_{c,d}} \partial_{z_1} \Gamma_{c,d}) \Re(\overline{\partial_d V} \Gamma_{c,d}), \\ I_5 &= \int_{B(d\vec{e}_1, d\varepsilon')} |\Gamma_{c,d}|^2 \Re(\overline{\partial_d V} (2\partial_{x_1} V_{-1} V_1 + \partial_{z_1} \Gamma_{c,d})). \end{aligned}$$

We infer that all these terms are $o_{d \rightarrow \infty} \left(\frac{1}{d^2}\right)$.

4.6 Proof of $\partial_c d_c = -\frac{1}{c^2} + o_{c \rightarrow 0} \left(\frac{1}{c^2}\right)$

Recall that d_c is defined by the implicit equation

$$\int_{B(d\vec{e}_1, d\varepsilon') \cup B(-d\vec{e}_1, d\varepsilon')} \Re(\overline{\partial_d V} \text{TW}_c(Q_{c,d})) = 0.$$

We showed in subsection 4.5 that

$$\partial_d \int_{B(d\vec{e}_1, d\varepsilon') \cup B(-d\vec{e}_1, d\varepsilon')} \Re(\overline{\partial_d V} \text{TW}_c(Q_{c,d}))|_{d=d_c} = \frac{-2\pi}{d_c^2} + o_{d_c \rightarrow \infty} \left(\frac{1}{d_c^2}\right).$$

Therefore, by the implicit function theorem,

$$\partial_c d_c = \frac{\partial_c \int_{B(d\vec{e}_1, d\varepsilon') \cup B(-d\vec{e}_1, d\varepsilon')} \Re(\overline{\partial_d V} \text{TW}_c(Q_{c,d}))|_{d=d_c}}{\frac{-2\pi}{d_c^2} + o_{d_c \rightarrow \infty} \left(\frac{1}{d_c^2}\right)}.$$

We compute for

$$\text{TW}_c(Q_{c,d}) = -ic\partial_{x_2} Q_{c,d} - \Delta Q_{c,d} - (1 - |Q_{c,d}|^2)Q_{c,d}$$

that, with $\partial_c Q_{c,d} = \partial_c(V + \Gamma_{c,d}) = \partial_c \Gamma_{c,d}$ at fixed d , we have (still at fixed d)

$$\partial_c(\text{TW}_c(Q_{c,d})) = -i\partial_{x_2} Q_{c,d} - L_{Q_{c,d}}(\partial_c \Gamma_{c,d}),$$

where

$$L_{Q_{c,d}}(h) := -\Delta h - ic\partial_{x_2} h - (1 - |Q_{c,d}|^2)h + 2\Re(\overline{Q_{c,d}} h)Q_{c,d}.$$

We are left with the computation of

$$\partial_c \int_{B(d\vec{e}_1, d\varepsilon') \cup B(-d\vec{e}_1, d\varepsilon')} \Re(\overline{\partial_d V} \text{TW}_c(Q_{c,d}))|_{d=d_c} =$$

$$\begin{aligned}
& - \int_{B(d\bar{e}_1^\dagger, d^{e'}) \cup B(-d\bar{e}_1^\dagger, d^{e'})} \Re(\bar{\partial}_d \bar{V}(i\partial_{x_2} Q_{c,d}))|_{d=d_c} \\
& - \int_{B(d\bar{e}_1^\dagger, d^{e'}) \cup B(-d\bar{e}_1^\dagger, d^{e'})} \Re(\bar{\partial}_d \bar{V} L_{Q_c}(\partial_c \Gamma_{c,d}))|_{d=d_c}.
\end{aligned}$$

As above, we omit the subscript in d_c for the computations.

Step 1. Proof of $\int_{B(d\bar{e}_1^\dagger, d^{e'}) \cup B(-d\bar{e}_1^\dagger, d^{e'})} \Re(\bar{\partial}_d \bar{V}(-i\partial_{x_2} Q_c))|_{d=d_c} = 2\pi + o_{c \rightarrow 0}(1)$.

We have $\partial_{x_2} Q_c = \partial_{x_2} V + \partial_{x_2} \Gamma_{c,d}$, hence

$$\begin{aligned}
& - \int_{B(d\bar{e}_1^\dagger, d^{e'}) \cup B(-d\bar{e}_1^\dagger, d^{e'})} \Re(\bar{\partial}_d \bar{V}(i\partial_{x_2} Q_c)) = \\
& - \int_{B(d\bar{e}_1^\dagger, d^{e'}) \cup B(-d\bar{e}_1^\dagger, d^{e'})} \Re(i\bar{\partial}_d \bar{V} \partial_{x_2} V) - \int_{B(d\bar{e}_1^\dagger, d^{e'}) \cup B(-d\bar{e}_1^\dagger, d^{e'})} \Re(i\bar{\partial}_d \bar{V} \partial_{x_2} \Gamma_{c,d}).
\end{aligned}$$

Since

$$|\partial_d V| \leq \frac{K}{(1+r_1)}$$

and

$$|\partial_{x_2} \Gamma_{c,d}| \leq \frac{K}{(1+r_1)^{1+1/2} d^{1/2}},$$

we have

$$\int_{B(d\bar{e}_1^\dagger, d^{e'})} \Re(i\bar{\partial}_d \bar{V} \partial_{x_2} \Gamma_{c,d}) = o_{c \rightarrow 0}(1).$$

Furthermore,

$$- \int_{B(d\bar{e}_1^\dagger, d^{e'})} \Re(i\bar{\partial}_d \bar{V} \partial_{x_2} V) = \int_{B(d\bar{e}_1^\dagger, d^{e'})} \Re(i\bar{\partial}_{x_1} \bar{V}_1 \partial_{x_2} V_1) + o_{c \rightarrow 0}(1),$$

and we already computed in (2.25) that

$$\int_{\mathbb{R}^2} \Re(i\partial_{x_2} V_1 \bar{\partial}_{x_1} \bar{V}_1) = -\pi + o_{c \rightarrow 0}(c^{1/4})$$

hence

$$\int_{B(d\bar{e}_1^\dagger, d^{e'}) \cup B(-d\bar{e}_1^\dagger, d^{e'})} \Re(\bar{\partial}_d \bar{V}(-i\partial_{x_2} Q_c))|_{d=d_c} = 2\pi + o_{c \rightarrow 0}(1).$$

Step 2. Proof of $\int_{B(d\bar{e}_1^\dagger, d^{e'}) \cup B(-d\bar{e}_1^\dagger, d^{e'})} \Re(\bar{\partial}_d \bar{V} L_{Q_c}(\partial_c \Gamma_{c,d}))|_{d=d_c} = o_{c \rightarrow 0}(1)$.

From the definition of $\Gamma_{c,d}$, at fixed d , we have

$$\partial_c \Gamma_{c,d} = \eta V \partial_c \Psi_{c,d} + (1-\eta) V \partial_c \Psi_{c,d} e^{\Psi_{c,d}}. \quad (4.23)$$

We have, by definition,

$$L_{Q_c}(\partial_c \Gamma_{c,d}) = -ic \partial_{x_2} \partial_c \Gamma_{c,d} - \Delta \partial_c \Gamma_{c,d} - (1 - |Q_c|^2) \partial_c \Gamma_{c,d} + 2\Re(\bar{Q}_c \partial_c \Gamma_{c,d}) Q_c,$$

and using $|\partial_d V| \leq \frac{K}{(1+r_1)}$ with $|\partial_{x_2} \partial_c \Gamma_{c,d}| \leq \frac{K c^{-1/2}}{(1+r_1)^{1+1/2}}$ since $\left\| \frac{\partial_c \Gamma_{c,d}}{V} \right\|_{*,1/2,d} \leq K c^{-3/4}$ from Lemma 3.9 and (4.23),

we have

$$\left| \int_{B(d\bar{e}_1^\dagger, d^{e'})} \Re(\bar{\partial}_d \bar{V}(-ic \partial_{x_2} \partial_c \Gamma_{c,d})) \right| \leq K \int_{B(d\bar{e}_1^\dagger, d^{e'})} \frac{c^{1/4}}{(1+r_1)^{2+1/2}} = o_{c \rightarrow 0}(1).$$

The estimate on $B(-d\bar{e}_1^\dagger, d^{e'})$ is similar.

We define

$$\tilde{L}_{Q_c}(h) := -\Delta h - (1 - |Q_c|^2)h + 2\Re(\overline{Q_c}h)Q_c$$

and we are then left with the computation of

$$\int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \tilde{L}_{Q_c}(\partial_c \Gamma_{c,d})),$$

the part on $B(-d\vec{e}_1, d^{\varepsilon'})$ being symmetrical. We want to put the linear operator onto $\partial_d V$ since $\tilde{L}_{Q_c}(\partial_d V)$ is close to $L_V(\partial_d V)$ which is itself small. We then integrate by parts:

$$\begin{aligned} & \left| \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \tilde{L}_{Q_c}(\partial_c \Gamma_{c,d})) \right| \leq \left| \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\tilde{L}_{Q_c}(\partial_d V) \overline{\partial_c \Gamma_{c,d}}) \right| \\ & + \left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \nabla \partial_c \Gamma_{c,d}) \right| + \left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\nabla \overline{\partial_d V} \partial_c \Gamma_{c,d}) \right|. \end{aligned}$$

We have on $\partial B(d\vec{e}_1, d^{\varepsilon'})$, that $|\partial_d V| \leq \frac{K}{d^{3/4}}$, $|\nabla \partial_d V| \leq \frac{K}{d^{3/2}}$ from Lemma 2.6. Moreover, by $\left\| \frac{\partial_c \Gamma_{c,d}}{V} \right\|_{*,1/2,d} \leq K(\sigma)c^{-1/2-\sigma}$ from Lemma 3.9 and (4.23), we deduce $|\nabla \partial_c \Gamma_{c,d}| \leq \frac{K(\sigma)d^{1/2+\sigma}}{d^{(3/4)(3/2)}} \leq \frac{K(\sigma)}{d^{5/8-\sigma}}$ and $|\partial_c \Gamma_{c,d}| \leq \frac{K(\sigma)d^{1/2-\sigma}}{d^{(3/4)(1/2)}} \leq K(\sigma)d^{1/8-\sigma}$. We then obtain, for $\sigma > 0$ small enough,

$$\begin{aligned} & \left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \nabla \partial_c \Gamma_{c,d}) \right| \leq \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} |\partial_d V| |\nabla \partial_c \Gamma_{c,d}| \leq d^{3/4} \frac{K(\sigma)d^{2\sigma}}{d^{3/4}d^{5/8}} = o_{c \rightarrow 0}(1), \\ & \left| \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} \Re(\nabla \overline{\partial_d V} \partial_c \Gamma_{c,d}) \right| \leq \int_{\partial B(d\vec{e}_1, d^{\varepsilon'})} |\nabla \partial_d V| |\partial_c \Gamma_{c,d}| \leq d^{3/4} \frac{K(\sigma)d^{1/8+\sigma}}{d^{3/2}} = o_{c \rightarrow 0}(1). \end{aligned}$$

Therefore,

$$\int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \tilde{L}_{Q_c}(\partial_c \Gamma_{c,d})) = \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\tilde{L}_{Q_c}(\partial_d V) \overline{\partial_c \Gamma_{c,d}}) + o_{c \rightarrow 0}(1).$$

Now, from (2.27), we have that that

$$|L_V(\partial_d V)| \leq \frac{K}{(1+r_1)d}$$

and by Lemma 3.9 and (4.23), we have $|\partial_c \Gamma_{c,d}| \leq \frac{Kd^{1/4}}{(1+r_1)^{1/2}}$, hence

$$\left| \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(L_V(\partial_d V) \overline{\partial_c \Gamma_{c,d}}) \right| \leq K \int_{B(d\vec{e}_1, d^{\varepsilon'})} \frac{1}{(1+r_1)^{2+1/2}d^{1/4}} = o_{c \rightarrow 0}(1).$$

We deduce from this that

$$\int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re(\overline{\partial_d V} \tilde{L}_{Q_c}(\partial_c \Gamma_{c,d})) = \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re((\tilde{L}_{Q_c} - L_V)(\partial_d V) \overline{\partial_c \Gamma_{c,d}}) + o_{c \rightarrow 0}(1).$$

We have $\tilde{L}_{Q_c}(h) = -\Delta h - (1 - |Q_c|^2)h + 2\Re(\overline{Q_c}h)Q_c$ and $L_V(h) = -\Delta h - (1 - |V|^2)h + 2\Re(\overline{V}h)V$, therefore

$$(\tilde{L}_{Q_c} - L_V)(\partial_d V) = (|Q_c|^2 - |V|^2)\partial_d V + 2\Re(\overline{V}\partial_d V)(Q_c - V) + 2\Re(\overline{Q_c - V}\partial_d V)Q_c.$$

We have by (3.13) that $\left| |Q_c|^2 - |V|^2 \right| \leq \frac{Kc^{3/4}}{(1+\tilde{r})^{1+1/4}}$, hence

$$\left| \int_{B(d\vec{e}_1, d^{\varepsilon'})} \Re((|Q_c|^2 - |V|^2)\partial_d V \overline{\partial_c \Gamma_{c,d}}) \right| \leq K \int_{B(d\vec{e}_1, d^{\varepsilon'})} \frac{c^{1/4}}{(1+r_1)^{2+3/4}} = o_{c \rightarrow 0}(1).$$

We have from (3.12) that $|Q_c - V| \leq \frac{c^{3/4}}{(1+\bar{r})^{1/4}}$, and, in $B(d\bar{e}_1^\rightarrow, d^{\varepsilon'})$, we have (by Lemmas 2.1 and 2.2) that $|\Re(\bar{V}\partial_d V)| \leq \frac{K}{(1+r_1)^3}$, therefore

$$\left| \int_{B(d\bar{e}_1^\rightarrow, d^{\varepsilon'})} \Re(2\Re(\bar{V}\partial_d V)(Q_c - V)\overline{\partial_c \Gamma_{c,d}}) \right| \leq K \int_{B(d\bar{e}_1^\rightarrow, d^{\varepsilon'})} \frac{c^{1/4}}{(1+r_1)^{3+3/4}} = o_{c \rightarrow 0}(1).$$

Finally, by using the same estimates, we have

$$\left| \int_{B(d\bar{e}_1^\rightarrow, d^{\varepsilon'})} \Re(2\Re(\overline{Q_c - V}\partial_d V)Q_c\overline{\partial_c \Gamma_{c,d}}) \right| \leq K \int_{B(d\bar{e}_1^\rightarrow, d^{\varepsilon'})} \frac{c^{3/4}}{(1+r_1)^{1+1/4}} |\Re(Q_c\overline{\partial_c \Gamma_{c,d}})|.$$

We compute

$$\Re(Q_c\overline{\partial_c \Gamma_{c,d}}) = \Re(V\overline{\partial_c \Gamma_{c,d}}) + \Re(\Gamma_{c,d}\overline{\partial_c \Gamma_{c,d}}).$$

By using $\left\| \frac{\partial_c \Gamma_{c,d}}{V} \right\|_{*,1/2,d} \leq K(\sigma)c^{-1/2-\sigma}$ from Lemma 3.2 and (4.23), we have $|\Re(V\overline{\partial_c \Gamma_{c,d}})| \leq \frac{K(\sigma)c^{-1/2-\sigma}}{(1+r_1)^{3/2}}$.

Furthermore, with $|\Gamma_{c,d}| \leq \frac{Kc^{1/2}}{(1+r_1)^{1/2}}$, we have $|\Re(\Gamma_{c,d}\overline{\partial_c \Gamma_{c,d}})| \leq \frac{K(\sigma)c^{-\sigma}}{(1+r_1)}$. With these estimates, we infer, taking $\sigma > 0$ small enough,

$$\left| \int_{B(d\bar{e}_1^\rightarrow, d^{\varepsilon'})} \Re(2\Re(\overline{Q_c - V}\partial_d V)Q_c\overline{\partial_c \Gamma_{c,d}}) \right| = o_{c \rightarrow 0}(1)$$

which ends the proof of

$$\int_{B(d\bar{e}_1^\rightarrow, d^{\varepsilon'})} \Re((\tilde{L}_{Q_c} - L_V)(\partial_d V)\overline{\partial_c \Gamma_{c,d}}) = o_{c \rightarrow 0}(1).$$

Step 3. Conclusion.

We showed that

$$\partial_c d_c = \frac{2\pi + o_{c \rightarrow 0}(1)}{\frac{-2\pi}{d_c^2} + o_{d_c \rightarrow \infty}\left(\frac{1}{d_c^2}\right)},$$

therefore, with $d_c = \frac{1+o_{c \rightarrow 0}(1)}{c}$ from Proposition 2.26 we have

$$\partial_c d_c = -\frac{1 + o_{c \rightarrow 0}(1)}{c^2}.$$

As a result of subsection 4.5, at fixed c ,

$$\partial_d \int_{B(d\bar{e}_1^\rightarrow, d^{\varepsilon'}) \cup B(-d\bar{e}_1^\rightarrow, d^{\varepsilon'})} \Re(\overline{\partial_d V} \text{TW}_c(Q_{c,d}))|_{d=d_c} \neq 0$$

for c small enough. By the implicit function theorem, taking some $0 < c_* < c_0(\sigma)$, we can construct a C^1 branch $c \mapsto d_c$ in a vicinity of c_* . We define \mathbf{C} as the set of $c_* > c_\otimes \geq 0$ such that there exists a C^1 branch $c \mapsto d_c$ on $]c_\otimes, c_*[$. We have just shown that \mathbf{C} is not empty. Let us suppose that $c_\otimes := \inf \mathbf{C} \neq 0$. Then, $c \mapsto d_c$ is uniformly bounded on $]c_\otimes, c_*[$ in C^1 by subsection 4.6, and can therefore be extended by continuity to c_\otimes , and we denote d_\otimes its value there. We can construct the perturbation $\Phi_{c_\otimes, d_\otimes}$ by continuity since $c, d \mapsto \Phi_{c,d}$ are C^1 functions in the Banach space $\{\Phi \in C^1(\mathbb{R}^2, \mathbb{C}), \|\Phi\|_{*,\sigma, d_\otimes} < +\infty\}$ for its canonical norm (which is equivalent to $\|\cdot\|_{*,\sigma, d}$ for any $d \in [d_\otimes, d_{c_*}]$). By passing to the limit, we have $\|\Phi_{c_\otimes, d_\otimes}\|_{*,\sigma, d_\otimes} \leq K_0(\sigma, \sigma')c_\otimes^{1-\sigma'}$ for $K_0(\sigma, \sigma')$ defined in Proposition 2.21. By continuity of λ , we check that we have $\lambda(c_\otimes, d_\otimes) = 0$ (for the perturbation $\Phi_{c_\otimes, d_\otimes}$). Therefore, by the implicit function theorem, there exists a unique branch $c \mapsto d_c$ in a vicinity of (c_\otimes, d_\otimes) such that $\lambda(c, d_c) = 0$. This branch, by uniqueness, corresponds to the branch we had on $]c_\otimes, c_*[$, and is also C^1 by the implicit function theorem. Therefore $\inf \mathbf{C} < c_\otimes$, which is in contradiction with $c_\otimes = \inf \mathbf{C}$, and thus $\inf \mathbf{C} = 0$.

In particular, the travelling wave Q_c on this branch is uniquely defined by this construction and is a C^1 function of c . Indeed, we shall now show that there is only one choice of d_c such that $\lambda(c, d_c) = 0$ in $]\frac{1}{2c}, \frac{2}{c}[$. If there exist $d_1 \neq d_2$ in $]\frac{1}{2c}, \frac{2}{c}[$ such that $\lambda(c, d_1) = \lambda(c, d_2) = 0$, by Subsection 4.5, we have

$$\partial_d(\lambda(c, d))|_{d=d_1} < 0 \quad \text{and} \quad \partial_d(\lambda(c, d))|_{d=d_2} < 0,$$

therefore, there exists d' such that $\lambda(c, d') = 0$ and $\partial_d(\lambda(c, d'))|_{d=d'} \geq 0$, but then, since $\lambda(c, d') = 0$, we have $\partial_d(\lambda(c, d))|_{d=d'} < 0$, which is in contradiction with $\partial_d(\lambda(c, d'))|_{d=d'} \geq 0$. Now that we have uniqueness in the choice of d_c (in $]\frac{1}{2c}, \frac{2}{c}[$), we have uniqueness of $\Phi_{c,d}$ in the set

$$\{\Phi \in C^1(\mathbb{R}^2, \mathbb{C}), \|\Phi\|_{*,\sigma,d_c} \leq K_0(\sigma, \sigma')c^{1-\sigma'}\}$$

for $K_0(\sigma, \sigma') > 0$ defined in Proposition 2.21.

4.7 Proof of the estimate on $\partial_c Q_c$

We conclude the proof of Theorem 1.1 with the following lemma.

Lemma 4.6 *For any $0 < \sigma < 1$, there exist $c_0(\sigma) > 0$ such that for any $c < c_0(\sigma)$,*

$$\left\| \frac{\partial_c Q_c}{V} + \left(\frac{1 + o_{c \rightarrow 0}(1)}{c^2} \right) \frac{\partial_d V|_{d=d_c}}{V} \right\|_{*,\sigma,d_c} = o_{c \rightarrow 0} \left(\frac{1}{c^2} \right).$$

With this estimate and by using the same computations as in the proof of Lemma 3.6, we show that

$$\left\| \partial_c Q_c + \left(\frac{1 + o_{c \rightarrow 0}(1)}{c^2} \right) \partial_d (V_1(\cdot - d\vec{e}_1) V_{-1}(\cdot + d\vec{e}_1))|_{d=d_c} \right\|_p = o_{c \rightarrow 0} \left(\frac{1}{c^2} \right).$$

for all $+\infty \geq p > 2$ if c is small enough, which ends the proof of Theorem 1.1.

Proof From subsection 4.5, we know that Q_c is a C^1 function of c . We have $Q_c = V + \Gamma_{c,d_c}$, hence

$$\partial_c Q_c = \partial_c V + \partial_c(\Gamma_{c,d_c}) = \frac{-1 + o_{c \rightarrow 0}(1)}{c^2} \partial_d V + \partial_c(\Gamma_{c,d_c}),$$

where we used $\partial_c V = \left(-\frac{1}{c^2} + o_{c \rightarrow 0}\left(\frac{1}{c^2}\right)\right) \partial_d V$ thanks to subsection 4.6. Γ_{c,d_c} depends on c directly and through d_c . We will write $\partial_c \Gamma_{c,d_c}$ for the derivatives with respect to c but at a fixed d_c , and $\partial_d \Gamma_{c,d_c}$ for the derivate with respect to d_c but at fixed c . In particular,

$$\partial_c(\Gamma_{c,d_c}) = \partial_c \Gamma_{c,d_c} + \partial_c d_c \partial_d \Gamma_{c,d_c}.$$

From Lemma 3.9 and (4.23), we showed that

$$\left\| \frac{\partial_c \Gamma_{c,d_c}}{V} \right\|_{*,\sigma,d_c} \leq K(\sigma, \sigma')c^{-\sigma'},$$

and from Lemma 3.3 with the definition of $\Gamma_{c,d}$, we show easily that

$$\left\| \frac{\partial_d \Gamma_{c,d_c}}{V} \right\|_{*,\sigma,d_c} \leq K(\sigma, \sigma')c^{1-\sigma'}.$$

Finally, from subsection 4.6, we have $\partial_c d_c = \frac{1 + o_{c \rightarrow 0}(1)}{c^2}$, therefore

$$\left\| \frac{\partial_c(\Gamma_{c,d_c})}{V} \right\|_{*,\sigma,d_c} \leq K(\sigma, \sigma')(c^{-\sigma'} + c^{-2}(1 + o_{c \rightarrow 0}(1))c^{1-\sigma'}) = o_{c \rightarrow 0} \left(\frac{1}{c^2} \right)$$

since $0 < \sigma < \sigma' < 1$. □

This concludes the proof of Lemma 4.6, which itself concludes the proof of Theorem 1.1.

References

- [1] Milton Abramowitz and Irene A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] Fabrice Bethuel, Robert L. Jerrard, and Didier Smets. On the NLS dynamics for infinite energy vortex configurations on the plane. *Rev. Mat. Iberoam.*, 24(2):671–702, 2008.
- [3] Fabrice Bethuel and Jean-Claude Saut. Travelling waves for the Gross-Pitaevskii equation. I. *Ann. Inst. H. Poincaré Phys. Théor.*, 70(2):147–238, 1999.
- [4] Fabrice Bethuel, Philippe Gravejat, and Jean-Claude Saut. Travelling waves for the Gross-Pitaevskii equation. II. *Comm. Math. Phys.*, 285(2):567–651, 2009.
- [5] Xinfu Chen, Charles M. Elliott, and Tang Qi. Shooting method for vortex solutions of a complex-valued Ginzburg-Landau equation. *Proc. Roy. Soc. Edinburgh Sect. A*, 124(6):1075–1088, 1994.
- [6] David Chiron and Mihai Mariş. Traveling waves for nonlinear Schrödinger equations with nonzero conditions at infinity. *Arch. Ration. Mech. Anal.*, 226(1):143–242, 2017.
- [7] Manuel del Pino, Patricio Felmer, and Michał Kowalczyk. Minimality and nondegeneracy of degree-one Ginzburg-Landau vortex as a Hardy’s type inequality. *Int. Math. Res. Not.*, (30):1511–1527, 2004.
- [8] Manuel del Pino, Michał Kowalczyk, and Monica Musso. Variational reduction for Ginzburg-Landau vortices. *J. Funct. Anal.*, 239(2):497–541, 2006.
- [9] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [10] V. L. Ginzburg and L. P. Pitaevskii. On the theory of superfluidity. *Soviet Physics. JETP*, 34 (7):858–861, 1958.
- [11] Philippe Gravejat. Decay for travelling waves in the Gross-Pitaevskii equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 21(5):591–637, 2004.
- [12] Philippe Gravejat. Asymptotics for the travelling waves in the Gross-Pitaevskii equation. *Asymptot. Anal.*, 45(3-4):227–299, 2005.
- [13] Philippe Gravejat. First order asymptotics for the travelling waves in the Gross-Pitaevskii equation. *Adv. Differential Equations*, 11(3):259–280, 2006.
- [14] Rose-Marie Hervé and Michel Hervé. Étude qualitative des solutions réelles d’une équation différentielle liée à l’équation de Ginzburg-Landau. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 11(4):427–440, 1994.
- [15] C A Jones and P H Roberts. Motions in a bose condensate. IV. axisymmetric solitary waves. *Journal of Physics A: Mathematical and General*, 15(8):2599–2619, aug 1982.
- [16] Fanghua Lin and Juncheng Wei. Traveling wave solutions of the Schrödinger map equation. *Comm. Pure Appl. Math.*, 63(12):1585–1621, 2010.
- [17] Yong Liu and Juncheng Wei. Multi-vortex traveling waves for the gross-pitaevskii equation and the adler-moser polynomials. 2018.
- [18] John C. Neu. Vortices in complex scalar fields. *Phys. D*, 43(2-3):385–406, 1990.

Laboratoire de Mathématiques J.A. Dieudonné
UMR 7351 CNRS UNS
Université de Nice - Sophia Antipolis 06108 Nice Cedex 02, France
david.chiron@univ-cotedazur.fr
eliot.pacherie@univ-cotedazur.fr