

ENTROPY OF NON-EQUILIBRIUM STATIONARY MEASURES OF BOUNDARY DRIVEN TASEP

CÉDRIC BERNARDIN, PATRÍCIA GONÇALVES, AND CLAUDIO LANDIM

ABSTRACT. We examine the entropy of non-equilibrium stationary states of boundary driven totally asymmetric simple exclusion processes. As a consequence, we obtain that the Gibbs-Shannon entropy of the non equilibrium stationary state converges to the Gibbs-Shannon entropy of the local equilibrium state. Moreover, we prove that its fluctuations are Gaussian, except when the mean displacement of particles produced by the bulk dynamics agrees with the particle flux induced by the density reservoirs in the maximal phase regime.

1. INTRODUCTION

The boundary driven simple exclusion process (SEP) is a fundamental model of non-equilibrium statistical physics and the progress accomplished during the last decade in our understanding of non-equilibrium stationary states (NESS) comes, for a large part, from its study. In this work we will be concerned with the one-dimensional boundary driven asymmetric simple exclusion process (ASEP) which is a lattice gas model connected to particle reservoirs and in the bulk particles perform biased random walks respecting the exclusion rule (that is at most one particle per site). Despite its simplicity, the ASEP shows a range of non trivial macroscopic phenomena such as phase transitions ([9]) which is one of the striking properties of non-equilibrium steady states, since as it is well known one-dimensional systems at equilibrium with short-range interactions cannot exhibit phase transitions.

In [3] the entropy function of NESS has been proposed in analogy with the usual Boltzmann entropy for equilibrium stationary states and it has been explicitly computed for the boundary driven symmetric simple exclusion process. In the present paper we extend that work to the boundary driven ASEP and show that the entropy function detects phase transitions.

The boundary driven ASEP is defined as follows. Let $p = 1 - q \in [0, 1] \neq 1/2$ and $0 < \rho_- < \rho_+ < 1$. The microstates are described by the vectors $\eta = (\eta_{-N}, \dots, \eta_N) \in \Omega_N := \{0, 1\}^{\{-N, \dots, N\}}$ where for $x \in \{-N, \dots, N\}$, $\eta_x = 1$ if the site x is occupied and $\eta_x = 0$ if the site x is empty. In the bulk of the system, each particle, independently from the others, performs a nearest-neighbor asymmetric random walk, where jumps to the right (resp. left) neighboring site occur at rate p (resp. rate q), with the convention that each time a particle attempts to jump to a site already occupied, the jump is suppressed in order to respect the exclusion constrain. At the two boundaries the dynamics is modified to mimic the coupling with reservoirs of particles: if the site $-N$ is empty (resp. occupied), a particle is injected at rate α (resp. removed at rate γ); similarly, if the site N is empty, a particle is injected at rate δ (resp. removed at rate β). For any sites $x \neq y$, we denote by $\sigma^{x,y}\eta$ (resp. $\sigma^x(\eta)$) the configuration obtained from $\eta \in \Omega_N$ by the exchange of the occupation variables η_x and η_y (resp. by the change of η_x into $1 - \eta_x$). The boundary

Key words and phrases. Non-equilibrium stationary states, phase transitions, large deviations, quasi-potential, boundary driven asymmetric exclusion processes.

driven (nearest neighbor) asymmetric simple exclusion process is the Markov process on Ω whose generator L is given by

$$L = L_0 + L_- + L_+,$$

where L_0, L_-, L_+ act on functions $f : \Omega_N \rightarrow \mathbb{R}$ as follows

$$(L_0 f)(\eta) = \sum_{x=-N}^{N-1} \{p\eta_x(1-\eta_{x+1}) + q\eta_{x+1}(1-\eta_x)\} [f(\sigma^{x,x+1}\eta) - f(\eta)],$$

$$(L_- f)(\eta) = c_-(\eta_{-N}) [f(\sigma^{-N}\eta) - f(\eta)], \quad (L_+ f)(\eta) = c_+(\eta_N) [f(\sigma^N\eta) - f(\eta)]$$

with $c_{\pm} : \Omega_N \rightarrow [0, +\infty)$ given by

$$c_-(\eta) = \alpha(1-\eta_{-N}) + \gamma\eta_{-N}, \quad c_+(\eta) = \delta(1-\eta_N) + \beta\eta_N.$$

The density of the left (resp. right) reservoir is denoted by ρ_- (resp. ρ_+) and can be explicitly computed as a function of p, q, α, γ (resp. p, q, β, δ). For simplicity we will focus only on the totally asymmetric simple exclusion process (TASEP) that corresponds to $p = 0$ or $p = 1$. Furthermore, if $p = 1 - q = 1$ we take $\alpha = \rho_-, \beta = 1 - \rho_+, \gamma = \delta = 0$. If $p = 1 - q = 0$, we take $\delta = \rho_+, \gamma = 1 - \rho_-$ and $\alpha = \beta = 0$. Since $\rho_- < \rho_+$ the reservoirs induce a flux of particles from the right to the left. On the other hand the bulk dynamics produces a mean displacement of the particles with a drift equal to $(p - q)$. For $p = 0$ both effects cooperate to push the particles to the left and we call the corresponding system the *cooperative* TASEP. If $p = 1$ the two effects push the particles in opposite directions and we call the corresponding system the *competitive* TASEP.

The unique non-equilibrium stationary state of the boundary driven TASEP is denoted by $\mu_{ss,N}$. In the case $\rho_- = \rho_+ = \rho \in (0, 1)$, $\mu_{ss,N}$ is given by the Bernoulli product measure ν_ρ on Ω_N . In the non-equilibrium situation, the steady state has a lot of non-trivial interesting properties. The phase diagram for the average density $\bar{\rho}$ is well known and one can distinguish three phases: the high-density phase (HD) for which $\bar{\rho} = \rho_+$, the low density phase (LD) for which $\bar{\rho} = \rho_-$ and the maximal current phase (MC) where $\bar{\rho} = 1/2$. The transition lines between these phases are second order phase transitions except for the boundary $\rho_- + \rho_+ = 1$ in the competitive case where the transition is of first order. On this line, the typical configurations are shocks between LD phase with density ρ_- at the left of the shock and HD phase with density ρ_+ at the right of the shock. The position of the shock is uniformly distributed along the system and the average profile $\bar{\rho}(x)$ is given by $\bar{\rho}(x) = \rho_- \mathbf{1}\{x \leq 0\} + \rho_+ \mathbf{1}\{x \geq 0\}$. This is summarized in Figure 1.




FIGURE 1. The phase diagram for the cooperative TASEP (resp. left) and the competitive TASEP (resp. right).

The entropy function of $\mu_{ss,N}$ introduced in [3] is the function $S_{\rho_-, \rho_+}^{p,q} : [0, +\infty) \rightarrow [0, \log(2)] \cup \{-\infty\}$ defined by

$$S(E) = \lim_{\delta \rightarrow 0} \lim_{N \rightarrow +\infty} \frac{1}{2N+1} \log \left(\sum_{\eta \in \Omega_N} \mathbf{1} \left\{ \left| \frac{1}{2N+1} \log(\mu_{ss,N}(\eta)) + E \right| \leq \delta \right\} \right)$$

if the limit exists.

Observe that $J_{\rho_-, \rho_+}^{p,q}(E) = E - S_{\rho_-, \rho_+}^{p,q}(E) \geq 0$ coincides with the large deviations function of the random variables $Y_N(\eta) := -\frac{1}{2N+1} \log(\mu_{ss,N}(\eta))$ under the probability measure $\mu_{ss,N}$. Then the (concave) Legendre transform

$$P(\theta) := P_{\rho_-, \rho_+}^{p,q}(\theta) = \inf_{E \geq 0} \{ \theta E - S_{\rho_-, \rho_+}^{p,q}(E) \} \quad (1.1)$$

of $S_{\rho_-, \rho_+}^{p,q}$, that we call the *pressure*, is by the Laplace-Varadhan theorem simply related to the cumulant generating function of the random variables $\{Y_N\}_N$, i.e.

$$\begin{aligned} P(\theta) &= - \lim_{N \rightarrow +\infty} \frac{1}{2N+1} \log \left(\int e^{(2N+1)(1-\theta)Y_N(\eta)} \mu_{ss,N}(d\eta) \right) \\ &= - \lim_{N \rightarrow +\infty} \frac{1}{2N+1} \log \left(\sum_{\eta \in \Omega_N} (\mu_{ss,N}(\eta))^\theta \right). \end{aligned} \quad (1.2)$$

In the case $\rho_- = \rho_+ = \rho \in (0, 1)$, denoting by

$$\varphi = \log\left(\frac{\rho}{1-\rho}\right) \in \mathbb{R} \quad (1.3)$$

the corresponding chemical potential, it is easy to show that the entropy function is given by

$$S_{\rho, \rho}(E) = -s \left(\frac{-E + \log(1 + e^\varphi)}{\varphi} \right) \quad (1.4)$$

where $s(\theta) = \theta \log(\theta) + (1-\theta) \log(1-\theta)$. The pressure $P(\varphi, \theta)$ is then given by

$$P(\varphi, \theta) := \theta \log(1 + e^\varphi) - \log(1 + e^{\varphi\theta}). \quad (1.5)$$

In the non-equilibrium case $\rho_- \neq \rho_+$, since $\mu_{ss,N}$ has not a simple form, the computation of the entropy function is much more difficult. It has been proved in [3] that if a strong form of local equilibrium holds (see Section 4 for a precise definition), then the entropy function $S := S_{\rho_-, \rho_+}^{p,q}$ can be expressed in a variational form involving the non-equilibrium free energy $V := V_{\rho_-, \rho_+}^{p,q}$ and the Gibbs-Shannon entropy \mathbb{S} :

$$S(E) = \sup_{\rho \in \mathcal{M}} \{ \mathbb{S}(\rho) ; V(\rho) + \mathbb{S}(\rho) = E \}. \quad (1.6)$$

The interval composed of the $E \in [0, +\infty)$ such that $S(E) \neq -\infty$ is called the *energy band*. The bottom and the top of the energy band are defined respectively by

$$E^- := \inf_{\rho \in \mathcal{M}} \{ \mathbb{S}(\rho) + V(\rho) \} \quad \text{and} \quad E^+ := \sup_{\rho \in \mathcal{M}} \{ \mathbb{S}(\rho) + V(\rho) \} \quad (1.7)$$

The set \mathcal{M} is given in (2.1). The Gibbs-Shannon entropy of the profile $\rho \in \mathcal{M}$ is defined by

$$\mathbb{S}(\rho) = -\frac{1}{2} \int_{-1}^1 s(\rho(x)) dx. \quad (1.8)$$

The non-equilibrium free energy is the large deviation function of the empirical density under $\mu_{ss,N}$. Its value does not depend on p nor q but only on the sign of $p - q$. The explicit computation of this functional has been obtained first in [8] and generalized to other systems in [2]. Thus we denote it by V^+ if $p - q > 0$ and by V^- if $p - q < 0$. Similarly, the entropy (resp. pressure) of the competitive TASEP is denoted by S^+ (resp. P^+) and the entropy (resp. pressure) of the cooperative TASEP by S^- (resp. P^-). It follows easily from (1.1) and (1.6) that

$$P(\theta) = \inf_{\rho \in \mathcal{M}} \{ \theta(V(\rho) + \mathbb{S}(\rho)) - \mathbb{S}(\rho) \}.$$

This formula can also be obtained starting from (1.2) and using the local equilibrium statement as it is done in [3] for the entropy function.

In this paper we compute explicitly S^+ and S^- (resp. Theorem 2.2 and Theorem 3.2) and P^+ and P_- (resp. Theorem 2.4 and Theorem 3.4). From those results we deduce several interesting consequences (see Theorem 2.5 and Theorem 3.5):

- We recover some results of [1] for the TASEP, showing that the Gibbs-Shannon entropy of the TASEP is the same, in the thermodynamic limit, as the Gibbs-Shannon entropy of the local Gibbs equilibrium measure, which is nothing but the Bernoulli product measure $\otimes_{x=-N}^N \mathcal{B}(\bar{\rho}(x))$ where $\mathcal{B}(r)$ is the one-site Bernoulli measure on $\{0, 1\}$ with density r .
- For the competitive TASEP, contrarily to what happens for the boundary driven symmetric simple exclusion process ([3, 6]), the first order corrections are identical for the non-equilibrium stationary state $\mu_{ss,N}$ and for the local equilibrium state $V_{\bar{\rho}}$.
- For the cooperative TASEP, the same occurs if $\rho_-, \rho_+ \leq 1/2$ or if $\rho_-, \rho_+ \geq 1/2$. But in the MC phase $\rho_- \leq 1/2 \leq \rho_+$, the fluctuations are not Gaussian. This is reminiscent of [5, 8] where it is shown that the fluctuations of the density are non-Gaussian¹.

Our last results concern the presence of phase transitions² for the competitive and the cooperative TASEP. For the cooperative TASEP the function S^- is a continuously differentiable concave function on its energy band but has linear parts. As a consequence the pressure function P^- is a concave function with a discontinuous derivative. The function P^- may also have a linear part due to the fact that the entropy S^- does not necessarily vanish at the boundaries of the energy band. The function S^+ is a smooth concave function on its energy band if $\rho_- \leq 1/2 \leq \rho_+$ but does not vanish at the top of the energy band. Consequently the pressure function P^+ is concave with a linear part on an infinite interval. If $\rho_-, \rho_+ \leq 1/2$ or $\rho_-, \rho_+ \geq 1/2$, the entropy function S^+ has a discontinuity of its derivative at some point in the interior of the energy band but vanishes at the boundaries of the energy band. Then, the pressure function P^+ has a linear part on a finite interval.

It would be interesting to see how these results extend to other asymmetric systems for which the quasi-potential has been explicitly computed ([2]). The form of the entropy function obtained for the TASEP is relatively simple but follows from long computations. We did not succeed in giving a simple intuitive explanation to the final formulas obtained. We also notice that extending these results to a larger class of systems would require to prove the strong form of local equilibrium for them in order to get (1.6). This seems to be

¹The non-Gaussian part of the fluctuations can be described in terms of the statistical properties of a Brownian excursion ([5]).

²We refer the interested reader to [10] for more informations about the implications of these facts from a physical viewpoint.

a difficult task.

The paper is organized as follows. In Section 2 we obtain the entropy and the pressure functions for the competitive TASEP and deduce some consequences of these computations. In Section 3 we obtain similar results for the cooperative TASEP. The local equilibrium statement is proved in Section 4. Technical parts are postponed to the Appendix.

2. COMPETITIVE TASEP

In this section we compute (1.6) for the competitive TASEP. Let $\chi : [0, 1] \rightarrow [0, 1]$ be defined by $\chi(\rho) = \rho(1 - \rho)$. The chemical potential corresponding to ρ_{\pm} is denoted by φ_{\pm} and satisfies $\rho_{\pm} = e^{\varphi_{\pm}} / (1 + e^{\varphi_{\pm}})$, see (1.3).

We consider the set $\mathbb{L}^{\infty}([-1, 1])$ equipped with the weak* topology and \mathcal{M} as the set

$$\mathcal{M} = \{\rho \in \mathbb{L}^{\infty}([-1, 1]) : 0 \leq \rho \leq 1\} \quad (2.1)$$

which is equipped with the relative topology. Let

$$\Phi = \left\{ \varphi := \varphi_y : x \in [-1, 1] \rightarrow \varphi_- \mathbf{1}\{-1 \leq x < y\} + \varphi_+ \mathbf{1}\{y \leq x \leq 1\} ; y \in [-1, 1] \right\}.$$

For $(\rho, \varphi) \in \mathcal{M} \times \Phi$ we define the functional

$$\mathcal{H}(\rho, \varphi) = \frac{1}{2} \int_{-1}^1 \left[(1 - \rho(x))\varphi(x) - \log(1 + e^{\varphi(x)}) \right] dx. \quad (2.2)$$

Then the quasi-potential of the competitive TASEP is given ([8], [2]) by

$$V^+(\rho) = -\mathbb{S}(\rho) + \inf_{\varphi \in \Phi} \mathcal{H}(\rho, \varphi) - \bar{V}^+$$

where

$$\bar{V}^+ = -\mathbb{S}(\bar{\rho}) + \inf_{\varphi \in \Phi} \mathcal{H}(\bar{\rho}, \varphi) = \log \left(\min_{\rho \in [\rho_-, \rho_+]} \chi(\rho) \right).$$

We recall that $\bar{\rho} = \rho_-$ for $\rho_+ < 1 - \rho_-$, i.e. $\varphi_+ < -\varphi_-$, $\bar{\rho} = \rho_+$ for $\rho_+ > 1 - \rho_-$, i.e. $\varphi_+ > -\varphi_-$ and $\bar{\rho}(x) = \rho_- \mathbf{1}\{x \leq 0\} + \rho_+ \mathbf{1}\{x > 0\}$ if $\rho_- + \rho_+ = 1$. Let $\bar{\varphi} = \sup(\varphi_+, -\varphi_-)$ so that $\bar{\rho} = e^{\bar{\varphi}} / (1 + e^{\bar{\varphi}})$ if $\rho_- + \rho_+ \neq 1$. Let us also introduce $\varphi_0 = \sup\{|\varphi_-|, |\varphi_+|\}$ and $\rho_0 = e^{\varphi_0} / (1 + e^{\varphi_0})$.

For each $E \geq 0$, $m \in [0, 2]$ and φ_-, φ_+ we define

$$\xi_0 := \frac{\log(1 + e^{\varphi_+}) - \log(1 + e^{\varphi_-})}{\varphi_+ - \varphi_-} \in (0, 1), \quad \hat{\xi}_0 := \frac{\log(1 + e^{\varphi_+}) + \log(1 + e^{\varphi_-})}{\varphi_+ - \varphi_-}. \quad (2.3)$$

2.1. Energy bands. In this section we determine the energy band of the competitive TASEP. This is summarized in Figure 2.

Proposition 2.1. *The bottom of the energy band is given by*

$$E_{+\infty}^- = -\bar{V}^+ - \log(1 + e^{\varphi_0})$$

and the top of the energy band is given by

$$E_{+\infty}^+ = -\bar{V}^+ + \begin{cases} \frac{\varphi_- \log(1 + e^{\varphi_+}) - \varphi_+ \log(1 + e^{\varphi_-})}{\varphi_+ - \varphi_-}, & \rho_- \leq \frac{1}{2} \leq \rho_+, \\ -\log(1 + e^{\varphi_+}), & \rho_- < \rho_+ \leq \frac{1}{2}, \\ -\log(1 + e^{-\varphi_-}), & \frac{1}{2} \leq \rho_- < \rho_+. \end{cases}$$

Proof. Recall from (1.7) that $E_{+\infty}^- = \inf_{\rho \in \mathcal{M}} \{\mathbb{S}(\rho) + V^+(\rho)\}$. The computation of the bottom of the energy band is easy since we have

$$\begin{aligned} E_{+\infty}^- &= \inf_{\rho \in \mathcal{M}} \inf_{\varphi \in \Phi} \mathcal{H}(\rho, \varphi) - \bar{V}^+ \\ &= \inf_{\varphi \in \Phi} \inf_{\rho \in \mathcal{M}} \mathcal{H}(\rho, \varphi) - \bar{V}^+ \\ &= \inf_{y \in [-1, 1]} \frac{1}{2} \left\{ (y+1)(\varphi_- \wedge 0 - \log(1 + e^{\varphi_-})) + (1-y)(\varphi_+ \wedge 0 - \log(1 + e^{\varphi_+})) \right\} - \bar{V}^+ \\ &= \inf \left\{ (\varphi_- \wedge 0 - \log(1 + e^{\varphi_-})), (\varphi_+ \wedge 0 - \log(1 + e^{\varphi_+})) \right\} - \bar{V}^+. \end{aligned}$$

We now compute the top of the energy band. Recall from (1.7) that $E_{+\infty}^+ = \sup_{\rho \in \mathcal{M}} \{\mathbb{S}(\rho) + V^+(\rho)\}$. For each profile $\rho \in \mathcal{M}$, we introduce the non-decreasing continuous and almost everywhere differentiable function

$$H_\rho(x) = \int_{-1}^x (1 - \rho(z)) dz \quad (2.4)$$

and the constant $m_\rho = H_\rho(1) \in [0, 2]$. Let $y_\rho \in [-1, 1]$ the infimum of the points of $[-1, 1]$ where the infimum of the continuous function $y \rightarrow y\xi_0 - H_\rho(y)$ is attained. Then,

$$\begin{aligned} E_{+\infty}^+ &= \sup_{\rho \in \mathcal{M}} \inf_{\varphi \in \Phi} \mathcal{H}(\rho, \varphi) - \bar{V}^+ \\ &= \frac{(\varphi_+ - \varphi_-)}{2} \sup_{m \in [0, 2]} \left\{ \frac{\varphi_+}{\varphi_+ - \varphi_-} m - \frac{2\bar{V}^+}{\varphi_+ - \varphi_-} - \hat{\xi}_0 + \mathbb{U}(m) \right\} \end{aligned}$$

with

$$\mathbb{U}(m) = \sup_{\substack{\rho \in \mathcal{M}, \\ H_\rho(1) = m}} \inf_{y \in [-1, 1]} \{y\xi_0 - H_\rho(y)\}.$$

We claim that $\mathbb{U}(m) = \inf\{-\xi_0, \xi_0 - m\}$. It is trivial that $\mathbb{U}(m) \leq \inf\{-\xi_0, \xi_0 - m\}$ (take $y = 1, -1$ in the variational formula). For $m \neq \xi_0$, the supremum can be obtained by taking the piecewise linear function H_ρ such that H_ρ is linear on $[-1, 0]$ and on $[0, 1]$, with $H_\rho(-1) = 0, H_\rho(1) = m$ and

$$H_\rho(0) = \begin{cases} \xi_0, & -\xi_0 \leq \xi_0 - m, \quad m > \xi_0, \\ 0, & -\xi_0 \leq \xi_0 - m, \quad m < \xi_0, \\ 1, & -\xi_0 \geq \xi_0 - m, \quad m - \xi_0 \geq 1, \\ m - \xi_0, & -\xi_0 \geq \xi_0 - m, \quad m - \xi_0 \leq 1. \end{cases}$$

In the case $m = \xi_0$, there are two profiles H_ρ for which the supremum is obtained, one with $H_\rho(0) = \xi_0$, the other one with $H_\rho(0) = 0$. It follows that

$$\begin{aligned} E_{+\infty}^+ &= \frac{(\varphi_+ - \varphi_-)}{2} \sup_{m \in [0, 2]} \left\{ \frac{\varphi_+}{\varphi_+ - \varphi_-} m - \frac{2\bar{V}^+}{\varphi_+ - \varphi_-} - \hat{\xi}_0 + \inf\{\xi_0 - m, -\xi_0\} \right\} \\ &= \frac{(\varphi_+ - \varphi_-)}{2} \left[\sup_{m \in [2\xi_0, 2]} \left\{ \frac{\varphi_-}{\varphi_+ - \varphi_-} m + (\xi_0 - \hat{\xi}_0) \right\} \vee \sup_{m \in [0, 2\xi_0]} \left\{ \frac{\varphi_+}{\varphi_+ - \varphi_-} m - (\xi_0 + \hat{\xi}_0) \right\} \right] \\ &\quad - \bar{V}^+. \end{aligned}$$

We have now to optimize a piecewise linear function and we get the result. \square




FIGURE 2. The phase diagram for the bottom of the energy band (resp. left) and the top of the energy band (resp. right) for the competitive TASEP. We have $\underline{A} = -\log(1 - \rho_-)$, $\underline{B} = -\log(\rho_+)$ and $\bar{A} = -\log(\chi(\rho_-)) + \log(1 - \rho_+)$, $\bar{B} = \frac{\varphi_-}{\varphi_+ - \varphi_-} \log\left(\frac{1 - \rho_-}{1 - \rho_+}\right) - \log \rho_-$, $\bar{C} = \frac{\varphi_+}{\varphi_+ - \varphi_-} \log\left(\frac{1 - \rho_-}{1 - \rho_+}\right) - \log \rho_+$, $\bar{D} = -\log(\chi(\rho_+)) + \log(\rho_-)$. All transitions are of first order.

2.2. **Entropy.** Now we compute the entropy function. We introduce

$$W(\rho_-, \rho_+) = \frac{\log(\rho_+) \log(1 - \rho_-) - \log(\rho_-) \log(1 - \rho_+)}{\log\left(\frac{\rho_+(1 - \rho_-)}{\rho_-(1 - \rho_+)}\right)} \quad (2.5)$$

which corresponds to

$$W(\varphi_-, \varphi_+) = \frac{\varphi_- \log(1 + e^{\varphi_+}) - \varphi_+ \log(1 + e^{\varphi_-})}{\varphi_+ - \varphi_-} \quad (2.6)$$

and coincides with the first coordinate of one of the (possible) two intersection points of the curves $x \rightarrow S_{\rho_+, \rho_+}(-x)$ and $x \rightarrow S_{\rho_-, \rho_-}(-x)$.

Our aim is to prove

Theorem 2.2. *The restriction of the entropy function S^+ on the energy band $[E_{+\infty}^-; E_{+\infty}^+]$ is given by*

$$S^+(E) = \begin{cases} S_{\rho_0, \rho_0}(-(E + \bar{V}^+)), & \rho_- \leq \frac{1}{2} \leq \rho_+, \\ S_{\rho_-, \rho_-}(-(E + \bar{V}^+)) \mathbf{1}\{(E + \bar{V}^+) \leq W(\rho_-, \rho_+)\} \\ + S_{\rho_+, \rho_+}(-(E + \bar{V}^+)) \mathbf{1}\{(E + \bar{V}^+) > W(\rho_-, \rho_+)\}, & \rho_- < \rho_+ \leq \frac{1}{2}, \\ S_{\rho_-, \rho_-}(-(E + \bar{V}^+)) \mathbf{1}\{(E + \bar{V}^+) \geq W(\rho_-, \rho_+)\} \\ + S_{\rho_+, \rho_+}(-(E + \bar{V}^+)) \mathbf{1}\{(E + \bar{V}^+) < W(\rho_-, \rho_+)\}, & \frac{1}{2} \leq \rho_- < \rho_+, \end{cases}$$

and is a concave function.

The supremum in the definition of $S^+(E)$, see (1.6), for $E \in [E_{+\infty}^-; E_{+\infty}^+]$ is realized for a unique profile whose value is given by

$$\left\{ \begin{array}{l} u_{\bar{\rho}}, \quad \rho_- \leq \frac{1}{2} \leq \rho_+, \\ u_{\rho_-} \mathbf{1}\{(E + \bar{V}^+) \leq W(\rho_-, \rho_+)\} \\ + u_{\rho_+} \mathbf{1}\{(E + \bar{V}^+) > W(\rho_-, \rho_+)\}, \quad \rho_- < \rho_+ \leq \frac{1}{2}, \\ u_{\rho_-} \mathbf{1}\{(E + \bar{V}^+) \geq W(\rho_-, \rho_+)\} \\ + u_{\rho_+} \mathbf{1}\{(E + \bar{V}^+) < W(\rho_-, \rho_+)\}, \quad \frac{1}{2} \leq \rho_- < \rho_+, \end{array} \right.$$

where for any ρ , the profile u_{ρ} is the constant profile equal to $\frac{\log(\rho) - (E + \bar{V}^+)}{\log(\rho) - \log(1 - \rho)}$.




FIGURE 3. Graph of the function S^+ and graphs of the functions $E \rightarrow S_{\rho_{\pm}, \rho_{\pm}}(- (E + \bar{V}^+))$ for $\rho_- = 0.1$, $\rho_+ = 0.7$ (resp. left) and for $\rho_- = 0.1$, $\rho_+ = 0.3$ (resp. right).

Once the form of S^+ is obtained its concavity is easy to establish. The computation of S^+ is accomplished in several steps. For any $E \geq 0$, let $D := D(E)$ be the (possibly empty) compact convex domain of \mathbb{R}^2 defined by

$$(y, m) \in D \Leftrightarrow \left\{ \begin{array}{l} -1 \leq y \leq 1, \quad 0 \leq m \leq 2, \\ \sup\{\xi_0, m - \xi_0\} \leq E(m) \leq m - (m - 1)\xi_0, \\ 0 \leq y\xi_0 + E(m) \leq y + 1, \quad 0 \leq m - (y\xi_0 + E(m)) \leq 1 - y, \end{array} \right. \quad (2.7)$$

where

$$E(m) := \frac{\varphi_+ m - 2\bar{V}^+ - 2E}{\varphi_+ - \varphi_-} - \hat{\xi}_0.$$

The fact that D is a convex compact domain follows from the fact that $E(m)$ is a linear function of m so that D is the intersection of half-planes of \mathbb{R}^2 .

Define now the function $F : D \rightarrow \mathbb{R}$ on $(y, m) \in D$ by

$$F(y, m) = -(y+1)s \left(\frac{y\xi_0 + E(m)}{y+1} \right) - (1-y)s \left(\frac{m - (y\xi_0 + E(m))}{1-y} \right).$$

It is understood here that if $y = \pm 1$ then the indefinite terms are equal to 0. The function F is continuous on D and smooth on \mathring{D} .

Proposition 2.3. *The entropy function S^+ is given by*

$$S^+(E) = \frac{1}{2} \sup_{(y, m) \in D} F(y, m).$$

Proof. With the notations introduced above, we have

$$S^+(E) = \sup_{m \in [0, 2]} \sup_{\rho \in \mathcal{M}} \left\{ \mathbb{S}(\rho) ; m_\rho = m, \inf_{y \in [-1, 1]} \{y\xi_0 - H_\rho(y)\} = -E(m) \right\}.$$

Assume that there exists a profile $\rho_m \in \mathcal{M}$ such that $\inf_{y \in [-1, 1]} \{y\xi_0 - H_{\rho_m}(y)\} = -E(m)$ and $H_{\rho_m}(1) = m$. Then, by taking $y = \pm 1$ in the infimum we see that this is possible only if $-E(m) \leq \inf\{-\xi_0, \xi_0 - m\} = -\sup\{\xi_0, m - \xi_0\}$. Moreover, the existence of ρ_m implies that

$$-E(m) = \inf_{y \in [-1, 1]} \{y\xi_0 - H_{\rho_m}(y)\} \geq \inf_{\substack{\rho \in \mathcal{M}, \\ H_\rho(1) = m}} \inf_{y \in [-1, 1]} \{y\xi_0 - H_\rho(y)\}.$$

By inverting the two infimums, the right hand side of the previous inequality can be rewritten as

$$\inf_{y \in [-1, 1]} \left\{ y\xi_0 - \sup_{\substack{\rho \in \mathcal{M}, \\ H_\rho(1) = m}} H_\rho(y) \right\}.$$

Since $0 \leq H'_\rho \leq 1$, we have that $\sup_{\substack{\rho \in \mathcal{M}, \\ H_\rho(1) = m}} H_\rho(y)$ is equal to $y+1$ if $y+1 \leq m$ or m if $y+1 \geq m$. It follows that

$$\inf_{\substack{\rho \in \mathcal{M}, \\ H_\rho(1) = m}} \inf_{y \in [-1, 1]} \{y\xi_0 - H_\rho(y)\} = (m-1)\xi_0 - m.$$

Thus the existence of ρ_m is only possible if $\sup\{\xi_0, m - \xi_0\} \leq E(m) \leq m - (m-1)\xi_0$.

Let us denote ρ_m by ρ . Recall that y_ρ is the smallest point in $[-1, 1]$ such that the infimum in $\inf_{y \in [-1, 1]} \{y\xi_0 - H_\rho(y)\}$ is realized. If $y_\rho \in (-1, 1)$ then we have $y_\rho\xi_0 - H_\rho(y_\rho) = -E(m)$ which implies that

$$\begin{aligned} \frac{y_\rho\xi_0 + E(m)}{y_\rho + 1} &= \frac{H_\rho(y_\rho) - H_\rho(-1)}{y_\rho + 1} \in [0, 1], \\ \frac{m - (y_\rho\xi_0 + E(m))}{1 - y_\rho} &= \frac{H_\rho(1) - H_\rho(y_\rho)}{1 - y_\rho} \in [0, 1]. \end{aligned}$$

If $y_\rho = -1$, then we have $\xi_0 = E(m)$ and for any $z \geq -1$, $H_\rho(z) \leq z\xi_0 + E(m)$ and in particular, for $z = 1$, we get $m \leq 2\xi_0$. If $y_\rho = 1$, similarly, we have $\xi_0 + E(m) = m$ and $2\xi_0 \leq m$.

Thus, if a profile ρ is such that $\bar{V}^+(\rho) + \mathbb{S}(\rho) = E$ then (y_ρ, m_ρ) belongs to the set composed of couples $(y, m) \in [-1, 1] \times [0, 2]$ satisfying

$$\left\{ \begin{array}{l} \sup\{\xi_0, m - \xi_0\} \leq E(m) \leq m - (m-1)\xi_0, \\ \frac{y\xi_0 + E(m)}{y+1} \in [0, 1], \quad \frac{m - (y\xi_0 + E(m))}{1-y} \in [0, 1], \quad \text{if } y \in (-1, 1), \\ -\xi_0 + E(m) = 0, \quad m \leq 2\xi_0, \quad \text{if } y = -1, \\ \xi_0 + E(m) = m, \quad m \geq 2\xi_0, \quad \text{if } y = 1. \end{array} \right.$$

These conditions are equivalent to $(y, m) \in D$. By concavity of the function $-s$ together with Jensen's inequality, if $y_\rho \in [-1, 1]$, we have

$$\begin{aligned} \mathbb{S}(\rho) = \mathbb{S}(H'_\rho) &= -\frac{1}{2} \int_{-1}^{y_\rho} s(H'_\rho(x)) dx - \frac{1}{2} \int_{y_\rho}^1 s(H'_\rho(x)) dx \\ &\leq -\frac{(y_\rho + 1)}{2} s\left(\frac{y_\rho \xi_0 + E(m)}{y_\rho + 1}\right) - \frac{(1 - y_\rho)}{2} s\left(\frac{m - (y_\rho \xi_0 + E(m))}{1 - y_\rho}\right) \end{aligned} \quad (2.8)$$

with the convention that if $y_\rho = \pm 1$ the indefinite terms have to be replaced by 0. This proves that

$$S^+(E) \leq \frac{1}{2} \sup_{(y, m) \in D} F(y, m).$$

To prove the opposite inequality, consider any $(y, m) \in D$ and let H^y be the continuous function, linear on $[-1, y]$ and on $[y, 1]$ such that $H^y(-1) = 0, H^y(1) = m, H^y(y) = y\xi_0 + E(m)$. Since $(y, m) \in D$, the profile ρ such that $H_\rho = H^y$ belongs to \mathcal{M} and satisfies $H_\rho(1) = m, y_\rho = y$ and $\inf_z \{z\xi_0 - H_\rho(z)\} = -E(m)$, i.e. $\bar{V}^+(\rho) + \mathbb{S}(\rho) = E$. Observe now that the equality in (2.8) is satisfied for $H_\rho := H^{y\rho}$. This shows that $F(y, m) = 2\mathbb{S}(\rho) \leq 2S^+(E)$ and finishes the proof. \square

To prove Theorem 2.2 it remains to compute the supremum appearing in the statement of Proposition 2.3. This is done in Appendix A. The last part of Theorem 2.2 concerning the values of the maximizers is also postponed to Appendix A.

2.3. Pressure. We recall that the pressure function P^+ is defined as the Legendre transform of the entropy function S^+ :

$$P^+(\theta) = \inf_{E \geq 0} \{\theta E - S^+(E)\}.$$

We introduce the two parameters $\theta_0^\pm := -\frac{1}{\varphi_\pm} s'(\xi_0)$, where ξ_0 is defined in (2.3).

Theorem 2.4. *The pressure function P^+ is given by:*

- If $\rho_+ < 1 - \rho_- \leq \frac{1}{2}$ then

$$P^+(\theta) = \begin{cases} P(\varphi_-, -\theta) - \theta \bar{V}^+, & \theta \geq \theta_0^-, \\ P^+(\theta_0) + E_{+\infty}^+(\theta - \theta_0^-), & \theta < \theta_0^-. \end{cases}$$

- If $\frac{1}{2} \leq \rho_+ < 1 - \rho_-$ then

$$P^+(\theta) = \begin{cases} P(\varphi_+, -\theta) - \theta \bar{V}^+, & \theta \geq \theta_0^+, \\ P^+(\theta_0) + E_{+\infty}^+(\theta - \theta_0), & \theta < \theta_0^+. \end{cases}$$

- If $\rho_- < \rho_+ \leq \frac{1}{2}$ then

$$P^+(\theta) = \begin{cases} P(\varphi_-, -\theta) - \theta \bar{V}^+, & \theta \geq \theta_0^-, \\ P(\varphi_+, -\theta) - \theta \bar{V}^+, & \theta \leq \theta_0^+, \\ P^+(\theta_0^+) + \frac{P^+(\theta_0^-) - P^+(\theta_0^+)}{\theta_0^- - \theta_0^+}(\theta - \theta_0^+), & \theta \in (\theta_0^+, \theta_0^-). \end{cases}$$

- If $\frac{1}{2} \leq \rho_- < \rho_+$ then

$$P^+(\theta) = \begin{cases} P(\varphi_+, -\theta) - \theta \bar{V}^+, & \theta \geq \theta_0^+, \\ P(\varphi_-, -\theta) - \theta \bar{V}^+, & \theta \leq \theta_0^-, \\ P^+(\theta_0^-) + \frac{P^+(\theta_0^+) - P^+(\theta_0^-)}{\theta_0^+ - \theta_0^-}(\theta - \theta_0^-), & \theta \in (\theta_0^-, \theta_0^+), \end{cases}$$

where $P(\rho, \varphi)$ is given by (1.5).

The proof of this theorem is postponed to Appendix B.

2.4. Consequences. Let $\rho \in (0, 1)$ and φ be the associated chemical potential, see (1.3). Let us first observe that the equation of the tangent to the curve of $E \rightarrow S_{\rho, \rho}(E)$ at $E = E_0(\rho) := -\rho \log(1 - \rho) - (1 - \rho) \log(\rho)$ is given by

$$Y = -E - \log(\chi(\rho)).$$

This is the unique point where the tangent has a slope equal to -1 . Since $S_{\rho, \rho}$ is a concave function, the curve of $S_{\rho, \rho}$ is strictly below the tangent apart from the point $(E_0, S_{\rho, \rho}(E_0)) = (E_0, (-s)(\rho))$. Moreover, $E_0(\rho) \geq -\frac{\rho}{2} + \log(1 + e^\rho)$, i.e. E_0 is to the right of the point where the function $S_{\rho, \rho}$ has its maximum.

This permits to show that $J^+(E) = E - S^+(E)$ is a non negative convex function which vanishes for a unique value of E equal to $\mathbb{S}(\bar{\rho})$. We recall that J^+ is the large deviations function of the random variables $\{Y_N\}_N$ under $\mu_{ss, N}$.

From this we recover the result of Bahadoran ([1]) in the case of the TASEP. We also extend some of the results of [6] to the asymmetric simple exclusion process.

Theorem 2.5. *In the thermodynamic limit, the Gibbs-Shannon entropy of the non-equilibrium stationary state defined by*

$$\mathcal{S}(\mu_{ss, N}) = \sum_{\eta \in \Omega_N} [-\mu_{ss, N}(\eta) \log(\mu_{ss, N}(\eta))]$$

is equal to the Gibbs-Shannon entropy of the local equilibrium state, i.e.

$$\lim_{N \rightarrow +\infty} \frac{\mathcal{S}(\mu_{ss, N})}{2N + 1} = \mathbb{S}(\bar{\rho}).$$

Moreover, the corresponding fluctuations are Gaussian with a variance σ equal to the one provided by a local equilibrium statement, i.e. $\sigma = S''_{\bar{\rho}, \bar{\rho}}(\mathbb{S}(\bar{\rho})) = \frac{1}{2} \int_{-1}^1 \chi(\bar{\rho}) [(-s)'(\bar{\rho}(u))]^2 du$.

Proof. We just give the proof in the case $1/2 < 1 - \rho_- < \rho_+$ which corresponds to $0 < -\varphi_- < \varphi_+$ since the other cases are similar. Then, we have $\bar{\rho} = \rho_+$ and $\bar{V}^+ = \log(\rho_+(1 - \rho_+))$. Since, in the energy band, $J^+(E) = E + \bar{V}^+ - S_{\rho_+, \rho_+}(-E + \bar{V}^+) - \bar{V}^+$, we conclude that $J^+(E) \geq 0$ with equality if and only if $-(E + \bar{V}^+) = E_0(\rho_+)$ and $-E_0(\rho_+) - \bar{V}^+$ belongs to $[E_{+\infty}^-, E_{+\infty}^+]$. This last condition is equivalent to

$$-E_0(\rho_+) \in [-\log(1 + e^{\varphi_+}), W(\varphi_-, \varphi_+)]. \quad (2.9)$$

Since $-E_0(\rho_+) = -(1 - \rho_+) \log(1 + e^{-\varphi_+}) - \rho_+ \log(1 + e^{\varphi_+})$ we get easily that $-E_0(\rho_+) \geq -\log(1 + e^{\varphi_+})$. To prove the other inequality we write $W(\varphi_-, \varphi_+) + E_0(\rho_+) = g(\varphi_-)$, where the function $g : [-\varphi_+, 0] \rightarrow \mathbb{R}$ is defined by

$$g(x) = (1 - \rho_+) \log(1 + e^{-\varphi_+}) + \rho_+ \log(1 + e^{\varphi_+}) + \frac{x \log(1 + e^{\varphi_+}) - \varphi_+ \log(1 + e^x)}{\varphi_+ - x}.$$

Since

$$g'(x) = \frac{\varphi_+}{(\varphi_+ - x)^2} \left\{ \log(1 + e^{\varphi_+}) + \log(1 + e^x) - (\varphi_+ - x) \frac{e^x}{1 + e^x} \right\}$$

there exists $\tilde{\varphi}_x \in [x, \varphi_+]$ such that

$$g'(x) = \frac{\varphi_+}{(\varphi_+ - x)} \left\{ \frac{e^{\tilde{\varphi}_x}}{1 + e^{\tilde{\varphi}_x}} - \frac{e^x}{1 + e^x} \right\} \geq 0.$$

The last inequality follows from the fact that the function $t \rightarrow e^t/(1 + e^t)$ is increasing. Thus $g(\varphi_-) \leq g(-\varphi_+)$ and $g(-\varphi_+) = (\rho_+ - 1/2)\varphi_+ \geq 0$ which proves (2.9). Thus, J^+ is a non-negative convex function vanishing only for $-E_0(\rho_+) - \bar{V}^+ = (-s)(\rho_+) = \mathbb{S}(\bar{\rho})$. The function J^+ is smooth around $\mathbb{S}(\bar{\rho})$ and $J^+(E) = E + \bar{V}^+ - S_{\rho_+, \rho_+}(-E + \bar{V}^+) - \bar{V}^+$. By performing a second Taylor expansion of J^+ around $\mathbb{S}(\bar{\rho})$ we can determine the value of the variance of the Gaussian fluctuations and we get the desired result. \square

3. COOPERATIVE TASEP

In this section we compute (1.6) for the cooperative TASEP. Let \mathcal{F} be the set

$$\mathcal{F} := \{ \varphi \in C^1([-1, 1]) : \varphi(\pm 1) = \varphi_{\pm}, \varphi' > 0 \}.$$

and recall that $\chi : [0, 1] \rightarrow [0, 1]$ is defined by $\chi(\rho) = \rho(1 - \rho)$.

The quasi-potential V^- of the cooperative TASEP ([8], [2]) is defined by

$$V^-(\rho) = -\mathbb{S}(\rho) + \sup_{\varphi \in \mathcal{F}} \mathcal{H}(\rho, \varphi) - \bar{V}^- \quad (3.1)$$

where $\mathbb{S}(\cdot)$ is defined in (1.8), \mathcal{H} is defined in (2.2) and

$$\bar{V}^- = \log \left(\max_{\rho \in [\rho_-, \rho_+]} \chi(\rho) \right).$$

3.1. Energy bands. In this section we determine the energy band of the cooperative TASEP. This is summarized in Figure 4.

Proposition 3.1. *The bottom of the energy band is given by*

$$E_{-\infty}^- = -\bar{V}^- + \begin{cases} -\log(2), & \rho_- \leq \frac{1}{2} \leq \rho_+, \\ -\log(1 + e^{-\varphi_+}), & \rho_- < \rho_+ \leq \frac{1}{2}, \\ -\log(1 + e^{\varphi_-}), & \frac{1}{2} \leq \rho_- < \rho_+, \end{cases}$$

and the top of the energy band by is given by

$$E_{-\infty}^+ = -\bar{V}^- - \log(1 + e^{-\varphi_0})$$

where $\varphi_0 = \sup\{|\varphi_-|, |\varphi_+|\}$.

Proof. Let us first compute the top of the energy band. Recall from (1.7) that $E_{-\infty}^+ = \sup_{\rho \in \mathcal{M}} \{\mathbb{S}(\rho) + V^-(\rho)\}$. We have

$$\sup_{\rho \in \mathcal{M}} \sup_{\varphi \in \mathcal{F}} \mathcal{H}(\rho, \varphi) = \sup_{\varphi \in \mathcal{F}} \sup_{\rho \in \mathcal{M}} \mathcal{H}(\rho, \varphi).$$

For any $\varphi \in \mathcal{F}$ we define $x_\varphi = \sup\{x \in [-1, 1] : \varphi(x) \leq 0\}$. Then, $\sup_{\rho \in \mathcal{M}} \mathcal{H}(\rho, \varphi)$ is realized for $\rho(x) = \mathbf{1}\{-1 \leq x \leq x_\varphi\}$. It follows that

$$\sup_{\rho \in \mathcal{M}} \sup_{\varphi \in \mathcal{F}} \mathcal{H}(\rho, \varphi) = \sup_{x \in [-1, 1]} \sup_{\varphi \in \mathcal{F}_x} \left\{ \frac{1}{2} \int_x^1 (\varphi(u) - \log(1 + e^{\varphi(u)})) du - \frac{1}{2} \int_{-1}^x \log(1 + e^{\varphi(u)}) du \right\}$$

where \mathcal{F}_x is the set of functions $\varphi \in \mathcal{F}$ such that $\varphi(x) = 0$ if $x \in (-1, 1)$, $\varphi(x) \leq 0$ if $x = 1$ and $\varphi(x) \geq 0$ if $x = -1$. Assume for example that $\varphi_- \leq 0 \leq \varphi_+$ (the other cases are similar). By using the fact that the function $t \rightarrow t - \log(1 + e^t)$ is increasing, we see that

$$\sup_{\rho \in \mathcal{M}} \sup_{\varphi \in \mathcal{F}} \mathcal{H}(\rho, \varphi) = \sup_{x \in [-1, 1]} \left\{ -\frac{(x+1)}{2} \log(1 + e^{\varphi_-}) + \frac{(1-x)}{2} (\varphi_+ - \log(1 + e^{\varphi_+})) \right\}$$

because the supremum over \mathcal{F}_x is realized by a sequence of functions in \mathcal{F}_x converging to the step function $\varphi_- \mathbf{1}\{-1 \leq u \leq x\} + \varphi_+ \mathbf{1}\{x \leq u \leq 1\}$. The last supremum is equal to $-\log(1 + e^{-\varphi_0})$ which concludes the computation of the top of the energy band.

We compute now the bottom of the energy band. Recall from (1.7) that $E_{-\infty}^- = \inf_{\rho \in \mathcal{M}} \{\mathbb{S}(\rho) + V^-(\rho)\}$. We first recall some results of [8]. Recall the definition of H_ρ from (2.4) and let G_ρ be the convex envelop of H_ρ , i.e. the biggest convex function G such that $G \leq H_\rho$. We recall that any convex function is almost everywhere differentiable. Then, the supremum in (3.1) is given by $-\mathbb{S}(\rho) + \mathcal{H}(\rho, \varphi_{G_\rho}) + \bar{V}^-$ where

$$\varphi_{G_\rho}(x) = \begin{cases} \varphi_-, & G'_\rho(x) \leq \rho_-, \\ \log\left(\frac{G'_\rho(x)}{1 - G'_\rho(x)}\right), & \rho_- \leq G'_\rho(x) \leq \rho_+, \\ \varphi_+, & G'_\rho(x) \geq \rho_+. \end{cases}$$

Moreover, by (2.2) we have that

$$\mathcal{H}(\rho, \varphi_{G_\rho}) = \frac{1}{2} \int_{-1}^1 [G'_\rho(x) \varphi_{G_\rho}(x) - \log(1 + e^{\varphi_{G_\rho}(x)})] dx.$$

This shows that $\mathcal{H}(\rho, \varphi_{G_\rho})$ does not depend on ρ but only on G_ρ . Since as ρ describes the set \mathcal{M} , G_ρ describes exactly the set of non-decreasing convex functions G such that $G(-1) = 0$ and $0 \leq G' \leq 1$, then we have

$$\inf_{\rho \in \mathcal{M}} \sup_{\varphi \in \mathcal{F}} \mathcal{H}(\rho, \varphi) = \inf_{\rho \in \mathcal{M}} \mathcal{H}(\rho, \varphi_{G_\rho}) = \inf_G \left\{ \frac{1}{2} \int_{-1}^1 [G'(x) \varphi_G(x) - \log(1 + e^{\varphi_G(x)})] dx \right\}$$

where the last infimum is carried over the set of non-decreasing convex functions G such that $G(-1) = 0$ and $0 \leq G' \leq 1$.

Let us now consider the set of non-decreasing functions g such that $g \in [0, 1]$ and for such g let $T(g) : [-1, 1] \rightarrow [\varphi_-, \varphi_+]$ be defined by

$$T(g)(x) = \begin{cases} \varphi_-, & g(x) \leq \rho_-, \\ \log\left(\frac{g(x)}{1-g(x)}\right), & \rho_- \leq g(x) \leq \rho_+, \\ \varphi_+, & g(x) \geq \rho_+. \end{cases}$$

We have then

$$\inf_{\rho \in \mathcal{M}} \sup_{\varphi \in \mathcal{F}} \mathcal{H}(\rho, \varphi) = \inf_g \left\{ \frac{1}{2} \int_{-1}^1 [g(x)T(g)(x) - \log(1 + e^{T(g)(x)})] dx \right\} \quad (3.2)$$

where the infimum is taken over the set of non-decreasing functions $g : [-1, 1] \rightarrow [0, 1]$. To each non-decreasing function $g : [-1, 1] \rightarrow [0, 1]$ we associate $-1 \leq x_- \leq x_+ \leq 1$ and $0 \leq y_- \leq \rho_- < \rho_+ \leq y_+ \leq 1$ defined by

$$x_- = \sup\{x \in [-1, 1] : g(x) \leq \rho_-\}, \quad x_+ = \inf\{x \in [-1, 1] : g(x) \geq \rho_+\},$$

$$y_- = \frac{1}{x_- + 1} \int_{-1}^{x_-} g(x) dx, \quad y_+ = \frac{1}{1 - x_+} \int_{x_+}^1 g(x) dx.$$

In the case $x_- = -1$ (resp. $x_+ = 1$) we adopt the convention that $y_- = \rho_-$ (resp. $y_+ = \rho_+$). With these definitions we can write

$$\begin{aligned} & \int_{-1}^1 [g(x)T(g)(x) - \log(1 + e^{T(g)(x)})] dx \\ &= \left\{ (x_- + 1)(\varphi_- y_- - \log(1 + e^{\varphi_-})) + (1 - x_+)(\varphi_+ y_+ - \log(1 + e^{\varphi_+})) \right\} + \int_{x_-}^{x_+} s(g(x)) dx. \end{aligned}$$

The infimum can be computed by fixing first x_-, x_+, y_-, y_+ , optimizing separately over the restrictions of g to $[-1, x_-]$, $[x_-, x_+]$ and $[x_+, 1]$ and then taking the infimum over x_-, x_+, y_-, y_+ . These parameters shall satisfy $-1 \leq x_- \leq x_+ \leq 1$ and $0 \leq y_- \leq \rho_- < \rho_+ \leq y_+ \leq 1$.

By convexity of the function $s(\cdot)$ and by Jensen's inequality we get that the infimum of $\int_{x_-}^{x_+} s(g(x)) dx$ is given by $(x_+ - x_-) \inf_{\rho \in [\rho_-, \rho_+]} s(\rho)$. It follows that (3.2) is equal to

$$\begin{aligned} & \inf_{\substack{-1 \leq x_- \leq x_+ \leq 1 \\ 0 \leq y_- \leq \rho_- < \rho_+ \leq y_+ \leq 1}} \frac{1}{2} \left\{ (x_- + 1)(\varphi_- y_- - \log(1 + e^{\varphi_-})) + (1 - x_+)(\varphi_+ y_+ - \log(1 + e^{\varphi_+})) \right. \\ & \quad \left. + (x_+ - x_-) \left(\inf_{\rho \in [\rho_-, \rho_+]} s(\rho) \right) \right\} \\ &= \inf_{-1 \leq x_- \leq x_+ \leq 1} \frac{1}{2} \left\{ (x_- + 1)((\varphi_- \wedge 0)\rho_- - \log(1 + e^{\varphi_-})) + (1 - x_+)(\varphi_+ \wedge (\varphi_+ \rho_+)) \right. \\ & \quad \left. - \log(1 + e^{\varphi_+}) + (x_+ - x_-) \left(\inf_{\rho \in [\rho_-, \rho_+]} s(\rho) \right) \right\} \end{aligned}$$

We observe now that $\rho_{\pm} = e^{\varphi_{\pm}} / (1 + e^{\varphi_{\pm}})$ and that the function $t \rightarrow te^t / (1 + e^t) - \log(1 + e^t)$ is even, increasing on $[0, +\infty)$ and negative. The result then follows. \square




FIGURE 4. The phase diagram for the bottom of the energy band (resp. left) and the top of the energy band (resp. right) for the cooperative TASEP. We have $\underline{A} = -\log(1 - \rho_+)$, $\underline{B} = \log(2)$, $\underline{C} = -\log(\rho_-)$ and $\bar{A} = -\log(\chi(\rho_+)) + \log(1 - \rho_-)$, $\bar{B} = 2\log(2) + \log(1 - \rho_-)$, $\bar{C} = 2\log(2) + \log(\rho_+)$, $\bar{D} = -\log(\chi(\rho_-)) + \log(\rho_+)$. All transitions are of first order.

3.2. **Entropy.** Our aim is to prove that

Theorem 3.2. *The restriction of the entropy function S^- on the energy band $[E_{-\infty}^-; E_{-\infty}^+]$ is given by*

$$S^-(E) = \begin{cases} -(E + \bar{V}^-) \mathbf{1}\{(E + \bar{V}^-) \leq s(\rho_0)\} \\ + S_{\rho_0, \rho_0}(-E + \bar{V}^-) \mathbf{1}\{(E + \bar{V}^-) > s(\rho_0)\}, & \rho_- \leq \frac{1}{2} \leq \rho_+, \\ \\ S_{\rho_+, \rho_+}(-E + \bar{V}^-) \mathbf{1}\{(E + \bar{V}^-) < s(\rho_+)\} \\ -(E + \bar{V}^-) \mathbf{1}\{s(\rho_+) \leq (E + \bar{V}^-) \leq s(\rho_-)\} \\ + S_{\rho_-, \rho_-}(-E + \bar{V}^-) \mathbf{1}\{(E + \bar{V}^-) > s(\rho_-)\}, & \rho_- < \rho_+ \leq \frac{1}{2}, \\ \\ S_{\rho_-, \rho_-}(-E + \bar{V}^-) \mathbf{1}\{(E + \bar{V}^-) < s(\rho_-)\} \\ -(E + \bar{V}^-) \mathbf{1}\{s(\rho_-) \leq (E + \bar{V}^-) \leq s(\rho_+)\} \\ + S_{\rho_+, \rho_+}(-E + \bar{V}^-) \mathbf{1}\{(E + \bar{V}^-) > s(\rho_+)\}, & \frac{1}{2} \leq \rho_- < \rho_+. \end{cases}$$

Moreover, the function S^- is concave and continuously differentiable on its energy band. Its derivative is not continuous at the points where it becomes linear.

The supremum in the definition of $S^-(E)$, see (1.6), for $E \in [E_-^\infty; E_+^\infty]$ is realized by the profiles:

$$\left\{ \begin{array}{ll} \mathcal{M}_-^E, & \text{if } E + \bar{V}^- \leq s(\rho_0) \\ \{u_{\rho_0}\}, & \text{if } E + \bar{V}^- > s(\rho_0), \end{array} \right. \quad \rho_- \leq \frac{1}{2} \leq \rho_+,$$

$$\left\{ \begin{array}{ll} \{u_{\rho_+}\}, & \text{if } E + \bar{V}^- < s(\rho_+), \\ \mathcal{M}_-^E, & \text{if } s(\rho_+) \leq E + \bar{V}^- \leq s(\rho_-), \\ \{u_{\rho_-}\}, & \text{if } E + \bar{V}^- > s(\rho_-), \end{array} \right. \quad \rho_- < \rho_+ \leq \frac{1}{2},$$

$$\left\{ \begin{array}{ll} \{u_{\rho_-}\}, & \text{if } E + \bar{V}^+ < s(\rho_-), \\ \mathcal{M}_-^E, & \text{if } s(\rho_-) \leq E + \bar{V}^- \leq s(\rho_+), \\ \{u_{\rho_+}\}, & \text{if } E + \bar{V}^- > s(\rho_+), \end{array} \right. \quad \frac{1}{2} \leq \rho_- < \rho_+,$$

where for any ρ , the profile u_ρ is the constant profile equal to $\frac{\log(\rho) - (E + \bar{V}^-)}{\log(\rho) - \log(1-\rho)}$ and \mathcal{M}_-^E is the set of non-increasing profiles $\rho : [-1, 1] \rightarrow [1 - \rho_+, 1 - \rho_-]$ such that $\mathbb{S}(\rho) = -(E + \bar{V}^-)$.




FIGURE 5. Graph of the function S^- and graphs of the functions $E \rightarrow S_{\rho_\pm, \rho_\pm}(-E + \bar{V}^-)$ for $\rho_- = 0.2$, $\rho_+ = 0.6$ (resp. left) and for $\rho_- = 0.25$, $\rho_+ = 0.4$ (resp. right).

Once the form of S^- is known, the fact that it is concave and continuously differentiable on its energy band is trivial. To obtain the form of S^- we start to reduce the computation of S^- to a 4 dimensional optimization problem. Some notations shall be introduced. Let $m = \min_{\rho \in [\rho_-, \rho_+]}(-s(\rho))$ and $M = \max_{\rho \in [\rho_-, \rho_+]}(-s(\rho))$. We define the linear functions $\gamma_\pm : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\gamma_\pm(y) = y\varphi_\pm - \log(1 + e^{\varphi_\pm}), \quad y \in \mathbb{R}.$$

For any $E \geq 0$, let $K := K(E)$ be the (possibly empty) compact convex domain of \mathbb{R}^4 composed of 4-tuples (x_-, x_+, y_-, y_+) such that the following conditions are satisfied

$$\begin{cases} 0 \leq y_- \leq \rho_- < \rho_+ \leq y_+ \leq 1, \\ -1 \leq x_- \leq x_+ \leq 1, \\ \frac{1}{2} \left((x_- + 1)(\gamma_-(y_-) + m) + (1 - x_+)(\gamma_+(y_+) + m) \right) \geq (E + \bar{V}^-) + m, \\ \frac{1}{2} \left((x_- + 1)(\gamma_-(y_-) + M) + (1 - x_+)(\gamma_+(y_+) + M) \right) \leq (E + \bar{V}^-) + M. \end{cases}$$

Let $F : K \rightarrow \mathbb{R}$ be the function defined by

$$F(x_-, x_+, y_-, y_+) = \frac{1}{2} \left((x_- + 1)((-s)(y_-) + \gamma_-(y_-)) + (1 - x_+)((-s)(y_+) + \gamma_+(y_+)) \right).$$

Proposition 3.3. *For any $E \geq 0$, we have that*

$$S^-(E) = \sup_{k \in K} F(k) - (E + \bar{V}^-).$$

Moreover $K \neq \emptyset$ if and only if $E \in [E_{-\infty}^-; E_{-\infty}^+]$.

Proof. The last part of the proposition follows directly from the computations performed during the determination of the energy band. We assume now that $E \in [E_{-\infty}^-; E_{-\infty}^+]$.

We use the notations introduced in the proof of Proposition 3.1. Then we have

$$S^-(E) = \sup_{\rho \in \mathcal{M}} \{ \mathbb{S}(\rho) ; \mathcal{H}(\rho, \varphi_{G_\rho}) = E + \bar{V}^- \}.$$

Since, by convexity of the function s , we have $\mathbb{S}(\rho) \leq \mathbb{S}(G'_\rho)$, we get

$$S^-(E) \leq \sup_G \left\{ \mathbb{S}(G') ; \frac{1}{2} \int_{-1}^1 G'(x) \varphi_G(x) - \log(1 + e^{\varphi_G(x)}) dx = E + \bar{V}^- \right\}$$

where the supremum is carried over the set of non-decreasing convex functions G such that $G(-1) = 0$, $0 \leq G' \leq 1$. On the other hand, given any non-decreasing convex function G such that $G(-1) = 0$ and $0 \leq G' \leq 1$, let $\rho = 1 - G' \in \mathcal{M}$. We have $H_\rho = G = G_\rho$ and $\mathbb{S}(\rho) = \mathbb{S}(H'_\rho) = \mathbb{S}(G)$. It follows that

$$S^-(E) = \sup_G \left\{ \mathbb{S}(G') ; \frac{1}{2} \int_{-1}^1 G'(x) \varphi_G(x) - \log(1 + e^{\varphi_G(x)}) dx = E + \bar{V}^- \right\}.$$

This can be written as

$$S^-(E) = \sup_g \left\{ \mathbb{S}(g) : \frac{1}{2} \int_{-1}^1 g(x) T(g)(x) - \log(1 + e^{T(g)(x)}) dx = E + \bar{V}^- \right\} \quad (3.3)$$

where the supremum is taken over the set of non-decreasing functions g such that $g \in [0, 1]$. Then, the constraint in (3.3) is given by

$$\begin{aligned} & \frac{1}{2} \int_{-1}^1 g(x) T(g)(x) - \log(1 + e^{T(g)(x)}) dx \\ &= \frac{1}{2} \left((x_- + 1)\gamma_-(y_-) + (1 - x_+)\gamma_+(y_+) + \int_{x_-}^{x_+} s(g(x)) dx \right) \\ &= E + \bar{V}^-. \end{aligned} \quad (3.4)$$

Fix x_-, x_+, y_-, y_+ . We decompose the integral appearing in $\mathbb{S}(g)$ into the three integrals corresponding to the intervals $[-1, x_-]$, $[x_-, x_+]$ and $[x_+, 1]$ so that we can optimize independently over the restrictions of g to $[-1, x_-]$ and $[x_+, 1]$. Then the value of the integral of $(-s)(g)$ over $[x_-, x_+]$ is fixed by the constraint (3.4).

By using that $\sup_g \int_{-1}^{x_-} (-s)(g(x))dx$ over the constraint that $\int_{-1}^{x_-} g(x)dx = (x_- + 1)y_-$ is given by $(x_- + 1)(-s)(y_-)$ (and similarly for $\sup_g \int_{x_+}^1 (-s)(g(x))dx$), we conclude that $S^-(E)$ is given by

$$S^-(E) = \sup \left\{ F(x_-, x_+, y_-, y_+) - (E + \bar{V}^-) : (x_-, x_+, y_-, y_+) \in \mathcal{A} \right\}.$$

Above \mathcal{A} is the set of 4-tuples (x_-, x_+, y_-, y_+) such that

$$-1 \leq x_- \leq x_+ \leq 1, \quad 0 \leq y_- \leq \rho_- < \rho_+ \leq y_+ \leq 1$$

and that there exists a non-decreasing function $h : [x_-, x_+] \rightarrow [\rho_-, \rho_+]$ satisfying

$$\frac{1}{2} \int_{x_-}^{x_+} (-s)(h(x))dx = -(E + \bar{V}^-) + \frac{1}{2} \left((x_- + 1)\gamma_-(y_-) + (1 - x_+)\gamma_+(y_+) \right).$$

This last condition can be stated as

$$-(E + \bar{V}^-) + \frac{1}{2} \left((x_- + 1)\gamma_-(y_-) + (1 - x_+)\gamma_+(y_+) \right) \in \left[\frac{x_+ - x_-}{2} m; \frac{x_+ - x_-}{2} M \right]. \quad (3.5)$$

It is easy to see that $\mathcal{A} = K$ and we have proved the proposition. \square

Assume from now on that E belongs to the energy band $[E_{-\infty}^-; E_{-\infty}^+]$. We have to compute the supremum of the function F over the non-empty convex compact set K . To do this we first fix $y_- \in [0, \rho_-]$ and $y_+ \in [\rho_+, 1]$ and optimize over the couples (x_-, x_+) such that $(x_-, x_+, y_-, y_+) \in K$.

Observe now that writing

$$(-s)(y_{\pm}) + \gamma_{\pm}(y_{\pm}) = (-s)(y_{\pm}) - [(-s)(\rho_{\pm}) + (-s')(\rho_{\pm})(y_{\pm} - \rho_{\pm})]$$

and using the concavity of $-s$, we get that $F(x_-, x_+, y_-, y_+) \leq 0$ for all (x_-, x_+, y_-, y_+) ; and $F(x_-, x_+, y_-, y_+) = 0$ if and only if $(x_{\pm} \mp 1)(y_{\pm} - \rho_{\pm}) = 0$.

If $-(E + \bar{V}^-)$ belongs to $[m, M]$ then by taking $x_{\pm} = \pm 1$, we conclude that $\sup_{k \in K} F(k) = 0$ and consequently that

$$S^-(E) = -(E + \bar{V}^-). \quad (3.6)$$

Consider now the case where E belongs to the energy band but $-(E + \bar{V}^-) \notin [m, M]$. Fix first y_{\pm} . In order to keep notation simple and since y_{\pm} are fixed, we use the notation $\gamma_{\pm} := \gamma_{\pm}(y_{\pm})$. We have first to maximize the function $F(\cdot, \cdot, y_-, y_+)$ in the compact convex domain $D := D(y_-, y_+)$ composed of (x_-, x_+) such that

$$D = \left\{ \begin{array}{l} -1 \leq x_- \leq x_+ \leq 1, \\ \frac{1}{2} \left((x_- + 1)(\gamma_- + m) + (1 - x_+)(\gamma_+ + m) \right) \geq (E + \bar{V}^-) + m, \\ \frac{1}{2} \left((x_- + 1)(\gamma_- + M) + (1 - x_+)(\gamma_+ + M) \right) \leq (E + \bar{V}^-) + M. \end{array} \right. \quad (3.7)$$

The two last conditions are obtained from (3.5).

Since $F(\cdot, \cdot, y_-, y_+)$ is an affine function, the supremum of $F(\cdot, \cdot, y_-, y_+)$ is attained at one of the extremal points of the domain D . Consider the lines D_m and D_M defined by

$$D_m = \left\{ (x_-, x_+) \in \mathbb{R}^2; \frac{1}{2} \left((x_- + 1)(\gamma_- + m) + (1 - x_+)(\gamma_+ + m) \right) = (E + \bar{V}^-) + m \right\},$$

$$D_M = \left\{ (x_-, x_+) \in \mathbb{R}^2; \frac{1}{2} \left((x_- + 1)(\gamma_- + M) + (1 - x_+)(\gamma_+ + M) \right) = (E + \bar{V}^-) + M \right\}.$$

There are 3, 4 or 5 of such extremal points. The line D_m intersects the lines $x_- = x_+$, $x_- = -1$ and $x_+ = 1$ at the points

$$X_0 := \left(\frac{2(E + \bar{V}^-) - (\gamma_- + \gamma_+)}{\gamma_- - \gamma_+}, \frac{2(E + \bar{V}^-) - (\gamma_- + \gamma_+)}{\gamma_- - \gamma_+} \right),$$

$$X_m := \left(-1, 1 - \frac{2(E + \bar{V}^-) + 2m}{\gamma_+ + m} \right)$$

$$Y_m := \left(\frac{2(E + \bar{V}^-) + 2m}{\gamma_- + m} - 1, 1 \right),$$

respectively. The line D_M intersects the same lines at the points X_0 ,

$$X_M := \left(-1, 1 - \frac{2(E + \bar{V}^-) + 2M}{\gamma_+ + M} \right)$$

$$Y_M := \left(\frac{2(E + \bar{V}^-) + 2M}{\gamma_- + M} - 1, 1 \right),$$

respectively. Observe that the point $(x_-, x_+) = (-1, 1)$ does not belong to the domain D because $-(E + \bar{V}^-) \notin [m, M]$.

The rest of the proof consists in determining what are the extremal points of D according to the position of E in the energy band, find what is the supremum of $F(\cdot, \cdot, y_-, y_+)$ among these extremal points, and then to maximize over y_-, y_+ . This is accomplished in Appendix C. The proof of the last statement of the theorem is also postponed to the Appendix.

3.3. Pressure. Let us define the function $m : \mathbb{R} \rightarrow (0, +\infty)$ by

$$m(\theta) = \inf\{f_\theta(t) : t \in [\varphi_-, \varphi_+]\}, \quad \theta \in \mathbb{R},$$

where for any $\theta \in \mathbb{R}$, $f_\theta : \mathbb{R} \rightarrow (0, +\infty)$ is given by

$$f_\theta(t) = \begin{cases} \exp\left\{\frac{1}{\theta}\log(1 + e^{\theta t}) + \log(1 + e^{-t})\right\}, & \theta \neq 0, \\ \exp\{t/2 + \log(1 + e^{-t})\}, & \theta = 0. \end{cases}$$

It is easy to check that f_θ is increasing (resp. decreasing) on $(-\infty, 0)$, decreasing (resp. increasing) on $(0, +\infty)$ if $\theta < -1$ (resp. $\theta > -1$) and is constant equal to 1 if $\theta = -1$. It follows that

$$m(\theta) = \begin{cases} \min(f_\theta(\varphi_-), f_\theta(\varphi_+)), & 0 \notin [\varphi_-, \varphi_+] \\ \min(f_\theta(\varphi_-), f_\theta(\varphi_+), f_\theta(0)), & 0 \in [\varphi_-; \varphi_+] \end{cases}.$$

Theorem 3.4. *The pressure function P^- is given by*

$$P^-(\theta) = -\theta(\log(m(\theta)) + \bar{V}^-).$$

The proof of this theorem is postponed to Appendix D.

3.4. Consequences.

Theorem 3.5. *In the thermodynamic limit, the Gibbs-Shannon entropy of the non-equilibrium stationary state defined by*

$$\mathcal{S}(\mu_{ss,N}) = \sum_{\eta \in \Omega_N} [-\mu_{ss,N}(\eta) \log(\mu_{ss,N}(\eta))]$$

is equal to the Gibbs-Shannon entropy of the local equilibrium state, i.e.

$$\lim_{N \rightarrow +\infty} \frac{\mathcal{S}(\mu_{ss,N})}{2N+1} = \mathbb{S}(\bar{\rho}).$$

Moreover, in the case $\rho_+ > \bar{\rho} \geq 1/2$ or $\bar{\rho} < \rho_+ \leq 1/2$, the fluctuations are Gaussian with a variance $\sigma = \chi(\bar{\rho})[(-s)'(\bar{\rho})]^2$. In the case $\bar{\rho} \leq 1/2 \leq \rho_+$ the fluctuations are not Gaussian.

Proof. We start by giving the proof in the case $1/2 \leq 1 - \rho_- < \rho_+$, which corresponds to $0 \leq -\varphi_- < \varphi_+$. The case $1/2 \leq \rho_+ < 1 - \rho_-$ is similar and for that reason is omitted.

By the definition of \bar{V}^- , we have that $\bar{V}^- = \log(\chi(1/2)) = -2\log(2)$. Also, by the results of the previous sections, defining $J^-(E) = E - S^-(E)$, we have for $E + \bar{V}^- \in I_1 = [-\log(2); -s(\rho_+)]$, that $J^-(E) = E + (E + \bar{V}^-) = 2E + \bar{V}^-$. Since J^- is linear and increasing, we conclude that for $E + \bar{V}^- \in I_1$ it holds that $J^-(E) \geq 0$ with equality if and only if $E_0 = -\frac{1}{2}\bar{V}^- = \log(2)$, which satisfies $E_0 + \bar{V}^- = -\log(2) \in I_1$. Now, for $E + \bar{V}^- \in I_2 = (s(\rho_+); -\log(1 + e^{-\varphi_+})]$, $J^-(E) = E - S_{\rho_+, \rho_+}(-(E + \bar{V}^-))$. As in the previous chapter we conclude that $J^-(E) \geq 0$ with equality if and only if $E_0 := E_0(\rho_+) = -\rho_+ \log(1 - \rho_+) - (1 - \rho_+) \log(\rho_+)$ and $-E_0 \in I_2$. Now, we notice that by a simple computation $-E_0$ can be written as $-\log(1 + e^{\varphi_+}) + \frac{\varphi_+}{1 + e^{\varphi_+}}$. Since $s(\rho_+) = -\log(1 + e^{\varphi_+}) + \frac{\varphi_+}{1 + e^{\varphi_+}} = -\log(1 + e^{\varphi_+}) + \frac{\varphi_+ e^{\varphi_+}}{1 + e^{\varphi_+}}$ and since $\varphi_+ > 0$ we easily conclude that $-E_0 \leq s(\rho_+)$ and as a consequence $-E_0 \notin I_2$. Then J^- vanishes for a unique value $E_0 := \log(2) = \mathbb{S}(\bar{\rho})$ for $\bar{\rho} = 1/2$. Thus, the function J^- is linear around $\log(2)$ and the fluctuations are not Gaussian.

Now we consider the case $\rho_+ \leq 1/2$, which corresponds to $\varphi_- \leq \varphi_+ < 0$. By the definition of \bar{V}^- , we have that $\bar{V}^- = \log(\chi(\rho_+))$. By the previous results, for $E + \bar{V}^- \in I_1 = [s(\rho_+); s(\rho_-)]$, we have that $J^-(E) = E + (E + \bar{V}^-) = 2E + \bar{V}^-$. We conclude that for $E + \bar{V}^- \in I_1$ it holds that $J^-(E) \geq 0$ with equality if and only if $E_0 = -\frac{1}{2}\bar{V}^- = \frac{1}{2}\log(\chi(\rho_+))$. But in this case a simple computation shows that $E_0 \notin I_1$. On the other hand for $E + \bar{V}^- \in I_2 = [-\log(1 + e^{-\varphi_+}); s(\rho_+)]$ we have that $J^-(E) = E - S_{\rho_+, \rho_+}(-(E + \bar{V}^-))$. As above we conclude that for $E + \bar{V}^- \in I_2$ it holds that $J^-(E) \geq 0$ with equality if and only if $E_0 := E_0(\rho_+) = -\rho_+ \log(1 - \rho_+) - (1 - \rho_+) \log(\rho_+)$. Repeating the same computations as above, one shows that $-E_0 < s(\rho_+)$ so that $-E_0 \in I_2$. In the remaining case, namely $E + \bar{V}^- \in I_3 = (s(\rho_-); -\log(1 + e^{\varphi_-})]$ we have $J^-(E) = E - S_{\rho_-, \rho_-}(-(E + \bar{V}^-))$ and $J^-(E_0) = 0$ for $E_0 := E_0(\rho_-) = -\rho_- \log(1 - \rho_-) - (1 - \rho_-) \log(\rho_-)$ but in this case $E_0 \notin I_3$. Thus, J^- vanishes for a unique value of $-E_0 - \bar{V}^- := -\rho_+ \log(\rho_+) - (1 - \rho_+) \log(1 - \rho_+) = \mathbb{S}(\bar{\rho})$ for $\bar{\rho} = \rho_+$.

The case $0 \leq \varphi_- < \varphi_+$ is analogous to the previous one and for that reason we omitted its proof. \square

4. A LOCAL EQUILIBRIUM STATEMENT

In this section we give a derivation of the strong form of local equilibrium that we need in order to establish the variational formula (1.6). For any $\varepsilon > 0$, we split the set $\{-N, \dots, N\}$ into $K = \varepsilon^{-1}$ boxes of size εN (we assume εN to be an integer to simplify). To each configuration $\eta \in \Omega_N$, let $\mathbf{M}(\eta) = (M_1(\eta), \dots, M_K(\eta))$ with $M_i(\eta)$ be the number of particles in the i^{th} box in the configuration η . For $\mathbf{M} = (M_1, \dots, M_K)$ fixed, we denote by $\Omega_N(\mathbf{M})$ the configurations η such that for any $i \in \{1, \dots, K\}$, the number of particles in the i^{th} box is M_i and by $Z_N(\mathbf{M})$ its cardinal.

The strong form of local equilibrium is the following statement:

$$\lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\mathbf{M}} \sup_{\eta \in \Omega_N(\mathbf{M})} \left| \log (Z_N(\mathbf{M}) \mu_{ss,N}(\eta | \mathbf{M})) \right| = 0. \quad (\mathbf{H})$$

The stationary state $\mu_{ss,N}$ can be expressed in terms of a product of (infinite) matrices ([4]). We consider the TASEP with $p = 1$ but we do not assume in this section that $\rho_- < \rho_+$. Thus, the case $\rho_- < \rho_+$ corresponds to the competitive TASEP and the case $\rho_- > \rho_+$ to the cooperative TASEP (up to a trivial left-right symmetry). Moreover, to have notations consistent with [4] we consider the boundary driven TASEP on the lattice $\{1, \dots, N\}$ rather than on $\{-N, \dots, N\}$. Let $\Sigma_N = \{0, 1\}^{\{1, \dots, N\}}$.

By [4], there exist matrices D, E and vectors $|V\rangle, \langle W|$ such that $\langle W | V \rangle = 1$,

$$\begin{aligned} DE &= D + E, \\ (1 - \rho_+)D|V\rangle &= |V\rangle, \\ \langle W| \rho_- E &= \langle W| \end{aligned}$$

and

$$\mu_{ss,N}(\eta) = \frac{\omega_N(\eta)}{\langle W | (D + E)^N | V \rangle},$$

where the weight $\omega_N(\eta)$ is given by

$$\omega_N(\eta) = \langle W | \prod_{x=1}^N \{ \eta(x)D + [1 - \eta(x)]E \} | V \rangle.$$

Lemma 4.1. *For any $N \geq 2$, any $\eta \in \Sigma_N$ such there exists a site $x \in \{1, \dots, N-1\}$ for which $\eta_x = 1, \eta_{x+1} = 0$, we have that*

$$s_N(x, \eta) = \omega_N(\eta) - \omega_N(\sigma^{x,x+1} \eta)$$

has the same sign as $\rho_- - \rho_+$.

Proof. Let us define $s = \frac{1}{\rho_-} + \frac{1}{1-\rho_+} - \frac{1}{(1-\rho_+)\rho_-}$ which has the same sign as $\rho_- - \rho_+$. We prove the lemma by induction. A configuration of Σ_N is identified with a sequence of 0's and 1's of length N . For example 011 is the configuration $\eta \in \Sigma_3$ such that $\eta_1 = 0, \eta_2 = 1, \eta_3 = 1$. For $N = 2$, the induction hypothesis is trivial since

$$\omega_2(10) - \omega_2(01) = \langle W | DE - ED | V \rangle = s.$$

Assume that the induction hypothesis is valid for $N-1$. Consider a configuration $\eta \in \Sigma_N$ such that $\eta_x = 1, \eta_{x+1} = 0, x \in \{1, \dots, N-1\}$. We write η in the form $\eta = \alpha 10\beta$ where

the 1 is at position x . If $\alpha = \alpha'1$ then by using the relation $DE = D + E$, we have

$$\begin{aligned} s_N(x, \eta) &= \omega_N(\alpha'110\beta) - \omega_N(\alpha'101\beta) \\ &= \omega_{N-1}(\alpha'11\beta) + \omega_{N-1}(\alpha'10\beta) - \omega_{N-1}(\alpha'11\beta) - \omega_{N-1}(\alpha'01\beta) \\ &= s_{N-1}(x-1, \eta') \end{aligned}$$

where $\eta' = \alpha'10\beta$. Thus, by the induction hypothesis applied to η' , $s_N(x, \eta)$ has the same sign as s . If $\beta = 0\beta'$, the same conclusion holds. Thus we can assume that η is in the form $\alpha 0101\beta$. If $\beta = 0\beta'$, by using the relation $DE = D + E$ applied at position $(x+2, x+3)$, we get

$$\begin{aligned} s_N(x, \eta) &= \omega_{N-1}(\alpha 0101\beta') - \omega_{N-1}(\alpha 0011\beta') + \omega_{N-1}(\alpha 0100\beta') - \omega_{N-1}(\alpha 0010\beta') \\ &= s_{N-1}(x, \eta^1) + s_{N-1}(x, \eta^0) \end{aligned}$$

where the $\eta^1 = \alpha 0101\beta'$ and $\eta^0 = \alpha 0100\beta'$. By the induction hypothesis, this has the same sign as s . The same conclusion holds if $\alpha = \alpha'1$. By iterating this procedure, one can prove that $s_N(x, \eta)$ has the same sign as s if there is a 1 to the left of $x-1$ or a 0 to the right of $x+2$. The only remaining case is if η is in the form $\eta = 0 \dots 0101 \dots 1$ with m zeroes to the left of the leftmost one and n ones to the right of the rightmost zero. But in this case we have

$$\begin{aligned} s_N(x, \eta) &= \langle W | E^m (DE - ED) D^n | V \rangle = \frac{1}{\rho_-^m (1 - \rho_+)^n} \langle W | (DE - ED) | V \rangle \\ &= \frac{s}{\rho_-^m (1 - \rho_+)^n} \end{aligned}$$

which has the same sign as s and the lemma is proved. \square

This lemma is sufficient to prove the local equilibrium statement as in [3].

APPENDIX A. PROOF OF THEOREM 2.2

Recall the definition of D from (2.7). We can rewrite the set D in a more convenient form by introducing

$$a_m = \frac{E(m) - 1}{1 - \xi_0} \quad \text{and} \quad b_m = \frac{m - E(m)}{\xi_0}.$$

It is clear that if $E(m) \geq \sup\{\xi_0, m - \xi_0\}$ then $a_m \geq -1$ and $b_m \leq 1$. Also, if $E(m) \leq m + (1 - m)\xi_0$ then $a_m \leq b_m$. We have $a_m = -1$ (resp. $b_m = 1$) if and only if $E(m) = \xi_0$ (resp. $m - E(m) = \xi_0$).

We have

$$(y, m) \in D \Leftrightarrow \begin{cases} 0 \leq m \leq 2, & \sup\{\xi_0, m - \xi_0\} \leq E(m) \leq m - (m-1)\xi_0, \\ a_m \leq y \leq b_m. \end{cases}$$

It is easy to check that $D \neq \emptyset$ is equivalent to $\{m \in [0, 2]; \sup\{\xi_0, m - \xi_0\} \leq E(m) \leq m - (m-1)\xi_0\} \neq \emptyset$ which is equivalent to $E \in [E_{+\infty}^-, E_{+\infty}^+]$.

Assume from now on that $E \in [E_{+\infty}^-, E_{+\infty}^+]$.

We denote by $\alpha := \alpha(E)$ (resp. $\beta := \beta(E)$, resp. $\gamma := \gamma(E)$) the solution to the linear equation $E(m) = \xi_0$ (resp. $m - E(m) = \xi_0$, resp. $E(m) - m + (m-1)\xi_0 = 0$). We have that

$\{m \in [0, 2]; \sup\{\xi_0, m - \xi_0\} \leq E(m) \leq m - (m - 1)\xi_0\}$ is given by $m \in [0, 2]$ such that

$$\begin{cases} E(\alpha) \leq E(m) \\ m - E(m) \leq \beta - E(\beta) \\ E(m) - m + (m - 1)\xi_0 \leq E(\gamma) - \gamma + (\gamma - 1)\xi_0. \end{cases}$$

Since $E(m)$ is a linear function of m , $\{m \in [0, 2]; \sup\{\xi_0, m - \xi_0\} \leq E(m) \leq m - (m - 1)\xi_0\}$ is a closed interval $[m_-, m_+]$ (with $m_- := m_-(E)$ and $m_+ := m_+(E)$) of $[0, 2]$ and it is easy to show by inspection of the different cases that we have:

$\frac{1}{2} \leq \rho_- < \rho_+$	$\rho_- < \rho_+ \leq \frac{1}{2}$	$\frac{1}{2} \leq 1 - \rho_- \leq \rho_+$	$\frac{1}{2} \leq \rho_+ \leq 1 - \rho_-$
$m_- = \sup\{\alpha, \beta\}$	$m_- = \gamma$	$m_- = \alpha$	$m_- = \gamma$
$m_+ = \gamma$	$m_+ = \inf\{\alpha, \beta\}$	$m_+ = \gamma$	$m_+ = \beta$
$\inf\{\alpha, \beta\} \notin (m_-, m_+)$	$\sup\{\alpha, \beta\} \notin (m_-, m_+)$	$\beta \notin (m_-, m_+)$	$\alpha \notin (m_-, m_+)$

TABLE 1. Values of m_- and m_+ in terms of α, β and γ .

This shows that γ is always equal to m_- or m_+ , that α or β is the other boundary of the interval $[m_-, m_+]$ and that the remaining point among $\{\alpha, \beta, \gamma\}$ does not belong to (m_-, m_+) .

Let $f : [m_-, m_+] \rightarrow \mathbb{R}$ be defined by

$$f(m) = \sup_{y \in [a_m, b_m]} F(y, m) \quad (\text{A.1})$$

so that

$$S^+(E) = \frac{1}{2} \sup_{m \in [m_-, m_+]} f(m). \quad (\text{A.2})$$

Observe that $\alpha \neq \beta \neq \gamma$ apart from a finite (at most three) number of explicit values of E . If E is different from these values we say that E is a regular value of the energy. For simplicity we restrict the study to the case where E is a regular value but the same analysis could be performed for the non-regular values of E . Since $\alpha = \beta$ is equivalent to $\alpha = \beta = 2\xi_0$ we will always assume that it is not the case.

Lemma A.1. *Let $E \in [E_{+\infty}^-; E_{+\infty}^+]$ be a regular value of the energy.*

For any $m \in (m_-, m_+)$, we have that $f(m) = F(y(m), m)$ for a unique $y(m) \in (a_m, b_m)$ which is solution to the equation $\partial_y F(y(m), m) = 0$.

If α (resp. β) belongs to $\{m_-, m_+\}$ then $f(\alpha) = -2s(\alpha/2)$ (resp. $f(\beta) = -2s(\beta/2)$) and the supremum appearing in the definition of f is uniquely realized for $y = -1$ (resp. $y = 1$). We have $f(\gamma) = 0$ and the supremum in the definition of f is uniquely realized for $y = \gamma - 1$.

Proof. We notice that if $m = \gamma$ then $a_m = b_m = m - 1$ and $f(\gamma) = 0$. This shows the last sentence of the lemma.

Now, let $m \in (m_-, m_+)$. Then $a_m < b_m$. For any $y \in (a_m, b_m)$, we have

$$\begin{aligned}\partial_y F(y, m) &= -\frac{\xi_0 - E(m)}{y+1} s' \left(\frac{y\xi_0 + E(m)}{y+1} \right) - s \left(\frac{y\xi_0 + E(m)}{y+1} \right) \\ &\quad - \frac{m - E(m) - \xi_0}{1-y} s' \left(\frac{m - (y\xi_0 + E(m))}{1-y} \right) + s \left(\frac{m - (y\xi_0 + E(m))}{1-y} \right), \\ \partial_y^2 F(y, m) &= -\frac{(\xi_0 - E(m))^2}{(y+1)^3} s'' \left(\frac{y\xi_0 + E(m)}{y+1} \right) \\ &\quad - \frac{(m - E(m) - \xi_0)^2}{(1-y)^3} s'' \left(\frac{m - (y\xi_0 + E(m))}{1-y} \right).\end{aligned}$$

Since s is a strictly convex function, $\partial_y^2 F(y, m) < 0$, i.e. $F(\cdot, m)$ is strictly concave on (a_m, b_m) so that $\sup_{[a_m, b_m]} F(\cdot, m)$ is attained for a unique point $y(m)$ of $[a_m, b_m]$. If $m \neq \alpha$ then $a_m \neq -1$ and if $m \neq \beta$ then $b_m \neq 1$. Noticing that $(y\xi_0 + E(m))/(y+1)$ (resp. $(m - (y\xi_0 + E(m)))/(1-y)$) goes to 1 (resp. 0) as y goes to a_m (resp. b_m), we conclude that $\partial_y F(y, m)$ goes to $+\infty$ (resp. $-\infty$). This implies that $y(m) \in (a_m, b_m)$ and $(\partial_y F)(y(m), m) = 0$. If $m = \alpha \in [m_-, m_+]$, then

$$F(y, \alpha) = -(y+1)s(\xi_0) - (1-y)s \left(\frac{\alpha - \xi_0(y+1)}{1-y} \right)$$

for $y \in [-1, b_\alpha] \subset [-1, 1)$, $F(\cdot, \alpha)$ is strictly concave on $[-1, b_\alpha]$ and

$$\lim_{y \rightarrow -1} \partial_y F(y, \alpha) = (-s)(\xi_0) - (-s)(\alpha/2) - (\xi_0 - \alpha/2)(-s)'(\alpha/2) < 0$$

because s is strictly convex and $\alpha \neq 2\xi_0$ (otherwise E is not regular). The maximum of $F(\cdot, \alpha)$ is uniquely attained for $y(\alpha) := -1$ and $F(y(\alpha), \alpha) = -2s(\alpha/2)$.

Similarly, if $m = \beta \in [m_-, m_+]$, the supremum is uniquely attained for $y(\beta) = 1$ and $F(y(\beta), \beta) = -2s(\beta/2)$. □

Lemma A.2. *Let $E \in [E_{+\infty}^-; E_{+\infty}^+]$ be a regular value of the energy. The function $f : [m_-, m_+] \rightarrow [0, 2\log(2)]$ is a continuously differentiable function on (m_-, m_+) and, when $\alpha, \beta \in [m_-, m_+]$,*

$$f'(\alpha) = -s'(\alpha/2) \quad \text{and} \quad f'(\beta) = -s'(\beta/2).$$

Proof. It is clear that f is smooth on (m_-, m_+) and that the implicit function theorem applies. Thus the continuity and differentiability problems are only around the points m_- and m_+ .

Let us prove that $f(m)$ goes to $f(\alpha)$ and that $f'(m)$ has a limit as m goes to α (assuming that $\alpha \in \{m_-, m_+\}$) equal to $-s'(\alpha/2)$. The other case can be treated similarly. We will show that

$$\lim_{m \rightarrow \alpha} y(m) = -1, \quad \lim_{m \rightarrow \alpha} \frac{y(m)\xi_0 + E(m)}{y(m)+1} = \lim_{m \rightarrow +\infty} \frac{m - (y(m)\xi_0 + E(m))}{1 - y(m)} = \alpha/2. \quad (\text{A.3})$$

By the implicit function theorem, for any $m \neq \alpha, \beta$ we have

$$f'(m) = \partial_m F(y(m), m) \\ = -\frac{\varphi_+}{\varphi_+ - \varphi_-} s' \left(\frac{y(m) \xi_0 + E(m)}{y(m) + 1} \right) + \frac{\varphi_-}{\varphi_+ - \varphi_-} s' \left(\frac{m - (y(m) \xi_0 + E(m))}{1 - y(m)} \right).$$

From (A.3) we deduce that $\lim_{m \rightarrow \alpha} f(m) = f(\alpha)$ and $\lim_{m \rightarrow \alpha} f'(m) = -s'(\alpha/2)$.

Let $(m_n)_{n \geq 0}$ be a sequence in (m_-, m_+) ($m_n \neq \alpha, \beta, \gamma$ for any n) converging to α . Since for any m ,

$$y(m) \in [-1, 1], \quad \frac{y(m) \xi_0 + E(m)}{y(m) + 1} \in [0, 1], \quad \frac{m - (y(m) \xi_0 + E(m))}{1 - y(m)} \in [0, 1],$$

up to a subsequence we can assume that $y_n := y(m_n)$ converges to some $a \in [-1, 1]$ and that $\frac{y_n \xi_0 + E(m_n)}{y_n + 1}$ converges to $u \in [0, 1]$ and $\frac{m_n - (y_n \xi_0 + E(m_n))}{1 - y_n}$ converges to $v \in [0, 1]$.

If $a \in (-1, 1)$, then $u = \xi_0 \in (0, 1)$, $v = \xi_0 + \frac{\alpha - 2\xi_0}{1 - a}$ and $v \neq u$ since $\alpha \neq 2\xi_0$. By continuity of the functions involved and taking into account that $\partial_y F(y_n, m_n) = 0$, we get that

$$-s(u) - \frac{\alpha - 2\xi_0}{1 - a} s'(v) + s(v) = 0.$$

The term on the left hand side of the previous equality can be written as $s(v) - s(u) - (v - u)s'(v)$ which is negative, by the convexity of the function s (if $v = 0$ or $v = 1$ then $s'(v) = -\infty$ or $s'(v) = +\infty$ and the inequality is still valid). Therefore $a \in \{-1, 1\}$.

If $a = 1$, then since $\frac{m_n - (y_n \xi_0 + E(m_n))}{1 - y_n}$ converges to $v \in [0, 1]$ and $y_n \rightarrow a = 1$, $m_n \rightarrow \alpha$ it implies that $\alpha = 2\xi_0$ that is in contradiction with our assumptions.

It follows that $a = -1$, so that $v = \alpha/2$. Observe that $\frac{E(m_n) - \xi_0}{y_n + 1}$ converges to $u - \xi_0$. Using the fact that $\partial_y F(y_n, m_n) = 0$, we get

$$0 = (u - \xi_0)s'(u) - s(u) - \frac{\alpha - 2\xi_0}{2} s'(\alpha/2) + s(\alpha/2).$$

Since s is convex, the function $z \rightarrow (z - \xi_0)s'(z) - s(z)$ is monotone, so that the equality is uniquely satisfied for $u = \alpha/2$. This proves (A.3). \square

Lemma A.3. *Let $E \in [E_{+\infty}^-; E_{+\infty}^+]$ be a regular value of the energy. The function f is strictly concave on (m_-, m_+) .*

Proof. On \mathring{D} we have

$$\begin{aligned} \partial_m^2 F(y, m) &= -\left(\frac{\varphi_+}{\varphi_+ - \varphi_-} \right)^2 \frac{1}{y+1} s'' \left(\frac{\xi_0 - E(m)}{y+1} \right) \\ &\quad - \left(\frac{\varphi_-}{\varphi_+ - \varphi_-} \right)^2 \frac{1}{1-y} s'' \left(\frac{m - (y \xi_0 + E(m))}{1-y} \right) \\ \partial_{y,m}^2 F(y, m) &= -\left(\frac{\varphi_+}{\varphi_+ - \varphi_-} \right) \frac{\xi_0 - E(m)}{(y+1)^2} s'' \left(\frac{\xi_0 - E(m)}{y+1} \right) \\ &\quad + \left(\frac{\varphi_-}{\varphi_+ - \varphi_-} \right) \frac{(m - E(m) - \xi_0)}{(1-y)^2} s'' \left(\frac{m - (y \xi_0 + E(m))}{1-y} \right). \end{aligned}$$

One easily checks that

$$\begin{aligned} & (\partial_y^2 F)(\partial_m^2 F) - (\partial_{y,m}^2 F)^2 \\ &= \frac{s''(a)s''(b)}{(\varphi_+ - \varphi_-)^2(y^2 - 1)^2} \left[\varphi_+ \sqrt{\frac{1+y}{1-y}}(m - E(m) - \xi_0) + \varphi_- \sqrt{\frac{1-y}{1+y}}(\xi_0 - E(m)) \right]^2, \end{aligned}$$

where

$$a = \frac{y\xi_0 - E(m)}{y+1} \quad \text{and} \quad b = \frac{m - (y\xi_0 + E(m))}{1-y}.$$

By convexity of the function s it follows that $(\partial_y^2 F)(\partial_m^2 F) - (\partial_{y,m}^2 F)^2 > 0$ if $m \neq \alpha, \beta$.

By the implicit function theorem, the function $m \rightarrow F(y(m), m)$ is smooth on (m_-, m_+) and

$$\begin{aligned} f''(m) &= \partial_m^2 F(y(m), m) + \partial_{y,m}^2 F(y(m), m) y'(m) \\ &\frac{d}{dm} [\partial_y F(y(m), m)] = 0 = (\partial_{m,y}^2 F)(y(m), m) + (\partial_y^2 F)(y(m), m) y'(m), \end{aligned}$$

so that

$$f''(m) = \frac{[(\partial_y^2 F)(\partial_m^2 F) - (\partial_{y,m}^2 F)^2](y(m), m)}{\partial_y^2 F(y(m), m)}.$$

Recalling from the proof of lemma A.1 that $\partial_y^2 F < 0$, we get that $f''(m) < 0$. \square

We have to compute $\sup_{m \in [0,2]} f(m)$. Since f is strictly concave there exists a unique $m_0 \in [m_-, m_+]$ for which the supremum of f is attained.

The point m_0 belongs to (m_-, m_+) if and only if there exists $m \in (m_-, m_+)$ such that $f'(m) = 0$. This is equivalent to the existence of $m \in (m_-, m_+)$ and $y \in (a_m, b_m)$ such that

$$\partial_m F(y, m) = 0, \quad \partial_y F(y, m) = 0. \quad (\text{A.4})$$

To simplify notations we introduce

$$\tilde{a} = s' \left(\frac{y\xi_0 + E(m)}{y+1} \right) \quad \text{and} \quad \tilde{b} = s' \left(\frac{m - (y\xi_0 + E(m))}{1-y} \right),$$

so that $\tilde{a} := s'(a)$ and $\tilde{b} := s'(b)$, where a and b were introduced above. Then (A.4) is equivalent to

$$\tilde{b} = \frac{\varphi_+}{\varphi_-} \tilde{a} \quad \text{and} \quad \xi_0 \left(1 - \frac{\varphi_-}{\varphi_+} \right) \tilde{b} + \log \left(\frac{1 + e^{\frac{\varphi_-}{\varphi_+} \tilde{b}}}{1 + e^{\tilde{b}}} \right) = 0.$$

There are two solutions to the second equation, $\tilde{b} = 0$ and $\tilde{b} = \varphi_+$.

If $\tilde{b} = 0$ then from $a = b = 1/2$ we get that $m = 1$. As a consequence we obtain that $y = (1/2 - E(1))/(\xi_0 - 1/2)$. The condition $y \in (a_1, b_1)$ implies that $E(1) < \sup\{1 - \xi_0, \xi_0\}$, which is in contradiction with the fact that $m = 1$ shall satisfy $\sup\{\xi_0, m - \xi_0\} \leq E(m) \leq m - (m - 1)\xi_0$.

If $\tilde{b} = \varphi_+$ then

$$a = \frac{y\xi_0 + E(m)}{y+1} = \rho_- \quad \text{and} \quad b = \frac{m - (y\xi_0 + E(m))}{1-y} = \rho_+.$$

As a consequence we obtain that $y = \frac{m - \rho_+ - \rho_-}{\rho_- - \rho_+}$. Then we get that

$$E(m) - \xi_0 = (\rho_- - \xi_0)(y+1), \quad (m - E(m) - \xi_0) = (\rho_+ - \xi_0)(1-y).$$

Thus the conditions $\sup\{\xi_0, m - \xi_0\} \leq E(m)$ and $y \in (-1, 1)$ imply $\rho_- \geq \xi_0 \geq \rho_+$. But we assumed $\rho_- < \rho_+$ and we have a contradiction.

Therefore $m_0 \in \{m_-, m_+\}$. Consequently, for any regular value $E \in [E_{+\infty}^-; E_{+\infty}^+]$,

$$S^+(E) = \frac{1}{2} \sup\{f(m_-), f(m_+)\}.$$

Recall that the set $\{m_-, m_+\}$ is equal to $\{\alpha, \gamma\}$ or to $\{\beta, \gamma\}$ and that $f(\gamma) = 0$, $f(\alpha) = -2s(\alpha/2)$ and $f(\beta) = -2s(\beta/2)$. Thus, by using the results in Table 1, we have

$$S^+(E) = \begin{cases} -s(\sup\{\alpha, \beta\}/2), & 0 < \varphi_- < \varphi_+, \\ -s(\inf\{\alpha, \beta\}/2), & \varphi_- < \varphi_+ < 0, \\ -s(\alpha/2), & 0 < -\varphi_- \leq \varphi_+, \\ -s(\beta/2), & 0 < \varphi_+ \leq -\varphi_-. \end{cases}$$

By definition of α and β we have that

$$\frac{\alpha}{2} = \frac{1}{\varphi_+} \{(E + \bar{V}^+) + \log(1 + e^{\varphi_+})\} \quad \text{and} \quad \frac{\beta}{2} = \frac{1}{\varphi_-} \{(E + \bar{V}^+) + \log(1 + e^{\varphi_-})\}.$$

Recall from (2.6) the definition of $W(\varphi_-, \varphi_+)$ and let $\varphi = \log(\rho/(1-\rho))$. Observe that the function $x \rightarrow S_{\rho, \rho}(-x)$ is a concave function equal to $-\infty$ outside $[i_1, i_2] := [-\log(1 + e^{-|\varphi|}); -\log(1 + e^{|\varphi|})]$, positive inside, vanishing at the boundaries of the interval, attaining its maximum equal to $\log(2)$ for $x_0 := (i_1 + i_2)/2 = \varphi/2 - \log(1 + e^\varphi)$. It is increasing on $[i_1, x_0]$ and decreasing on $[x_0, i_2]$.

A.1. The case $\frac{1}{2} \leq 1 - \rho_- \leq \rho_+$:

Recall that the energy band is given by

$$-\bar{V}^+ + [-\log(1 + e^{\varphi_+}); W(\varphi_-, \varphi_+)].$$

The entropy function S^+ , in the energy band, is given by $S^+(E) = S_{\rho_+, \rho_+}(-(E + \bar{V}^+))$. Remark that we have $W(\varphi_-, \varphi_+) + \log(1 + e^{\varphi_-}) = \varphi_- \xi_0 < 0$ since $\xi_0 \in (0, 1)$. Thus, $W(\varphi_-, \varphi_+) < -\log(1 + e^{-\varphi_+})$ and the function S^+ is concave, smooth in the interior of the energy band, but does not vanish at the top of the energy band.

A.2. The case $\frac{1}{2} \leq \rho_+ \leq 1 - \rho_-$:

Recall that the energy band is given by

$$-\bar{V}^+ + [-\log(1 + e^{-\varphi_-}); W(\varphi_-, \varphi_+)].$$

The entropy function S^+ , in the energy band, is given by $S^+(E) = S_{\rho_-, \rho_-}(-(E + \bar{V}^+))$. The function S^+ is concave, smooth in the interior of the energy band, but does not vanish at the top of the energy band.

It remains now to prove the last statement of Theorem 2.2. Let us assume that ρ is a maximizer of $S^+(E)$, E belonging to the energy band. We use the notations of the proof of Proposition 2.3. In the proof of this proposition, we have seen that ρ being a maximizer of $S^+(E)$ is equivalent to the fact that $(y_\rho, H_\rho(1)) \in D(E)$ being a maximizer of the function F over $D(E)$ and ρ is such that H_ρ is linear on $[-1, y_\rho]$ and on $[y_\rho, 1]$ with $H_\rho(y_\rho) = y_\rho \xi_0 + E(m_\rho)$. Moreover, we have seen above that such a maximizer (y, m) satisfies $y = \pm 1$ and

$$\begin{aligned} \text{i) } m &= \sup(\{\alpha, \beta\}) = \begin{cases} \beta & \text{if } E + \bar{V}^+ > W(\rho_-, \rho_+), \\ \alpha & \text{if } E + \bar{V}^+ \leq W(\rho_-, \rho_+), \end{cases} \quad , \quad 1/2 \leq \rho_- < \rho_+, \\ \text{ii) } m &= \inf(\{\alpha, \beta\}) = \begin{cases} \beta & \text{if } E + \bar{V}^+ < W(\rho_-, \rho_+), \\ \alpha & \text{if } E + \bar{V}^+ \geq W(\rho_-, \rho_+), \end{cases} \quad , \quad \rho_- < \rho_+ \leq 1/2, \\ \text{iii) } m &= \alpha \quad 1/2 \leq 1 - \rho_- \leq \rho_+, \\ \text{iv) } m &= \beta \quad 1/2 \leq \rho_+ \leq 1 - \rho_-. \end{aligned}$$

This implies in particular that if ρ is a maximizer of $S^+(E)$ then H_ρ is linear with a slope equal to $m_\rho/2$, i.e. ρ is constant equal to $1 - m_\rho/2$. Since, by definition, we have

$$\alpha = 2 \frac{\log(1 + e^{\varphi_+}) + (E + \bar{V}^+)}{\varphi_+} \quad \text{and} \quad \beta = 2 \frac{\log(1 + e^{\varphi_-}) + (E + \bar{V}^+)}{\varphi_-},$$

we get the result.

A.3. The case $\rho_- < \rho_+ \leq \frac{1}{2}$:

Recall that the energy band is given by

$$-\bar{V}^+ + [-\log(1 + e^{-\varphi_-}); -\log(1 + e^{\varphi_+})].$$

The condition $\alpha/2 \leq \beta/2$ is equivalent to

$$E + \bar{V}^+ \geq W(\varphi_-, \varphi_+).$$

Observe that $W(\varphi_-, \varphi_+) + \log(1 + e^{\varphi_+}) = \varphi_+ \xi_0$ and $\xi_0 \in (0, 1)$, so that

$$-\log(1 + e^{-\varphi_-}) < -\log(1 + e^{-\varphi_+}) \leq W(\varphi_-, \varphi_+) \leq -\log(1 + e^{\varphi_+}) < -\log(1 + e^{\varphi_-}).$$

Moreover, we have $\xi_0 < 1/2$ because there exists $\tilde{\varphi} \in [\varphi_-, \varphi_+]$ such that $\xi_0 = \frac{e^{\tilde{\varphi}}}{1 + e^{\tilde{\varphi}}}$. This implies that $W(\varphi_-, \varphi_+) \geq \frac{\varphi_+}{2} - \log(1 + e^{\varphi_+})$. We recall that $\frac{\varphi_+}{2} - \log(1 + e^{\varphi_+})$ is the first coordinate of the point for which the concave function $x \rightarrow S_{\rho_+, \rho_+}(-x)$ attains its maximum given by $\log(2)$. It follows that S^+ is a concave function. On the energy band it is given by

$$\begin{aligned} S^+(E) &= S_{\rho_-, \rho_-}(-(E + \bar{V}^+)) \mathbf{1}\{(E + \bar{V}^+) \leq W(\varphi_-, \varphi_+)\} \\ &\quad + S_{\rho_+, \rho_+}(-(E + \bar{V}^+)) \mathbf{1}\{(E + \bar{V}^+) > W(\varphi_-, \varphi_+)\}. \end{aligned}$$

The function S^+ is not differentiable at the point $W(\varphi_-, \varphi_+) - \bar{V}^+$.

A.4. The case $\frac{1}{2} \leq \rho_- < \rho_+$:

Recall that the energy band is given by

$$-\bar{V}^+ + [-\log(1 + e^{\varphi_+}); -\log(1 + e^{-\varphi_-})].$$

The condition $\alpha/2 \leq \beta/2$ is equivalent to $E + \bar{V}^+ \geq W(\varphi_-, \varphi_+)$. Observe that $W(\varphi_-, \varphi_+) + \log(1 + e^{\varphi_-}) = \varphi_- \xi_0$ and $\xi_0 \in (0, 1)$, so that

$$-\log(1 + e^{\varphi_+}) \leq -\log(1 + e^{\varphi_-}) \leq W(\varphi_-, \varphi_+) \leq -\log(1 + e^{-\varphi_-}) \leq -\log(1 + e^{-\varphi_+}).$$

Moreover, we have $\xi_0 > 1/2$ because there exists $\tilde{\varphi} \in [\varphi_-, \varphi_+]$ such that $\xi_0 = \frac{e^{\tilde{\varphi}}}{1+e^{\tilde{\varphi}}}$. This implies that $W(\varphi_-, \varphi_+) \geq \frac{\varphi_-}{2} - \log(1 + e^{\varphi_-})$. We recall that $\frac{\varphi_-}{2} - \log(1 + e^{\varphi_-})$ is the first coordinate of the point for which the concave function $x \rightarrow S_{\rho_-, \rho_-}(-x)$ attains its maximum given by $\log(2)$. It follows that S^+ is a concave function. On the energy band it is given by

$$S^+(E) = S_{\rho_-, \rho_-}(-(E + \bar{V}^+)) \mathbf{1}\{(E + \bar{V}^+) \geq W(\varphi_-, \varphi_+)\} \\ + S_{\rho_+, \rho_+}(-(E + \bar{V}^+)) \mathbf{1}\{(E + \bar{V}^+) < W(\varphi_-, \varphi_+)\}.$$

The function S^+ is not differentiable at the point $W(\varphi_-, \varphi_+) - \bar{V}^+$.

APPENDIX B. PROOF OF THEOREM 2.4

In order to prove the theorem, we have simply to compute the Legendre transform of S^+ whose explicit form is given in Theorem 2.2. Recall also that the Legendre transform of the function $S_{\rho, \rho}$ defined by (1.4) is given by the function $P(\varphi, \cdot)$ defined by (1.5).

B.1. The case $\frac{1}{2} \leq 1 - \rho_- < \rho_+$:

For any $E \in (E_{+\infty}^-, E_{+\infty}^+)$ we have that $\frac{dS^+}{dE}$ is a decreasing function and

$$\lim_{E \rightarrow E_{+\infty}^-} \frac{dS^+}{dE} = +\infty, \quad \lim_{E \rightarrow E_{+\infty}^+} \frac{dS^+}{dE} = \theta_0^-.$$

We get that

$$P^+(\theta) = \begin{cases} P(\varphi_-, -\theta) - \theta \bar{V}^+, & \theta \geq \theta_0^-, \\ P^+(\theta_0^-) + E_{+\infty}^+(\theta - \theta_0^-), & \theta < \theta_0^-. \end{cases}$$

B.2. The case $\frac{1}{2} \leq \rho_+ < 1 - \rho_-$:

For any $E \in (E_{+\infty}^-, E_{+\infty}^+)$ we have that $\frac{dS^+}{dE}$ is a decreasing function and

$$\lim_{E \rightarrow E_{+\infty}^-} \frac{dS^+}{dE} = +\infty, \quad \lim_{E \rightarrow E_{+\infty}^+} \frac{dS^+}{dE} = \theta_0^+.$$

We get that

$$P^+(\theta) = \begin{cases} P(\varphi_+, -\theta) - \theta \bar{V}^+, & \theta \geq \theta_0^+, \\ P^+(\theta_0^+) + E_{+\infty}^+(\theta - \theta_0^+), & \theta < \theta_0^+. \end{cases}$$

B.3. The case $\rho_- < \rho_+ \leq \frac{1}{2}$:

In this case we have that $\theta_0^+ \leq \theta_0^-$ because $\xi_0 \in (0, 1/2)$. We get similarly that

$$P^+(\theta) = \begin{cases} P(\varphi_-, -\theta) - \theta \bar{V}^+, & \theta \geq \theta_0^-, \\ P(\varphi_+, -\theta) - \theta \bar{V}^+, & \theta \leq \theta_0^+, \\ P^+(\theta_0^+) + \frac{P^+(\theta_0^-) - P^+(\theta_0^+)}{\theta_0^- - \theta_0^+}(\theta - \theta_0^+), & \theta \in (\theta_0^+, \theta_0^-). \end{cases}$$

B.4. The case $\frac{1}{2} \leq \rho_- < \rho_+$:

The function S^+ is differentiable everywhere in the interior of the energy band apart from the point $W(\varphi_-, \varphi_+) - \bar{V}^+$. For $E = W(\varphi_-, \varphi_+) - \bar{V}^+$, S^+ has a left-tangent and a right-tangent. Moreover, $\frac{dS^+}{dE}$ is decreasing on $(E_{+\infty}^-; W(\varphi_-, \varphi_+) - \bar{V}^+)$ and increasing on $(W(\varphi_-, \varphi_+) - \bar{V}^+; E_{+\infty}^+)$. We have

$$\lim_{E \rightarrow E_{+\infty}^\pm} \frac{dS^+}{dE} = \mp\infty \quad \text{and} \quad \lim_{E \rightarrow [W(\varphi_-, \varphi_+) - \bar{V}^+]^\pm} \frac{dS^+}{dE} = \theta_0^\pm.$$

Observe that $\theta_0^- \leq \theta_0^+$ because $\xi_0 \in (1/2, 1)$. We get easily that

$$P^+(\theta) = \begin{cases} P(\varphi_+, -\theta) - \theta \bar{V}^+, & \theta \geq \theta_0^+, \\ P(\varphi_-, -\theta) - \theta \bar{V}^+, & \theta \leq \theta_0^-, \\ P^+(\theta_0^-) + \frac{P^+(\theta_0^+) - P^+(\theta_0^-)}{\theta_0^+ - \theta_0^-}(\theta - \theta_0^-), & \theta \in (\theta_0^-, \theta_0^+). \end{cases}$$

APPENDIX C. PROOF OF THEOREM 3.2

In this section we determine the extremal points of the domain Δ according to the position of E along the energy band, we find the supremum of $F(\cdot, \cdot, y_-, y_+)$ among those points and then we maximize over y_- and y_+ .

C.1. The case $\rho_- \leq \frac{1}{2} \leq \rho_+$:

This case corresponds to $\varphi_- \leq 0 \leq \varphi_+$, therefore $m = (-s)(\rho_0)$ and $M = (-s)(1/2) = \log(2) = -E_{-\infty}^- - \bar{V}^-$. Since $\gamma'_+(y_+) = \varphi_+ > 0$, then the function γ_+ is increasing. On the other hand, since $\gamma'_-(y_-) = \varphi_- < 0$, the function γ_- is decreasing. Since $\gamma_+(\rho_+) := \varphi_+/(1 + e^{-\varphi_+}) - \log(1 + e^{\varphi_+})$ and the function $t \rightarrow t/(1 + e^{-t}) - \log(1 + e^t)$ is increasing in $(0, +\infty)$, we obtain that $-M \leq \gamma_+(\rho_+) < \gamma_+(1)$. On the other hand, since $\gamma_-(\rho_-) = \varphi_-/(1 + e^{-\varphi_-}) - \log(1 + e^{\varphi_-})$ and the function $t \rightarrow t/(1 + e^{-t}) - \log(1 + e^t)$ is decreasing in $(-\infty, 0)$ we obtain that $-M \leq \gamma_-(\rho_-) \leq \gamma_-(0)$.

We only consider the case $\frac{1}{2} \leq 1 - \rho_- < \rho_+$ (which corresponds to $\varphi_0 = \varphi_+$), the case $\frac{1}{2} \leq \rho_+ < 1 - \rho_-$ (which corresponds to $\varphi_0 = -\varphi_-$) being similar.

Since, $-m = \gamma_+(\rho_+)$ we have that $-M < -m < \gamma_+(1) = E_{-\infty}^+ + \bar{V}^-$. As a consequence, $\gamma_+(y_+) \geq -m$ for all $y_+ \in [\rho_+, 1]$. Notice that the function $t \rightarrow t/(1 + e^{-t}) - \log(1 + e^t)$ is even and increasing in $(0, +\infty)$. Therefore, $\gamma_-(\rho_-) \leq \gamma_+(\rho_+)$. On the other hand, $\gamma_-(0) \leq \gamma_+(1)$, which implies that $-M \leq \gamma_-(\rho_-) \leq -m = \gamma_+(\rho_+) \leq \gamma_+(1)$. Now, two things can happen, either $\gamma_-(0) > -m$ or $\gamma_-(0) < -m$. We start by the former.

(a) $\gamma_-(0) > -m$: Since we do not know the sign of $\gamma_- - \gamma_+$ we split again into two cases: $\gamma_- < \gamma_+$ and $\gamma_- > \gamma_+$. We start by the former.

(a.1) Case $\gamma_- > \gamma_+$: Recall the intersection points of the lines D_m and D_M from Section 3.2. In this case X_0 is in the square $[-1, 1]^2$ if and only if $\gamma_+ \leq E + \bar{V}^- \leq \gamma_-$. We first restrict to the case $X_0 \in [-1, 1]^2$, i.e. $\gamma_+ \leq E + \bar{V}^- \leq \gamma_-$. Now, we check whether X_m, X_M, Y_m and Y_M are in the domain D . We start with X_m , and the same computations holds for X_M . For that purpose, it is enough to notice that X_m satisfies the third equation in (3.7), that

is

$$\frac{(E + \bar{V}^-) + m}{\gamma_+ + m} \leq \frac{(E + \bar{V}^-) + M}{\gamma_+ + M}.$$

Since the function $t \rightarrow (E + \bar{V}^- + t)/(\gamma_+ + t)$ is decreasing and $m < M$ we conclude that X_m is not in the domain D . Analogously one shows that X_M is not in D . By replacing γ_+ by γ_- in the computations above, one shows that Y_m and Y_M are in the domain D .

Now, if $X_0 \notin [-1, 1]^2$, i.e. $E + \bar{V}^- > \gamma_-$ then the same computation as done above shows that Y_m and Y_M are not in D and as a consequence D is empty; and if $E + \bar{V}^- < \gamma_+$ then X_m and X_M are not in D and as a consequence D is empty.

So, we are restricted to the case $\gamma_+ \leq E + \bar{V}^- \leq \gamma_-$. It remains to compute

$$\sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(Y_m, y_-, y_+), F(Y_M, y_-, y_+) \right\},$$

where $\Gamma := \{(y_-, y_+) : \gamma_+ \leq E + \bar{V}^- \leq \gamma_-\}$. Observe now that whatever the value of y_- is, we have that

$$\begin{aligned} F(Y_m, y_-, y_+) &= \frac{(E + \bar{V}^-) + m}{\gamma_- + m} ((-s)(y_-) + \gamma_-), \\ F(Y_M, y_-, y_+) &= \frac{(E + \bar{V}^-) + M}{\gamma_- + M} ((-s)(y_-) + \gamma_-). \end{aligned}$$

Now we notice that:

$$\begin{aligned} (-s)(y_-) + \gamma_- &= -\left(s \left(\frac{\gamma_- + \log(1 + e^{\varphi_-})}{\varphi_-} \right) - \gamma_- \right) \\ &= -\left(s \left(\frac{-\gamma_- - \log(1 + e^{-\varphi_-})}{\varphi_-} \right) - \gamma_- \right) \\ &= -\left(-S_{1-\rho_-, 1-\rho_-}(-\gamma_-) - \gamma_- \right) \\ &= -J_{\rho_-, \rho_-}(-\gamma_-). \end{aligned}$$

In the second equality above, we wrote y_- in terms of γ_- , in the third equality we used the fact that $s(\theta) = s(1 - \theta)$ for all $\theta \in (0, 1)$ and in fourth equality we used that for any $\rho \in [0, 1]$,

$$S_{\rho, \rho} = S_{1-\rho, 1-\rho} \tag{C.1}$$

together with the definition of $J_{\rho, \rho}$ given above. The same argument also shows that $(-s)(y_+) + \gamma_+ = -J_{\rho_+, \rho_+}(-\gamma_+)$. Then we conclude that

$$\begin{aligned} F(Y_m, y_-, y_+) &= -\frac{(E + \bar{V}^-) + m}{\gamma_- + m} J_{\rho_-, \rho_-}(-\gamma_-) \\ F(Y_M, y_-, y_+) &= -\frac{(E + \bar{V}^-) + M}{\gamma_- + M} J_{\rho_-, \rho_-}(-\gamma_-). \end{aligned}$$

Observe that since the function $t \rightarrow -((E + \bar{V}^-) + t)/(\gamma_- + t)$ is decreasing, $m \leq M$ and $J_{\rho_-, \rho_-}(-\gamma_-) \geq 0$, we get that $F(Y_m, y_-, y_+) \geq F(Y_M, y_-, y_+)$.

Now we recall some properties of the function $J_{\rho, \rho}$, for $\rho \in [0, 1]$. At first we notice that by (C.1), we have that $J_{\rho, \rho} = J_{1-\rho, 1-\rho}$. The function $J_{\rho, \rho}$ is convex and positive apart from the point $-s(\rho)$ where it vanishes. As a consequence the function J_{ρ_-, ρ_-} is convex, non-negative and finite on $[\log(1 + e^{\varphi_-}); \log(1 + e^{-\varphi_-})]$ and has a minimum equal to 0 at the point $-s(\rho_-) = \gamma_-(\rho_-)$. Analogously, the function J_{ρ_+, ρ_+} is convex, non-negative and finite on $[\log(1 + e^{-\varphi_+}); \log(1 + e^{\varphi_+})]$ and has a minimum equal to 0 at the point

$-s(\rho_+) = \gamma_+(\rho_+)$. Since $-s(\rho_-) \leq m = -s(\rho_+)$ then for all $y \in I := [\log(1 + e^{-\phi_+}); m]$ we have that $J_{\rho_-, \rho_-}(y) \geq J_{\rho_+, \rho_+}(y)$. In particular, since $\gamma_- \in I$ we obtain that $J_{\rho_-, \rho_-}(-\gamma_-) \geq J_{\rho_+, \rho_+}(-\gamma_-)$. Putting together the previous observations, the fact that $E + \bar{V}^- \geq \gamma_+ \geq m$ and $\gamma_- > -m$, we conclude that

$$F(Y_m, y_-, y_+) = -\frac{(E + \bar{V}^-) + m}{\gamma_- + m} J_{\rho_-, \rho_-}(-\gamma_-) \leq -\frac{(E + \bar{V}^-) + m}{\gamma_- + m} J_{\rho_+, \rho_+}(-\gamma_-).$$

On the other hand, by computing the derivative of the function

$$G(\gamma_-) = -\frac{(E + \bar{V}^-) + m}{\gamma_- + m} J_{\rho_+, \rho_+}(-\gamma_-)$$

with respect to γ_- and using the fact that the function $J_{\rho_+, \rho_+}(-\gamma_-)$ is convex at the point γ_- , we conclude that $G(\cdot)$ is decreasing. Then,

$$\begin{aligned} \sup_{(y_-, y_+) \in \Gamma} F(Y_m, y_-, y_+) &\leq \sup_{\gamma_- \geq E + \bar{V}^-} G(\gamma_-) \\ &= G(E + \bar{V}^-) \\ &= -J_{\rho_+, \rho_+}(- (E + \bar{V}^-)). \end{aligned}$$

Now we rewrite $F(X_0, y_-, y_+)$ as

$$F(X_0, y_-, y_+) = -\frac{(E + \bar{V}^-) - \gamma_+}{\gamma_- - \gamma_+} J_{\rho_-, \rho_-}(-\gamma_-) - \frac{\gamma_- - (E + \bar{V}^-)}{\gamma_- - \gamma_+} J_{\rho_+, \rho_+}(-\gamma_+).$$

By computing the derivative with respect to γ_+ of $F(X_0, y_-, y_+)$, and noticing that both J_{ρ_+, ρ_+} and J_{ρ_-, ρ_-} are convex, we obtain that $F(X_0, y_-, y_+)$ is increasing as a function of γ_+ . Then

$$\begin{aligned} \sup \left\{ F(X_0, y_-, y_+) : (y_-, y_+) \in \Gamma \right\} &\leq \sup \left\{ F(X_0, y_-, y_+) : \gamma_- \geq E + \bar{V}^- \right\} \\ &= -J_{\rho_+, \rho_+}(- (E + \bar{V}^-)). \end{aligned}$$

Putting together the previous computations we obtain that

$$\begin{aligned} \sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(Y_m, y_-, y_+), F(Y_M, y_-, y_+) \right\} &= \sup_{\gamma_+ \leq E + \bar{V}^- \leq \gamma_-} F(X_0, y_-, y_+) \\ &= \sup_{\gamma_+ = E + \bar{V}^-} \left\{ F(X_0, y_-, y_+) \right\} \\ &= -J_{\rho_+, \rho_+}(- (E + \bar{V}^-)). \end{aligned}$$

(a.2) Case $\gamma_- < \gamma_+$: A simple computation shows that X_0 belongs to the square $[-1, 1]^2$ if and only if $\gamma_- \leq E + \bar{V}^- \leq \gamma_+$. As above, we first restrict to $\gamma_- \leq E + \bar{V}^- \leq \gamma_+$. A simple computation as performed above, shows that X_m and X_M are in D and Y_m and Y_M are not in D .

Now, if $X_0 \notin [-1, 1]^2$, i.e. $E + \bar{V}^- > \gamma_+$ then the same computation as done above shows that X_m and X_M are not in D and as a consequence D is empty; and if $E + \bar{V}^- < \gamma_-$ then Y_m and Y_M are not in D and as a consequence D is empty.

So, we are restricted to the case $\gamma_- \leq E + \bar{V}^- \leq \gamma_+$. Then, it remains to compute

$$\sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(X_m, y_-, y_+), F(X_M, y_-, y_+) \right\},$$

where $\Gamma := \{(y_-, y_+) : \gamma_- \leq E + \bar{V}^- \leq \gamma_+\}$. By inverting the role of ρ_- with ρ_+ and of γ_- with γ_+ in the proof of the previous case, we obtain here that the previous supremum equals to $-J_{\rho_+, \rho_+} \left(-(E + \bar{V}^-) \right)$.

(b) $\gamma_-(0) < -m$: In this case we have that $\gamma_- < -m < \gamma_+$. As above, we have to check whether the points X_0, X_m, X_M, Y_m, Y_M are in the domain D or not.

The point X_0 belongs to $[-1, 1]^2$ if and only if $\gamma_- \leq E + \bar{V}^- \leq \gamma_+$. If $X_0 \notin [-1, 1]^2$, i.e. $E + \bar{V}^- > \gamma_+$, then X_m and X_M are not in the domain D and as a consequence D is empty.

Then we restrict to $\gamma_- \leq E + \bar{V}^- \leq \gamma_+$. A simple computation shows that X_m and X_M belong to the domain D , but Y_m, Y_M are not in D . So we have to compute

$$\sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(X_m, y_-, y_+), F(X_M, y_-, y_+) \right\},$$

where $\Gamma := \{(y_-, y_+) : \gamma_- \leq E + \bar{V}^- \leq \gamma_+\}$.

As above easily we can show that $F(X_m, y_-, y_+) \geq F(X_M, y_-, y_+)$. Now we have to compare $F(X_m, y_-, y_+)$ with $F(X_0, y_-, y_+)$. A simple computation shows that $F(X_0, y_-, y_+)$ can be written as

$$F(X_0, y_-, y_+) = -\frac{\gamma_+ - (E + \bar{V}^-)}{\gamma_+ - \gamma_-} J_{\rho_-, \rho_-}(-\gamma_-) - \frac{\gamma_- - (E + \bar{V}^-)}{\gamma_- - \gamma_+} J_{\rho_+, \rho_+}(-\gamma_+).$$

Since J_{ρ_-, ρ_-} is a positive function, $E + \bar{V}^- \leq \gamma_-$ and $\gamma_+ > \gamma_-$ we have that

$$F(X_0, y_-, y_+) \leq -\frac{\gamma_- - (E + \bar{V}^-)}{\gamma_- - \gamma_+} J_{\rho_+, \rho_+}(-\gamma_+).$$

Now, since the function $t \rightarrow -(E + \bar{V}^- + t)/(\gamma_+ + t)$ is decreasing and $m > -\gamma_-$ we obtain that $F(X_0, y_-, y_+) \leq F(X_m, y_-, y_+)$.

It follows that

$$\begin{aligned} & \sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(X_m, y_-, y_+), F(X_M, y_-, y_+) \right\} \\ &= \sup_{(y_-, y_+) \in \Gamma} F(X_m, y_-, y_+) \\ &= \sup_{\gamma_+ \geq E + \bar{V}^-} F(X_m, y_-, y_+) \\ &= -J_{\rho_+, \rho_+} \left(-(E + \bar{V}^-) \right). \end{aligned}$$

By the conclusions above together with (3.6), we obtain that the restriction of the entropy function S^- to $[E_{-\infty}^-; E_{-\infty}^+]$ is given by

$$\begin{aligned} S^-(E) &= \\ &= \begin{cases} -(E + \bar{V}^-), & -\log(2) \leq E + \bar{V}^- \leq \frac{\varphi_0}{1+e^{-\varphi_0}} - \log(1 + e^{\varphi_0}), \\ S_{\rho_0, \rho_0} \left(-(E + \bar{V}^-) \right), & \frac{\varphi_0}{1+e^{-\varphi_0}} - \log(1 + e^{\varphi_0}) < E + \bar{V}^- \leq -\log(1 + e^{-\varphi_0}). \end{cases} \end{aligned}$$

Above we used the equality $S_{\rho_+, \rho_+}(E) = E - J_{\rho_+, \rho_+}(E)$. To conclude, we notice that for any $\rho = e^\varphi / (1 + e^\varphi) \in [0, 1]$,

$$s(\rho) = \frac{\varphi}{1 + e^{-\varphi}} - \log(1 + e^\varphi) = \frac{-\varphi}{1 + e^\varphi} - \log(1 + e^{-\varphi}).$$

Now, we prove the last assertion of the theorem. As above, we consider the case $\varphi_0 = \varphi_+$ the other case being similar. We have to split now into two cases, whether $E + \bar{V}^- > s(\rho_+)$ or $E + \bar{V}^- \leq s(\rho_+)$. We start by the later.

Assume $E + \bar{V}^- \leq s(\rho_+)$ and let ρ be a profile such that $\mathbb{S}(\rho) = S^-(E)$ and $\mathbb{S}(\rho) + V^-(\rho) = E + \bar{V}^-$. With the notations of Proposition 3.3, we have $S^-(E) = \mathbb{S}(\rho) = \mathbb{S}(H'_\rho) \leq \mathbb{S}(G'_\rho)$. Moreover, we have seen in the proof of Proposition 3.1 that $(\mathbb{S} + V^-)(G'_\rho) = (\mathbb{S} + V^-)(\rho)$. We claim now that $\rho = G'_\rho$. Indeed, let $(a, b) \in [-1, 1]$ be a maximal interval where $G_\rho < H_\rho$ (which implies $H_\rho(a) = G_\rho(a)$ and $H_\rho(b) = G_\rho(b)$). Since G_ρ is the convex envelope of H_ρ , it implies that G_ρ is linear on $[a, b]$. By Jensen's inequality one has that

$$\begin{aligned} \int_a^b (-s)(H'_\rho(x))dx &> (b-a)(-s) \left(\frac{1}{b-a} \int_a^b H'_\rho(x)dx \right) \\ &= (b-a)(-s) \left(\frac{H_\rho(b) - H_\rho(a)}{b-a} \right) = (b-a)(-s) \left(\frac{G_\rho(b) - G_\rho(a)}{b-a} \right) \\ &= \int_a^b (-s)(G'_\rho(x))dx. \end{aligned}$$

Thus, if $H_\rho \neq G_\rho$ we can find a profile $\bar{\rho}$ (i.e. $1 - G'_\rho$) which satisfies the constraint $\mathbb{S}(\bar{\rho}) + V^-(\bar{\rho}) = E + \bar{V}^-$ and such that $\mathbb{S}(\bar{\rho}) > \mathbb{S}(\rho) = S^-(E)$. Since this is not possible we get that $H_\rho = G_\rho$, i.e. ρ is a non-increasing profile. In particular it implies that $1 - \rho = g$ where g is a maximizer of (3.3). In the proof of Proposition 3.3 we have seen that $S^-(E) = -(E + \bar{V}^-)$ corresponds to the case where the supremum $\sup_{k \in K} F(k) = 0$, which is equivalent to $(x_\pm \mp 1)(y_\pm - \rho_\pm) = 0$. For such a 4-tuple $(x_-, x_+, y_-, y_+) \in K$, a maximizer g of (3.3) is then given by any non-decreasing function on $[x_-, x_+]$ taking values in $[\rho_-, \rho_+]$, constant equal to ρ_- on $[-1, x_-]$ and to ρ_+ on $[x_+, 1]$ and such that (3.4) is satisfied. Thus the set of maximizers ρ of $S^-(E)$, when $E + \bar{V}^- \in [s(\rho_+), s(\rho_-)]$ is given by the set of non-increasing profiles ρ such that $1 - \rho_+ \leq \rho \leq 1 - \rho_-$ and satisfying $\mathbb{S}(\rho) = -(E + \bar{V}^-)$.

Now assume that $E + \bar{V}^- > s(\rho_+)$. In all the cases (a_1) , (a_2) and (b) above, the supremum of F is attained at X_0, X_0 and X_m , respectively, and $\gamma_+ = E + \bar{V}^-$, which implies that $x_- = x_+ = -1$. Then, the function g realizing the supremum $\sup_\rho \int_{-1}^1 (-s)(g(x))dx$ with the constraint $\int_{-1}^1 g(x)dx = (1 - x_+)y_+$ is constant and equals to y_+ . Therefore, the profile u_{ρ_+} is given by $1 - y_+$. Using the definition of y_+ and the fact that $\gamma_+ = E + \bar{V}^-$, it follows that

$$u_{\rho_+} \equiv \frac{\log(\rho_+) - E + \bar{V}^-}{\log(\rho_+) - \log(1 - \rho_+)}.$$

Finally, putting together (1.6), (1.8), (1.4) and the expression for u_{ρ_+} we recover the expression for $S^-(E)$ that is $S^-(E) = S_{\rho_+, \rho_+}(-E + \bar{V}^-)$.

C.2. The case $\rho_- < \rho_+ \leq \frac{1}{2}$:

This case corresponds to $\varphi_- < \varphi_+ \leq 0$, therefore $m = (-s)(\rho_-)$ and $M = (-s)(\rho_+)$. Since $\gamma'(y_\pm) = \varphi_\pm < 0$, then both functions γ_\pm are decreasing. Notice that $\gamma_+(1) = E_{-\infty} + \bar{V}^-$ and $\gamma_-(0) = E_{-\infty}^+ + \bar{V}^-$. Then $-M = \gamma_+(\rho_+) \geq \gamma_+(1) = -\log(1 + e^{-\varphi_+})$. Analogously, $-m = \gamma_-(\rho_-) \leq \gamma_-(0) = -\log(1 + e^{\varphi_-})$. As a consequence we have the following inequalities:

$$-\log(1 + e^{-\varphi_+}) = \gamma_+(1) \leq \gamma_+(\rho_+) = -M < -m \leq \gamma_-(\rho_-) \leq \gamma_-(0) = -\log(1 + e^{\varphi_-}).$$

These inequalities imply that $\gamma_+ < \gamma_-$. As in the previous case we have to check whether the intersection points are in the domain D . At first we notice that X_0 belongs to $[-1, 1]^2$ if and only if $\gamma_+ \leq E + \bar{V}^- \leq \gamma_-$. Since $-(E + \bar{V}^-) \notin [m, M]$ we have only to distinguish two cases: $\gamma_+ \leq E + \bar{V}^- < -M$ and $-m \leq E + \bar{V}^- < \gamma_-$.

If X_0 is not in $[-1, 1]^2$, i.e. $E + \bar{V}^- > \gamma_-$ then X_m and X_M are not in D and as a consequence D is empty; and if $E + \bar{V}^- < \gamma_+$ then Y_m and Y_M are not in D and as a consequence D is empty. So we restrict to $\gamma_+ \leq E + \bar{V}^- < -M$ and $-m \leq E + \bar{V}^- < \gamma_-$. We start by the former.

(a) $\gamma_+ \leq E + \bar{V}^- < -M$: In this case, a simple computation shows that X_m and X_M are in D , but Y_m and Y_M are not in D . Then we have to compute

$$\sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(X_m, y_-, y_+), F(X_M, y_-, y_+) \right\},$$

where $\Gamma := \{(y_-, y_+) : \gamma_+ \leq E + \bar{V}^- < -M\}$.

As above, by noticing that the function $t \rightarrow -((E + \bar{V}^-) + t)/(\gamma_+ + t)$ is increasing, $m \leq M$ and $J_{\rho_+, \rho_+}(-\gamma_+) \geq 0$, we get that $F(X_m, y_-, y_+) \leq F(X_M, y_-, y_+)$. On the other hand

$$F(X_0, y_-, y_+) = -\frac{(E + \bar{V}^-) - \gamma_+}{\gamma_- - \gamma_+} J_{\rho_-, \rho_-}(-\gamma_-) - \frac{\gamma_- - (E + \bar{V}^-)}{\gamma_- - \gamma_+} J_{\rho_+, \rho_+}(-\gamma_+).$$

Since the function J_{ρ_-, ρ_-} is positive, $\gamma_- > \gamma_+$ and $E + \bar{V}^- > \gamma_+$, we have that

$$F(X_0, y_-, y_+) \leq -\frac{\gamma_- - (E + \bar{V}^-)}{\gamma_- - \gamma_+} J_{\rho_+, \rho_+}(-\gamma_+) = -\frac{(E + \bar{V}^-) - \gamma_-}{\gamma_+ - \gamma_-} J_{\rho_+, \rho_+}(-\gamma_+).$$

Finally, since the function $t \rightarrow -((E + \bar{V}^-) + t)/(\gamma_+ + t)$ is increasing, $-\gamma_- < M$ and $J_{\rho_+, \rho_+}(-\gamma_+) \geq 0$ we obtain that $F(X_0, y_-, y_+) \leq F(X_M, y_-, y_+)$. Since the function J_{ρ_+, ρ_+} is convex, with a unique minimum at M equal to 0, it follows that

$$\begin{aligned} \sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(X_m, y_-, y_+), F(X_M, y_-, y_+) \right\} &= \sup_{(y_-, y_+) \in \Gamma} F(X_M, y_-, y_+) \\ &= \sup_{\gamma_+ \leq (E + \bar{V}^-)} F(X_M, y_-, y_+) \\ &= -J_{\rho_+, \rho_+} \left(-(E + \bar{V}^-) \right). \end{aligned}$$

(b) $-m < E + \bar{V}^- \leq \gamma_-$: In this case, a simple computation shows that Y_m and Y_M are in D , but X_m and X_M are not in D . Then we have to compute

$$\sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(Y_m, y_-, y_+), F(Y_M, y_-, y_+) \right\},$$

where $\Gamma := \{(y_-, y_+) : -m < E + \bar{V}^- \leq \gamma_-\}$.

As above, by noticing that the function $t \rightarrow -((E + \bar{V}^-) + t)/(\gamma_- + t)$ is increasing, $m \leq M$ and $J_{\rho_-, \rho_-}(-\gamma_-) \geq 0$, we get $F(Y_m, y_-, y_+) \leq F(Y_M, y_-, y_+)$.

As above we can show that $F(X_0, y_-, y_+) \leq F(Y_M, y_-, y_+)$ and as a consequence

$$\begin{aligned} \sup_{(y_-, y_+) \in \Gamma} \left\{ F(X_0, y_-, y_+), F(Y_m, y_-, y_+), F(Y_M, y_-, y_+) \right\} &= \sup_{(y_-, y_+) \in \Gamma} F(Y_M, y_-, y_+) \\ &= \sup_{\gamma_- \leq (E + \bar{V}^-)} F(Y_M, y_-, y_+) \\ &= -J_{\rho_-, \rho_-} \left(-(E + \bar{V}^-) \right). \end{aligned}$$

Collecting the previous facts and by (3.6), we have that the restriction of the entropy function S^- to $[E_{-\infty}^-; E_{-\infty}^+]$ is given by

$$S^-(E) = \begin{cases} S_{\rho_+, \rho_+} \left(-(E + \bar{V}^-) \right), & -\log(1 + e^{-\varphi_+}) \leq E + \bar{V}^- < \frac{-\varphi_+}{1 + e^{\varphi_+}} - \log(1 + e^{-\varphi_+}), \\ -(E + \bar{V}^-), & \frac{-\varphi_+}{1 + e^{\varphi_+}} - \log(1 + e^{-\varphi_+}) \leq E + \bar{V}^- \leq \frac{-\varphi_-}{1 + e^{\varphi_-}} - \log(1 + e^{-\varphi_-}), \\ S_{\rho_-, \rho_-} \left(-(E + \bar{V}^-) \right), & \frac{-\varphi_-}{1 + e^{\varphi_-}} - \log(1 + e^{-\varphi_-}) < E + \bar{V}^- \leq -\log(1 + e^{\varphi_-}) \end{cases}$$

Now, we prove the last assertion of the theorem. As above, we have to split into several cases, whether $E + \bar{V}^- < s(\rho_+)$, $s(\rho_+) \leq E + \bar{V}^- \leq s(\rho_-)$ or $E + \bar{V}^- > s(\rho_-)$. We start by the first case, the third being completely similar. Analogously to what we have done for $\rho_- \leq 1/2 \leq \rho_+$, it is enough to notice that in the cases (a) and (b) above, the supremum is attained at X_M and Y_M , respectively, with $\gamma_+ = E + \bar{V}^-$, which implies that $x_- = x_+ = -1$. The rest of the argument follows as above. The second case, follows by reasoning as in the case $\rho_- \leq \frac{1}{2} \leq \rho_+$.

C.3. The case $\frac{1}{2} \leq \rho_- < \rho_+$:

Repeating the same computations as performed in the previous situation, we can show that the restriction of the entropy function S^- to $[E_{-\infty}^-; E_{-\infty}^+]$ is given by

$$S^-(E) = \begin{cases} S_{\rho_-, \rho_-} \left(-(E + \bar{V}^-) \right), & -\log(1 + e^{\varphi_-}) \leq E + \bar{V}^- < \frac{-\varphi_-}{1 + e^{\varphi_-}} - \log(1 + e^{-\varphi_-}), \\ -(E + \bar{V}^-), & \frac{-\varphi_-}{1 + e^{\varphi_-}} - \log(1 + e^{-\varphi_-}) \leq E + \bar{V}^- \leq \frac{-\varphi_+}{1 + e^{\varphi_+}} - \log(1 + e^{-\varphi_+}) \\ S_{\rho_+, \rho_+} \left(-(E + \bar{V}^-) \right), & \frac{-\varphi_+}{1 + e^{\varphi_+}} - \log(1 + e^{-\varphi_+}) < E + \bar{V}^- \leq -\log(1 + e^{-\varphi_+}). \end{cases}$$

We notice that to prove the last assertion of the theorem is is enough to invert the role of ρ_- with ρ_+ in the proof of the previous case.

APPENDIX D. PROOF OF THEOREM 3.4

To prove this theorem we compute explicitly the Legendre transform of P^- , namely, $\tilde{S}^-(E) = \inf_{\theta \in \mathbb{R}} \{\theta E - P^-(\theta)\}$ and we show that it coincides with the expression for $S^-(E)$ obtained in the previous section. Since the Legendre transform is a one to one correspondence between concave functions, this is sufficient to conclude. We denote by P_0 the function defined by $P_0(\theta) = -\theta \log(m(\theta))$.

D.1. **The case $\rho_- \leq \frac{1}{2} \leq \rho_+$:**

This case corresponds to $\varphi_- \leq 0 \leq \varphi_+$. Recall that $\varphi_0 = \sup(|\varphi_-|, |\varphi_+|)$. Since

$$m(\theta) = \min(f_\theta(\varphi_-), f_\theta(\varphi_+)) = \begin{cases} f_\theta(\varphi_0), & \theta < -1, \\ 1, & \theta = -1, \\ f_\theta(0), & \theta > -1, \end{cases},$$

we have that

$$P_0(\theta) = \begin{cases} -\theta \log(1 + e^{-\varphi_0}) - \log(1 + e^{\theta\varphi_0}), & \theta < -1, \\ 0, & \theta = -1, \\ -(\theta + 1)\log 2, & \theta > -1. \end{cases}$$

As a consequence

$$P'_0(\theta) = \begin{cases} -\log(1 + e^{-\varphi_0}) - \frac{\varphi_0}{1 + e^{-\theta\varphi_0}}, & \theta < -1, \\ -\log(2), & \theta > -1. \end{cases}$$

The function P_0 is differentiable everywhere except for $\theta = -1$. A simple computation show that P'_0 is decreasing in $(-\infty, -1)$. We have that

$$\begin{aligned} P'_0(-1^-) &= \lim_{\substack{\theta \rightarrow -1 \\ \theta < -1}} P'_0(\theta) = -\log(1 + e^{-\varphi_0}) - \frac{\varphi_0}{1 + e^{\varphi_0}}, \\ P'_0(-\infty) &= \lim_{\theta \rightarrow -\infty} P'_0(\theta) = -\log(1 + e^{-\varphi_0}). \end{aligned}$$

Observe that $P'_0(-\infty) > P'_0(-1^-) > -\log(2)$, which is a consequence of the function $t \rightarrow -\log(1 + e^{-t}) - t/(1 + e^t)$ being decreasing on $(-\infty, 0]$. This implies that the function P_0 is concave. Now we compute $\tilde{S}^-(E)$. By the previous observations we have that

$$\tilde{S}^-(E) = \inf \left\{ \inf_{\theta > -1} \{ \theta(E + \bar{V}^-) - P_0(\theta) \}, -(E + \bar{V}^-), \inf_{\theta < -1} \{ \theta(E + \bar{V}^-) - P_0(\theta) \} \right\}.$$

Now, for $\theta > -1$ we have that

$$\inf_{\theta > -1} \{ \theta(E + \bar{V}^-) - P_0(\theta) \} = \log(2) + \inf_{\theta > -1} \{ \theta((E + \bar{V}^-) + \log(2)) \}.$$

This equals to $-(E + \bar{V}^-)$ if $(E + \bar{V}^-) > -\log(2)$ and equals $-\infty$ if $(E + \bar{V}^-) < -\log(2)$. On the other hand, for $\theta < -1$ we have that

$$\inf_{\theta < -1} \{ \theta(E + \bar{V}^-) - P_0(\theta) \} = \inf_{\theta < -1} \{ \theta(E + \bar{V}^-) - P(-\varphi_0, -\theta) \},$$

where $P(\cdot, \cdot)$ is defined in (1.5). Let $I_1 := \left[\frac{-\varphi_0}{1+e^{\varphi_0}} - \log(1+e^{-\varphi_0}); -\log(1+e^{-\varphi_0}) \right] \subset (-\infty, 0)$. We have that

$$\begin{aligned} \inf_{\theta \in \mathbb{R}} \left\{ \theta(E + \bar{V}^-) - P(-\varphi_0, -\theta) \right\} &= \frac{1}{2} \inf_{\theta \in \mathbb{R}} \left\{ \theta 2(E + \bar{V}^-) - 2P(-\varphi_0, -\theta) \right\} \\ &= \frac{1}{2} \inf_{\theta \in \mathbb{R}} \left\{ \theta(-2(E + \bar{V}^-)) - 2P(-\varphi_0, \theta) \right\} \\ &= -s \left(\frac{-(E + \bar{V}^-) - \log(1+e^{-\varphi_0})}{\varphi_0} \right) \\ &= S_{\rho_0, \rho_0} \left(-(E + \bar{V}^-) \right), \end{aligned}$$

for any $E + \bar{V}^- \in I_1$. Then, we conclude that

$$\inf_{\theta > -1} \left\{ \theta(E + \bar{V}^-) - P_0(\theta) \right\} = S_{\rho_0, \rho_0} \left(-(E + \bar{V}^-) \right),$$

for $E + \bar{V}^- \in I_1$. Now we look for $\tilde{S}^-(E)$ for $E + \bar{V}^-$ outside I_1 . Let $I_1^- := \left(-\infty; \frac{-\varphi_0}{1+e^{\varphi_0}} - \log(1+e^{-\varphi_0}) \right)$ and $I_1^+ := \left(-\log(1+e^{-\varphi_0}); +\infty \right)$. Then, if $E + \bar{V}^- \in I_1^-$, we have that $\text{sign}(E + \bar{V}^- - P'_0(\theta)) = \text{sign}((E + \bar{V}^-) - P'_0(-\infty)) = \text{sign}((E + \bar{V}^-) + \log(1+e^{-\varphi_0})) < 0$, and as a consequence the infimum is attained at $\theta = -1$ and in this case $\tilde{S}^-(E) = -(E + \bar{V}^-)$. On the other hand if $E + \bar{V}^- \in I_1^+$, we have that $\text{sign}(E + \bar{V}^- - P'_0(\theta)) = \text{sign}((E + \bar{V}^-) - P'_0(-\infty)) > 0$, and as a consequence the infimum is attained at $\theta = -\infty$ and in this case $\tilde{S}^-(E) = -\infty$.

Finally we conclude that \tilde{S}^- when restricted to the energy band $[E_{-\infty}^-; E_{-\infty}^+]$ is given by

$$\begin{aligned} \tilde{S}^-(E) &= \\ &= \begin{cases} -(E + \bar{V}^-), & -\log(2) \leq E + \bar{V}^- \leq \frac{-\varphi_0}{1+e^{\varphi_0}} - \log(1+e^{-\varphi_0}), \\ S_{\rho_0, \rho_0} \left(-(E + \bar{V}^-) \right), & \frac{-\varphi_0}{1+e^{\varphi_0}} - \log(1+e^{-\varphi_0}) < E + \bar{V}^- \leq -\log(1+e^{-\varphi_0}) \end{cases} \end{aligned}$$

and equal to $-\infty$ outside the energy band. Thus \tilde{S}^- coincides with S^- .

D.2. The case $\rho_- < \rho_+ \leq \frac{1}{2}$:

In this case we have that

$$m(\theta) = \min(f_\theta(\varphi_-), f_\theta(\varphi_+)) = \begin{cases} f_\theta(\varphi_-), & \theta < -1, \\ 1, & \theta = -1, \\ f_\theta(\varphi_+), & \theta > -1, \end{cases}$$

and as a consequence

$$P'_0(\theta) = \begin{cases} -\frac{\varphi_-}{1+e^{-\theta\varphi_-}} - \log(1+e^{-\varphi_-}), & \theta < -1, \\ -\frac{\varphi_+}{1+e^{-\theta\varphi_+}} - \log(1+e^{-\varphi_+}), & \theta > -1. \end{cases}$$

The function P_0 is differentiable everywhere except for $\theta = -1$. We have that P'_0 is decreasing on $(-\infty, -1)$ and

$$P'_0(-\infty) > P'_0(-1^-) = \lim_{\substack{\theta \rightarrow -1 \\ \theta < -1}} P'_0(\theta) > \lim_{\substack{\theta \rightarrow -1 \\ \theta > -1}} P'_0(\theta) = P'_0(-1^+) > P'_0(+\infty)$$

which is a consequence of the function $t \rightarrow \log(1 + e^{-t}) + t/(1 + e^t)$ being increasing on $(-\infty, 0)$. This implies that the function P_0 is concave. Now we compute $\tilde{S}^-(E)$. From the previous observations it follows that

$$\tilde{S}^-(E) = \inf \left\{ \inf_{\theta > -1} \{ \theta(E + \bar{V}^-) - P_0(\theta) \}, -(E + \bar{V}^-), \inf_{\theta < -1} \{ \theta(E + \bar{V}^-) - P_0(\theta) \} \right\}.$$

We start by the case $\theta > -1$. Since P_0 is concave, then P'_0 is decreasing. A simple computation shows that $P'_0(\theta) = E + \bar{V}^-$ for

$$\theta(E) := -\frac{1}{\varphi_+} \log \left(\frac{-\varphi_+ - (E + \bar{V}^-) - \log(1 + e^{-\varphi_+})}{E + \bar{V}^- + \log(1 + e^{-\varphi_+})} \right).$$

Since P'_0 is decreasing, its image is given by $I_1 := [P'_0(+\infty); P'_0(-1)]$, that is $I_1 = [-\log(1 + e^{-\varphi_+}); \frac{-\varphi_+}{1 + e^{\varphi_+}} - \log(1 + e^{-\varphi_+})]$. Then, for $E + \bar{V}^- \in I_1$ we have that

$$\tilde{S}^-(E) = \theta(E)(E + \bar{V}^-) - P_0(\theta(E)) = S_{\rho_+, \rho_+} \left(-(E + \bar{V}^-) \right).$$

Now we look for $\tilde{S}^-(E)$ for $E + \bar{V}^-$ outside I_1 . Let $I_1^- := (-\infty; -\log(1 + e^{-\varphi_+}))$ and $I_1^+ := (\frac{-\varphi_+}{1 + e^{\varphi_+}} - \log(1 + e^{-\varphi_+}); +\infty)$. Then, if $E + \bar{V}^- \in I_1^-$, we have that $\text{sign}(E + \bar{V}^- - P'_0(\theta)) = \text{sign}(E + \bar{V}^- - P'_0(+\infty)) = \text{sign}(E + \bar{V}^- + \log(1 + e^{-\varphi_+})) < 0$, and as a consequence the infimum is attained at $\theta = +\infty$ and in this case $\tilde{S}^-(E) = -\infty$. On the other hand if $E + \bar{V}^- \in I_1^+$, we have that $\text{sign}(E + \bar{V}^- - P'_0(\theta)) = \text{sign}(E + \bar{V}^- - P'_0(+\infty)) > 0$, and as a consequence the infimum is attained at $\theta = -1$ and in this case $\tilde{S}^-(E) = -\infty$.

Now we look to the case $\theta < -1$. Since P_0 is concave, then P'_0 is decreasing. A simple computation shows that $P'_0(\theta) = E + \bar{V}^-$ for

$$\theta(E) := -\frac{1}{\varphi_-} \log \left(\frac{-\varphi_- - (E + \bar{V}^-) - \log(1 + e^{-\varphi_-})}{E + \bar{V}^- + \log(1 + e^{-\varphi_-})} \right).$$

Since P'_0 is decreasing, its image is given by $I_2 := [P'_0(-1); P'_0(-\infty)]$, that is $I_2 = [\frac{-\varphi_-}{1 + e^{\varphi_-}} - \log(1 + e^{-\varphi_-}); -\log(1 + e^{\varphi_-})]$. Then, for $E + \bar{V}^- \in I_2$ we have that $\tilde{S}^-(E) := \theta(E)(E + \bar{V}^-) - P_0(\theta(E))$ which can be written as $S_{\rho_-, \rho_-} \left(-(E + \bar{V}^-) \right)$. Now we look for $\tilde{S}^-(E)$ for $E + \bar{V}^-$ outside I_2 . Let $I_2^- := (-\infty; \frac{-\varphi_-}{1 + e^{\varphi_-}} - \log(1 + e^{-\varphi_-}))$ and $I_2^+ := (-\log(1 + e^{\varphi_-}); +\infty)$. Then, if $E + \bar{V}^- \in I_2^-$, we have that $\text{sign}(E + \bar{V}^- - P'_0(\theta)) = \text{sign}(E + \bar{V}^- - P'_0(-\infty)) = \text{sign}(E + \bar{V}^- + \log(1 + e^{\varphi_-})) < 0$, and as a consequence the infimum is attained at $\theta = -1$ and in this case $\tilde{S}^-(E) = -(E + \bar{V}^-)$. On the other hand if $E + \bar{V}^- \in I_2^+$, we have that $\text{sign}(E + \bar{V}^- - P'_0(\theta)) = \text{sign}(E + \bar{V}^- - P'_0(-1)) > 0$, and as a consequence the infimum is attained at $\theta = -\infty$ and in this case $\tilde{S}^-(E) = -\infty$.

Now, a simple computation shows that the function $t \rightarrow -\log(1 + e^{-t}) - t/(1 + e^t)$ is decreasing on $(-\infty, 0]$ so that the sets I_1 and I_2 do not intersect. The restriction of the

function \tilde{S}^- to $[E_{-\infty}^-; E_{-\infty}^+]$ has the expression:

$$\begin{aligned} \tilde{S}^-(E) &= \\ &= \begin{cases} S_{\rho_+, \rho_+} \left(-(E + \bar{V}^-) \right), & -\log(1 + e^{-\varphi_+}) \leq E + \bar{V}^- < \frac{-\varphi_+}{1+e^{\varphi_+}} - \log(1 + e^{-\varphi_+}), \\ -(E + \bar{V}^-), & \frac{-\varphi_+}{1+e^{\varphi_+}} - \log(1 + e^{-\varphi_+}) \leq E + \bar{V}^- \leq \frac{-\varphi_-}{1+e^{\varphi_-}} - \log(1 + e^{-\varphi_-}), \\ S_{\rho_-, \rho_-} \left(-(E + \bar{V}^-) \right), & \frac{-\varphi_-}{1+e^{\varphi_-}} - \log(1 + e^{-\varphi_-}) < E + \bar{V}^- \leq -\log(1 + e^{\varphi_-}) \end{cases} \end{aligned}$$

and is equal to $-\infty$ outside the energy band. Thus \tilde{S}^- coincides with S^- .

D.3. The case $\frac{1}{2} \leq \rho_- < \rho_+$:

In this case, by inverting the role of ρ_- with ρ_+ and of φ_- with φ_+ in the previous case, we obtain that \tilde{S}^- restricted to $[E_{-\infty}^-; E_{-\infty}^+]$ is given by

$$\begin{aligned} \tilde{S}^-(E) &= \\ &= \begin{cases} S_{\rho_-, \rho_-} \left(-(E + \bar{V}^-) \right), & -\log(1 + e^{\varphi_-}) \leq E + \bar{V}^- < \frac{-\varphi_-}{1+e^{\varphi_-}} - \log(1 + e^{-\varphi_-}), \\ -(E + \bar{V}^-), & \frac{-\varphi_-}{1+e^{\varphi_-}} - \log(1 + e^{-\varphi_-}) \leq E + \bar{V}^- \leq \frac{-\varphi_+}{1+e^{\varphi_+}} - \log(1 + e^{-\varphi_+}), \\ S_{\rho_+, \rho_+} \left(-(E + \bar{V}^-) \right), & \frac{-\varphi_+}{1+e^{\varphi_+}} - \log(1 + e^{-\varphi_+}) < E + \bar{V}^- \leq -\log(1 + e^{-\varphi_+}) \end{cases} \end{aligned}$$

and is equal to $-\infty$ outside the energy band. Thus \tilde{S}^- coincides with S^- .

Acknowledgments.

The authors are very grateful to Bernard Derrida for very useful discussions. They acknowledge the support of Égide (France) and FCT (Portugal) through the research project "Fluctuations of weakly and strongly asymmetric systems" no. FCT/1560/25/1/2012/S. CB acknowledges the support of the French Ministry of Education through the grants ANR-10-BLAN 0108 (SHEPI). PG thanks FCT for support through the research project PTDC/MAT/109844/2009 and the Research Centre of Mathematics of the University of Minho, for the financial support provided by "FEDER" through the "Programa Operacional Factores de Competitividade - COMPETE" and FCT through the research project PEst-C/MAT/UI0013/2011.

REFERENCES

- [1] Bahadoran, C., On the convergence of entropy for stationary exclusion processes with open boundaries, *J. Stat. Phys.* **126** (2007), no. 4-5, 1069–1082.
- [2] Bahadoran C., A quasi-potential for conservation laws with boundary conditions, (2010), eprint arXiv:1010.3624.
- [3] Bernardin C., Landim C., Entropy of stationary nonequilibrium measures of boundary driven symmetric simple exclusion processes. *J. Stat. Phys.* **141** (2010), no. 6, 1014–1038.
- [4] Derrida, B., Non-equilibrium steady states: fluctuations and large deviations of the density and of the current, *J. Stat. Mech.* P07023 (2007)
- [5] Derrida B., Enaud C. and Lebowitz J.L., The asymmetric exclusion process and Brownian excursions *J. Stat. Phys.* **115** (2004), 365–382.
- [6] Derrida, B., Lebowitz, J.L., Speer, E.R., Large Deviation of the Density Profile in the Steady State of the Open Symmetric Simple Exclusion, *J. Statist. Phys.* **107** (2002), no. 3-4, 599–634.
- [7] Derrida, B.; Lebowitz, J. L.; Speer, E. R.; Entropy of open lattice systems, *J. Stat. Phys.* **126** (2007), no. 4-5, 1083–1108.

- [8] Derrida, B., Lebowitz, J.L., Speer, E.R., Exact large deviation functional of a stationary open driven diffusive system: the asymmetric exclusion process *J. Stat. Phys.* **110** (2003), 775–810.
- [9] Kafri Y., Levine E., Mukamel D., Schütz G. M. and Torok J., Criterion for phase separation in one-dimensional driven systems (2002) *Phys. Rev. Lett.* **89** 035702
- [10] Touchette H., The large deviation approach to statistical mechanics, *Physics Reports* **478**, (2009), 1–69.

UNIVERSITÉ DE LYON AND CNRS, UMPA, UMR-CNRS 5669, ENS-LYON, 46, ALLÉE D'ITALIE,
69364 LYON CEDEX 07 - FRANCE.

E-MAIL: cedric.bernardin@ens-lyon.fr

CMAT, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO MINHO, CAMPUS DE GUALTAR, 4710-057
BRAGA, PORTUGAL.

E-MAIL: patg@math.uminho.pt

IMPA, ESTRADA DONA CASTORINA 110, CEP 22460 RIO DE JANEIRO, BRASIL AND CNRS UMR
6085, UNIVERSITÉ DE ROUEN, AVENUE DE L'UNIVERSITÉ, BP.12, TECHNOPÔLE DU MADRILLET, F76801
SAINT-ÉTIENNE-DU-ROUVRAY, FRANCE.

E-MAIL: landim@impa.br