

From Normal Diffusion to Superdiffusion of Energy in the Evanescent Flip Noise Limit

Cédric Bernardin¹ · Patrícia Gonçalves^{2,3} · Milton Jara⁴ · Makiko Sasada⁵ · Marielle Simon^{2,6}

Received: 23 September 2014 / Accepted: 5 March 2015
© Springer Science+Business Media New York 2015

Abstract We consider a harmonic chain perturbed by a stochastic noise which conserves the energy and a second quantity called the volume, and destroys all the other ones. We then add to this model a second energy conserving noise depending on a parameter γ , that annihilates the volume conservation. When γ is of order one, the energy diffuses according to the standard heat equation after a space-time diffusive scaling. On the other hand, when $\gamma = 0$, the energy superdiffuses according to a $3/4$ -fractional heat equation after a subdiffusive space-time scaling. In this paper, we study the existence of a crossover between these two regimes as a function of γ .

✉ Cédric Bernardin
cbernard@unice.fr

Patrícia Gonçalves
patg@math.uminho.pt; patricia@mat.puc-rio.br

Milton Jara
mjara@impa.br

Makiko Sasada
sasada@math.keio.ac.jp

Marielle Simon
marielle.simon@mat.puc-rio.br

- ¹ Laboratoire J.A. Dieudonné, UMR CNRS 7351, Université de Nice Sophia-Antipolis, Parc Valrose, 06108 Nice cedex 02, France
- ² Departamento de Matemática, PUC-RIO, Rua Marquês de São Vicente, No. 225, Rio de Janeiro, RJ 22453-900, Brazil
- ³ CMAT, Centro de Matemática da Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal
- ⁴ IMPA, Estrada Dona Castorina 110, Jardim Botânico, Rio de Janeiro CEP 22460-340, Brazil
- ⁵ Department of Mathematics, Keio University, 3-14-1, Hiyoshi, Kohoku-ku, Yokohama-shi, Kanagawa 223-8522, Japan
- ⁶ UMPA ENS de Lyon, 46 allée d'Italie, 69007 Lyon, France

Keywords Anomalous diffusion · Fourier's law · Equilibrium fluctuations

1 Introduction

Over the last few years, superdiffusion of energy in one-dimensional hamiltonian systems conserving momentum has attracted a lot of interest. In particular it is expected to hold for one-dimensional chains of oscillators when unpinned, and more generically for dynamical systems which preserve momentum in addition to the energy (we refer to the review papers [10, 17]). A proof of this behavior is still unreached even if some recent progresses have been accomplished [19].

In order to get rigorous results in this field, it has been proposed to perturb hamiltonian systems by a stochastic noise conserving energy and momentum. In [2] the thermal conductivity is proved to be infinite for an unpinned harmonic chain of oscillators perturbed by an energy-momentum conserving noise. When the stochastic perturbations of harmonic systems do not conserve momentum or if the chain is pinned (so that momentum is not conserved), the thermal conductivity is always finite [4, 8] and energy diffuses. The progresses in the anharmonic case are still modest (see [6] for the case of exponential interactions).

In this work we focus on the harmonic chain of oscillators perturbed by an energy conserving noise, as considered in [7]. After a simple symplectic change of variables, the dynamical state of the harmonic chain is given by $(\omega_x)_{x \in \mathbb{Z}} \in \mathbb{R}^{\mathbb{Z}}$, the energy is $\sum_{x \in \mathbb{Z}} \omega_x^2$ and Newton's equations write as

$$d\omega_x(t) = (\omega_{x+1}(t) - \omega_{x-1}(t))dt, \quad x \in \mathbb{Z}. \quad (1.1)$$

As explained in [7], one can see the variable ω_{2x} as the velocity of the atom x and the variable ω_{2x+1} as the interdistance particle between the atom x and the atom $x + 1$.

There are several ways to perturb the dynamics in order to conserve the energy. The *flip* noise changes ω_x into $-\omega_x$ independently on each site x at random exponential times with intensity $\gamma \geq 0$, and conserves the total energy. The *exchange* noise exchanges nearest neighbor values ω_x and ω_y at random exponential times with intensity $\lambda \geq 0$. Apart from the energy, this second perturbation conserves the so-called *total volume* of the chain $\sum_{x \in \mathbb{Z}} \omega_x$. The harmonic chain perturbed by the flip noise is diffusive [3, 18] while the harmonic chain perturbed by the exchange noise is superdiffusive [7]. This drastic difference is due to the volume conservation which plays a role similar to the momentum conservation in chains of oscillators (see [7] for more explanations).

In this work we consider the harmonic chain perturbed by the flip noise with intensity $\gamma \geq 0$ and the exchange noise with intensity $\lambda > 0$. If $\gamma = 0$, the volume is conserved, the energy transport is superdiffusive and described by a Levy process governed by a fractional Laplacian. This has been recently proved in [5]. If $\gamma > 0$, the volume is no longer conserved and one can prove that the energy transport is diffusive and described by a Brownian motion. Our aim is to study the case $\gamma \rightarrow 0$, with λ of order 1, and to obtain a crossover in a suitable time scale between these two very different regimes.

The decay of energy fluctuations is studied at spatial scale n , $n \in \mathbb{N}$ and temporal scale n^a , $a > 0$, assuming that the flip noise is given by $\gamma = cn^{-b}$, ($c, b \geq 0$). The exchange noise is taken to be always present, with $\lambda > 0$ independent of n , and one studies the limits of energy and volume correlation functions, as $n \rightarrow \infty$. In other words, the system consists of the harmonic chain with exchange noise, which is furthermore perturbed by the flip noise of strength $\gamma \ll 1$, and then inspected at spatial scale $\gamma^{-1/b}$ and temporal scale $\gamma^{-a/b}$.

When γ is sufficiently small, i.e. b sufficiently large, we follow the approach of [5] and show that the energy superdiffuses. This can be proved for $b > 1$ and the superdiffusion is described by the same Levy process as in the $\gamma = 0$ case. If γ is not sufficiently small, i.e. b is too small, the techniques of [5] do not straightforwardly apply and we use Varadhan’s approach which consists in decomposing the energy current $j_{x,x+1}$ between the atom x and the atom $x + 1$ as a sum of a discrete gradient term ∇f_x and a small fluctuating term. This decomposition is known in the mathematical literature as a *fluctuation–dissipation* equation and can be understood as a microscopic version of Fourier’s law. In our case, some new interesting features appear with respect to the models for which normal behavior is usually observed when writing the fluctuation–dissipation equation. Indeed, the function f_x involved is no longer local (see also [1] for similar features). We are able to write the fluctuation–dissipation equation for any value of $b \geq 0$ but we are only able to identify the limit of the energy fluctuation field for $b \in [0, 2/3)$. The case $b \in [2/3, 1]$ remains open.

2 Model and Notations

We consider an infinite chain of harmonic oscillators at equilibrium perturbed by an energy conserving noise. The space of configurations is given by $\Omega = \mathbb{R}^{\mathbb{Z}}$. We say that a function $f : \Omega \rightarrow \mathbb{R}$ is *local* if there exists a finite subset Λ of \mathbb{Z} such that the support of f is included in Λ . Let $n \geq 1$ be a scaling parameter. The Liouville operator corresponding to the harmonic chain (1.1) is given by

$$\mathcal{A} = \sum_{x \in \mathbb{Z}} (\omega_{x+1} - \omega_{x-1}) \partial_{\omega_x}.$$

The generator of the perturbed harmonic chain under investigation is given by

$$\mathcal{L}_n = \mathcal{A} + \gamma_n \mathcal{S}^{\text{flip}} + \lambda \mathcal{S}^{\text{exch}}$$

where for all smooth local bounded functions $f : \Omega \rightarrow \mathbb{R}$ we define

$$\begin{aligned} (\mathcal{S}^{\text{flip}} f)(\omega) &= \sum_{x \in \mathbb{Z}} (f(\omega^x) - f(\omega)), \\ (\mathcal{S}^{\text{exch}} f)(\omega) &= \sum_{x \in \mathbb{Z}} (f(\omega^{x,x+1}) - f(\omega)). \end{aligned}$$

Here, the configuration ω^x is the configuration obtained from ω by flipping the variable ω_x , i.e. $(\omega^x)_z = \omega_z$, if $z \neq x$, and $(\omega^x)_x = -\omega_x$. The configuration $\omega^{x,x+1}$ is obtained from ω by exchanging ω_x and ω_{x+1} , i.e. $(\omega^{x,x+1})_z = \omega_z$, if $z \neq x, x + 1$, and $(\omega^{x,x+1})_x = \omega_{x+1}$, $(\omega^{x,x+1})_{x+1} = \omega_x$. We denote by $\mathcal{S}_n := \gamma_n \mathcal{S}^{\text{flip}} + \lambda \mathcal{S}^{\text{exch}}$ the total generator of the perturbation, where $\gamma_n \geq 0, \lambda > 0$ are two parameters which regulate the respective strengths of the noises. We assume that

$$\gamma_n = \frac{c}{n^b}, \quad c, b \geq 0.$$

The process with generator \mathcal{L}_n is denoted by $\{\omega(t)\}_{t \geq 0} := \{\omega_x(t); x \in \mathbb{Z}\}_{t \geq 0}$.

The Gibbs equilibrium measures of $\{\omega(t)\}_{t \geq 0}$ are given by the Gaussian product probability measures

$$\mu_\beta(d\omega) = \prod_{x \in \mathbb{Z}} \sqrt{\frac{\beta}{2\pi}} \exp\left(-\frac{\beta \omega_x^2}{2}\right) d\omega_x,$$

where $\beta > 0$ stands for the inverse temperature. In the following, the expectation of a function f with respect to μ_β is denoted by $\langle f \rangle_\beta$ and the covariance between functions f, g with respect to μ_β is denoted by $\langle f; g \rangle_\beta$. We also denote by $\| \cdot \|_{\mathbf{L}^2(\mu_\beta)}$ the usual \mathbf{L}^2 -norm in the Hilbert space $\mathbf{L}^2(\mu_\beta)$.

We consider the dynamics starting from a Gibbs equilibrium measure at a fixed temperature β^{-1} in the time scale tn^a , $a > 0$. The existence of the infinite dynamics under this initial distribution can be proved by following for instance [16] or [12]. We denote by $\mathcal{S}(\mathbb{R})$ the Schwartz space of rapidly decreasing functions, and by $\mathcal{S}'(\mathbb{R})$ its topological dual, the space of tempered distributions.

Let us fix a time horizon line $T > 0$. We define the energy fluctuation field $\{\mathcal{E}_t^n ; t \in [0, T]\}$ in the time scale tn^a as the $\mathcal{S}'(\mathbb{R})$ -valued process given by

$$\mathcal{E}_t^n(f) = \frac{1}{\sqrt{n}} \sum_{x \in \mathbb{Z}} f\left(\frac{x}{n}\right) \left\{ \omega_x^2(tn^a) - \beta^{-1} \right\}.$$

Similarly, the volume fluctuation field $\{\mathcal{V}_t^n ; t \in [0, T]\}$ in the time scale tn^a is the $\mathcal{S}'(\mathbb{R})$ -valued process given by

$$\mathcal{V}_t^n(f) = \frac{1}{\sqrt{n}} \sum_{x \in \mathbb{Z}} f\left(\frac{x}{n}\right) \omega_x(tn^a).$$

It is not difficult to check that these two fields are well-defined almost surely.

3 Statement of the Results

Given two functions $f, g \in \mathcal{S}(\mathbb{R})$, we look at the time evolution of the space-time correlation function of the energy fluctuation field

$$\begin{aligned} \sigma_t^n(f, g) &= \langle \mathcal{E}_t^n(f); \mathcal{E}_0^n(g) \rangle_\beta \\ &= \frac{1}{n} \sum_{x, y \in \mathbb{Z}} f\left(\frac{x}{n}\right) g\left(\frac{y}{n}\right) \left\langle \left(\omega_x^2(tn^a) - \beta^{-1} \right) \left(\omega_y^2(0) - \beta^{-1} \right) \right\rangle_\beta \\ &= \frac{1}{n} \sum_{y, z \in \mathbb{Z}} f\left(\frac{y+z}{n}\right) g\left(\frac{y}{n}\right) \langle \omega_z^2(tn^a) (\omega_0^2(0) - \beta^{-1}) \rangle_\beta, \end{aligned}$$

when $n \rightarrow \infty$. The choice of $a > 0$ fixes the time scale.

Theorem 3.1 *Assume $c > 0$ and $b < 2/3$, and set $a = 2 - b/2$. Let us fix $t > 0$. Then, for $f, g \in \mathcal{S}(\mathbb{R})$*

$$\lim_{n \rightarrow \infty} \sigma_t^n(f, g) = \frac{1}{\beta^2 \sqrt{\pi t \kappa}} \iint_{\mathbb{R}^2} dudv f(u) g(v) \exp\left(-\frac{(u-v)^2}{4t\kappa}\right),$$

where

$$\kappa = \begin{cases} \frac{1}{\sqrt{2\lambda c}} & \text{if } b \in (0, 2/3), \\ \lambda + \frac{1}{\sqrt{2\lambda c}} & \text{if } b = 0. \end{cases}$$

One can also prove that the fluctuation field \mathcal{E}_t^n converges in law to the infinite dimensional Ornstein–Uhlenbeck (OU) process \mathcal{E}_t , solution to the linear stochastic partial differential equation

$$\partial_t \mathcal{E} = \kappa \partial_u^2 \mathcal{E} dt + \sqrt{4\kappa\beta^{-2}} \partial_u B(u, t),$$

where B is the standard normalized space–time white noise. This extension of Theorem 3.1 is standard, and we refer to [14] for more details.

For the sake of simplicity, hereafter we assume $\lambda = 1$. The same computations for any $\lambda > 0$ could be done, but become significantly more technical. In Sect. 5 we prove the following theorem.

Theorem 3.2 *Let $\{P_t\}_{t \geq 0}$ be the semigroup generated by the infinitesimal generator*

$$\mathbb{L} := -\frac{1}{\sqrt{2}} \left((-\Delta)^{3/4} - \nabla(-\Delta)^{1/4} \right). \tag{3.1}$$

We assume $a = 3/2$. If $c = 0$ or $b > 1$, then, for $f, g \in \mathcal{S}(\mathbb{R})$,

$$\lim_{n \rightarrow +\infty} \sigma_t^n(f, g) = \frac{2}{\beta^2} \iint_{\mathbb{R}^2} dudv f(u)g(v)P_t(u - v). \tag{3.2}$$

As for Theorem 3.1, with a little more effort we could also prove that the energy fluctuation field \mathcal{E}_t^n converges to an infinite dimensional 3/4-fractional Ornstein–Uhlenbeck (fOU) process.

The case $b \in [2/3, 1]$ remains open. One possible behavior is the following: $b = 1$ would be a field interpolating the standard OU process and the fOU process, and $b \in [2/3, 1)$ would correspond to the same diffusive behavior as for $b \in [0, 2/3)$. In any cases, it is easy, by scaling considerations, to see that the limiting energy field $\mathcal{E}(t, x; c, b) := \lim_{n \rightarrow \infty} \mathcal{E}_t^n$ in the time scale tn^a shall satisfy the scaling relation (in law)

$$\mathcal{E}(t, x; c, b) = \varepsilon^{1/2} \mathcal{E}(t\varepsilon^a, \varepsilon x; c\varepsilon^{-b}, b), \quad \varepsilon > 0. \tag{3.3}$$

The only result we are able to prove in the window $b \in [2/3, 1]$ is that the time scale necessary to see some macroscopic evolution of the energy is at least $tn^{4/3}$. The proof is the same as in [6] and we refer the interested reader to that paper.

We turn now to the volume fluctuation field. The space–time correlation function of the volume fluctuation field in the time-scale tn^a is defined for $f, g \in \mathcal{S}(\mathbb{R})$ as

$$\begin{aligned} \eta_t^n(f, g) &:= \frac{1}{n} \sum_{x, y \in \mathbb{Z}} f\left(\frac{x}{n}\right) g\left(\frac{y}{n}\right) \langle \omega_x(tn^a) \omega_y(0) \rangle_\beta \\ &= \frac{1}{n} \sum_{z, y \in \mathbb{Z}} f\left(\frac{y+z}{n}\right) g\left(\frac{y}{n}\right) \langle \omega_z(tn^a) \omega_0(0) \rangle_\beta. \end{aligned} \tag{3.4}$$

We obtain complete results, in the sense that all time scales a and all exponents b are covered. The behavior of the volume fluctuations is of three different types:

- *Relaxation* this means that η_t^n converges to η_t which is solution to

$$\partial_t \eta_t(f, g) = -C \eta_t(f, g),$$

for some constant $C > 0$.

- *Transport* this means that η_t^n converges to η_t which is solution to

$$\partial_t \eta_t(f, g) = C \eta_t(f', g),$$

for some constant $C > 0$.

- *Heat* this means that η_t^n converges to η_t which is solution to

$$\partial_t \eta_t(f, g) = C \eta_t(f'', g),$$

for some constant $C > 0$.

Moreover, for any parameter $b > 0$, we will see that the hyperbolic time scale $a = 1$ yields to a transport equation whose speed is constantly equal to 2. In other words, the limit volume fluctuation field at time t is a translation of the initial one. As a consequence, in higher time scales, the fluctuation field should be redefined in order to take into account this transport phenomenon. When the time scale a satisfies $a > 1$, we redefine the space-time correlation function of the volume fluctuation field as

$$\tilde{\eta}_t^n(f, g) = \frac{1}{n} \sum_{x, y \in \mathbb{Z}} f\left(\frac{x-2tn^a}{n}\right) g\left(\frac{y}{n}\right) \langle \omega_x(tn^a) \omega_y(0) \rangle_\beta$$

Let us remark that the two fluctuation fields initially coincide: $\eta_0^n = \tilde{\eta}_0^n$.

Theorem 3.3 *The behavior of the volume fluctuation field depends on the time scale in the following way:*

(A) *Case $b \leq 1$.*

- (i) *If $a < b$, then $\partial_t \eta_t^n(f, g)$ vanishes, and*

$$\lim_{n \rightarrow \infty} \eta_t^n(f, g) = \eta_0(f, g) = \beta^{-1} \iint_{\mathbb{R}^2} dudv f(u)g(v).$$

- (ii) *If $a = b$, then*

$$\lim_{n \rightarrow \infty} \eta_t^n(f, g) = \iint_{\mathbb{R}^2} dudv f(u)g(v)P_t(u - v),$$

where $\{P_t\}_{t \geq 0}$ is the semigroup generated by the infinitesimal generator

$$\mathbf{1}_{b=1} \times 2\nabla - 2c\text{Id}.$$

- (iii) *If $a > b$, $\lim_{n \rightarrow \infty} \eta_t^n(f, g) = 0$.*

(B) *Case $1 < b < 2$.*

- (i) *If $a < b$ and $a \in (0, 1)$, then $\partial_t \eta_t^n(f, g)$ vanishes, and*

$$\lim_{n \rightarrow \infty} \eta_t^n(f, g) = \eta_0(f, g) = \beta^{-1} \iint_{\mathbb{R}^2} dudv f(u)g(v).$$

- (ii) *If $a < b$ and $a \in (1, 2)$, then $\partial_t \tilde{\eta}_t^n(f, g)$ vanishes, and*

$$\lim_{n \rightarrow \infty} \tilde{\eta}_t^n(f, g) = \eta_0(f, g) = \beta^{-1} \iint_{\mathbb{R}^2} dudv f(u)g(v).$$

- (iii) *If $a = 1$,*

$$\lim_{n \rightarrow \infty} \eta_t^n(f, g) = \iint_{\mathbb{R}^2} dudv f(u)g(v)P_t(u - v),$$

where $\{P_t\}_{t \geq 0}$ is the semigroup generated by the infinitesimal generator 2∇ .

(iv) If $a = b$, then $\tilde{\eta}_t^n$ converges to η_t where η_t is the solution of

$$\begin{cases} \partial_t \eta_t(f, g) = -2c\eta_t(f, g), \\ \eta_0(f, g) = \beta^{-1} \iint_{\mathbb{R}^2} dudv f(u)g(v). \end{cases}$$

(v) If $a > b$, $\lim_{n \rightarrow \infty} \eta_t^n(f, g) = 0$.

(C) Case $b \geq 2$.

(i) If $a \in (0, 1)$, then $\partial_t \eta_t^n(f, g)$ vanishes, and

$$\lim_{n \rightarrow \infty} \eta_t^n(f, g) = \eta_0(f, g) = \beta^{-1} \iint_{\mathbb{R}^2} dudv f(u)g(v).$$

(ii) If $a \in (1, 2)$, then $\partial_t \tilde{\eta}_t^n(f, g)$ vanishes, and

$$\lim_{n \rightarrow \infty} \tilde{\eta}_t^n(f, g) = \eta_0(f, g) = \beta^{-1} \iint_{\mathbb{R}^2} dudv f(u)g(v).$$

(iii) If $a = 1$,

$$\lim_{n \rightarrow \infty} \eta_t^n(f, g) = \iint_{\mathbb{R}^2} dudv f(u)g(v)P_t(u - v),$$

where $\{P_t\}_{t \geq 0}$ is the semigroup generated by the infinitesimal generator 2∇ .

(iv) If $a = 2$,

$$\lim_{n \rightarrow \infty} \tilde{\eta}_t^n(f, g) = \iint_{\mathbb{R}^2} dudv f(u)g(v)P_t(u - v),$$

where $\{P_t\}_{t \geq 0}$ is the semigroup generated by the infinitesimal generator

$$\lambda \Delta - \mathbf{1}_{b=2} \times 2c\text{Id}.$$

(v) If $a > 2$, $\lim_{n \rightarrow \infty} \eta_t^n(f, g) = 0$.

We summarize in Figs. 1 and 2 all the results of this paper.

4 Diffusive Domain for the Energy

Before giving the proof of Theorem 3.1 we recall some facts about the notion of the fluctuation–dissipation equation. Hereafter, τ_x denotes the translated operator that acts on a function $h : \Omega \rightarrow \mathbb{R}$ as $(\tau_x h)(\omega) := h(\tau_x \omega)$, and $\tau_x \omega$ is the configuration obtained from ω by shifting: $(\tau_x \omega)_y = \omega_{x+y}$. The microscopic current of energy is defined by the local conservation law $\mathcal{L}_n(\omega_x^2) = \nabla(j_{x-1,x})$, where the discrete gradient ∇ is defined for a function $h : \Omega \rightarrow \mathbb{R}$ as

$$(\nabla h)(\omega) := (\tau_1 h)(\omega) - h(\omega).$$

Here, the energy current is explicitly given by

$$j_{x,x+1}(\omega) = j_{x,x+1}^A(\omega) + j_{x,x+1}^S(\omega) = 2\omega_x \omega_{x+1} + \lambda (\omega_{x+1}^2 - \omega_x^2). \tag{4.1}$$

Let us notice that the current does not depend on n (since it does not involve the intensity γ_n of the flip noise). Recall that we are interested in the energy transport in the limit $\gamma \rightarrow 0$. More exactly, we assume that $\gamma = \frac{c}{n^b}$ with $n \rightarrow \infty$.




Fig. 1 Energy fluctuations




Fig. 2 Volume fluctuations

If $b = 0$, i.e. $\gamma_n = c$ is a constant independent of n , and $\lambda = 0$, the volume is not conserved and we can show that the energy fluctuation field \mathcal{E}_t^n converges as $n \rightarrow \infty$, when $a = 2$, towards an infinite dimensional Ornstein–Uhlenbeck process: the energy transport is diffusive. The argument is based on two ingredients: (1) the existence of a fluctuation–dissipation equation and (2) the Boltzmann–Gibbs principle. A fluctuation–dissipation equation for the

current is the decomposition of the current as a sum of a discrete gradient ∇f_x and a term in the form $\mathcal{L}_n(g_x)$, where f_x and g_x are two functions of the configuration. The *Boltzmann–Gibbs principle* was first introduced by Brox and Rost [9] and, roughly speaking, states that the space-time fluctuations of any non-conserved field can be written as a linear functional of the conserved field plus a small fluctuating term. These two properties are sufficient to close the evolution equation of the energy fluctuation field (see [14] for more details).

In the case $b = \lambda = 0$, a straightforward computation shows that

$$j_{x,x+1} = \nabla (\omega_x^2 + \omega_{x-1}\omega_{x+1}) - \mathcal{L}_n (\omega_x\omega_{x+1}),$$

where ∇ is the discrete gradient. Thus an exact fluctuation–dissipation equation involving local functions holds¹. If $b = 0$ but $\lambda > 0$, then a fluctuation–dissipation equation still holds, but the functions involved in the decomposition are no longer local. However, they are exponentially localized so that this does not really matter and a normal diffusion of energy takes place (see e.g. [1]). On the other hand, if $b = \infty$ (or equivalently $c = 0$) and $\lambda > 0$, the situation is very different since energy superdiffuses [5]. In this case, it is not clear if such a fluctuation–dissipation equation can be obtained: if it exists, it should involve very non-local functions.

It turns out that if $b < \infty$, an exact fluctuation–dissipation equation still holds with exponentially localized functions. Nevertheless the length scale of the localization is typically of order γ_n^{-1} which goes to ∞ , as $n \rightarrow \infty$. The “standard technique” [14] to obtain convergence of the fluctuation field to an Ornstein–Uhlenbeck process has to be adapted with care and, in fact, we are only able to do it for $b < 2/3$.

Along the paper we shall use letters C, C', C_0, \dots to denote constants that do not depend on n . The value of the constant can change from line to line.

4.1 The Fluctuation–Dissipation Equation

Strategy In the sequel, we erase the dependence on the parameters n, b and c whenever no confusion arises. We consider a function u in the form

$$u(\omega) = \sum_{x \in \mathbb{Z}} \sum_{k \geq 1} \rho_k(x) \omega_x \omega_{x+k}, \tag{4.2}$$

where $\{\rho_k(x); x \in \mathbb{Z}, k \geq 1\}$ is a real sequence that satisfies the condition

$$\sum_{x \in \mathbb{Z}} \sum_{k \geq 1} |\rho_k(x)|^2 < +\infty, \tag{4.3}$$

so that u is a function in $L^2(\mu_\beta)$. Observe first that $\mathcal{A}u$ is a sum of gradient terms. Indeed, we have

$$\mathcal{A}u(\omega) = \sum_{x \in \mathbb{Z}} \sum_{k \geq 1} \rho_k(x) \nabla [\omega_{x-1}\omega_{x+k} + \omega_x\omega_{x+k-1}]. \tag{4.4}$$

Our aim is now to solve the equation

$$(\gamma_n \mathcal{S}^{\text{flip}} + \lambda \mathcal{S}^{\text{exch}})(u)(\omega) = 2\omega_0\omega_1 = j_{0,1}^A(\omega). \tag{4.5}$$

Then it will follow that

$$\mathcal{L}_n u - v = j_{0,1}^A \tag{4.6}$$

¹ The situation is exactly similar to the velocity-flip model investigated in [18] and the results of [18] can be adapted to this model mutatis mutandis.

where $v = \mathcal{A}u$ given in (4.4) is a sum of gradient functions. One can easily check that v and $\mathcal{L}_n u$ are also in the space $\mathbf{L}^2(\mu_\beta)$.

Resolution Straightforward computations show that

$$(\gamma_n \mathcal{S}^{\text{flip}} + \lambda \mathcal{S}^{\text{exch}})(u)(\omega) = \sum_{x \in \mathbb{Z}} \sum_{k \geq 1} F_k(x) \omega_x \omega_{x+k},$$

with

$$\begin{aligned} F_1(x) &= -2(2\gamma_n + \lambda) \rho_1(x) + \lambda(\rho_2(x) + \rho_2(x - 1)), \\ F_k(x) &= -4(\gamma_n + \lambda) \rho_k(x) + \lambda(\rho_{k-1}(x) + \rho_{k-1}(x + 1) + \rho_{k+1}(x) \\ &\quad + \rho_{k+1}(x - 1)), \quad k \geq 2. \end{aligned}$$

For the sake of clarity, we forget that the coefficients $F_k(x)$ should depend on n . After identifying the coefficients in front of the different terms, it follows that (4.5) holds if, for all $x \in \mathbb{Z}$,

$$F_k(x) = 2\mathbf{1}_{\{k=1, x=0\}}. \tag{4.7}$$

In Appendix 7.2 we recall the definition of the bijective Fourier transform $\widehat{h} \in \mathbf{L}^2(\mathbb{T})$ of a given function $h \in \ell^2(\mathbb{Z}, \mathbb{R})$. Hereafter \mathbb{T} denotes the one-dimensional torus $[0, 1)$. The identity (4.7) can be equivalently reformulated in terms of the Fourier transform as follows: for $\theta \in \mathbb{T}$,

$$\begin{cases} -2(2\gamma_n + \lambda) \widehat{\rho}_1(\theta) + \lambda(1 + e^{2i\pi\theta}) \widehat{\rho}_2(\theta) = 2, \\ -4(\gamma_n + \lambda) \widehat{\rho}_k(\theta) + \lambda(1 + e^{-2i\pi\theta}) \widehat{\rho}_{k-1}(\theta) + \lambda(1 + e^{2i\pi\theta}) \widehat{\rho}_{k+1}(\theta) = 0, \quad k \geq 2. \end{cases} \tag{4.8}$$

By the Parseval–Plancherel’s identity, the condition (4.3) is equivalent to

$$\sum_{k \geq 1} \int_{\mathbb{T}} |\widehat{\rho}_k(\theta)|^2 d\theta < +\infty.$$

Then, one can easily show that (4.8) and the above integrability condition lead to

$$\widehat{\rho}_k(\theta) = \widehat{\rho}_1(\theta)(X(\theta))^{k-1},$$

with

$$X(\theta) = \frac{2}{1 + e^{2i\pi\theta}} \left\{ 1 + \frac{\gamma_n}{\lambda} - \sqrt{\left(1 + \frac{\gamma_n}{\lambda}\right)^2 - \cos^2(\pi\theta)} \right\}$$

and

$$\widehat{\rho}_1(\theta) = -\frac{1}{\gamma_n + \lambda \sqrt{\left(1 + \frac{\gamma_n}{\lambda}\right)^2 - \cos^2(\pi\theta)}}.$$

Sharp Estimates In the following, we will need sharp estimates on $\widehat{\rho}_1$ and X , precisely:

Lemma 4.1 *For sufficiently large n , and for any $\theta \in \mathbb{T}$, we have*

$$|X(\theta)| \leq \frac{1}{1 + \sqrt{\frac{\gamma_n}{\lambda}}} \quad \text{and} \quad |\widehat{\rho}_1(\theta)| \leq \frac{1}{\lambda \sqrt{\frac{\gamma_n}{\lambda} + \sin^2(\pi\theta)}}.$$

Proof We only prove the first estimate, since the second one is straightforward. Let us define $\mathbf{c} := \gamma_n/\lambda$. Then,

$$|X(\theta)| = \varphi\left(\frac{|\cos(\pi\theta)|}{1 + \mathbf{c}}\right),$$

where φ is the function defined by $\varphi(y) = (1 - \sqrt{1 - y^2})/y$. Since φ is increasing on $[0, 1/(1 + \mathbf{c})]$ we get

$$|X(\theta)| \leq \varphi\left(\frac{1}{1 + \mathbf{c}}\right).$$

Observing that $(1 + \sqrt{\mathbf{c}})\varphi(1/(1 + \mathbf{c})) = 1 - (\sqrt{2} - 1)\sqrt{\mathbf{c}} + O(\mathbf{c})$ whenever $\mathbf{c} \rightarrow 0$. Therefore, for n large enough, we have

$$\forall \theta \in \mathbb{T}, \quad |X(\theta)| \leq \frac{1}{1 + \sqrt{\mathbf{c}}}.$$

□

4.2 Strategy of the Proof of Theorem 3.1

We have

$$\sigma_t^n(f, g) = \sigma_0^n(f, g) - n^{a-2} \int_0^t \sum_{y,z \in \mathbb{Z}} (\nabla_n f)\left(\frac{y+z}{n}\right) g\left(\frac{y}{n}\right) \left\langle j_{z,z+1}(sn^a) (\omega_0^2(0) - \beta^{-1}) \right\rangle_\beta ds,$$

where ∇_n is the discretization of the derivative w.r.t. the lattice $n^{-1}\mathbb{Z}$, that is,

$$(\nabla_n f)\left(\frac{y}{n}\right) := n \left[f\left(\frac{y+1}{n}\right) - f\left(\frac{y}{n}\right) \right].$$

The discretization Δ_n of the Laplacian is defined in a similar way. Let us recall from (4.1) that the current writes as $j_{z,z+1} = j_{z,z+1}^A + j_{z,z+1}^S$.

If $a < 2$, it is easy to see that the contribution coming from $j_{z,z+1}^S$ vanishes as $n \rightarrow \infty$, since a second integration by parts can be performed and the time scale is subdiffusive. If $a = 2$ and $b = 0$, the symmetric part of the current gives a non trivial contribution. More precisely, after an integration by parts we get the term

$$\int_0^t \sigma_s^n(\lambda f'', g) ds,$$

and the coefficient λ appears in the thermal conductivity.

We now assume $a < 2$. By using the fluctuation–dissipation equation (4.6) for the contribution coming from $j_{0,1}^A$, we obtain

$$\begin{aligned} \sigma_t^n(f, g) &= \sigma_0^n(f, g) + n^{a-2} \int_0^t \sum_{y,z \in \mathbb{Z}} (\nabla_n f)\left(\frac{y+z}{n}\right) g\left(\frac{y}{n}\right) \left\langle (\tau_z v)(sn^a) (\omega_0^2(0) - \beta^{-1}) \right\rangle_\beta ds \\ &\quad - n^{a-2} \int_0^t \sum_{y,z \in \mathbb{Z}} (\nabla_n f)\left(\frac{y+z}{n}\right) g\left(\frac{y}{n}\right) \left\langle (\mathcal{L}_n \tau_z u)(sn^a) (\omega_0^2(0) - \beta^{-1}) \right\rangle_\beta ds \\ &\quad + o_n(1) \\ &=: \sigma_0^n(f, g) + V_t^n(f, g) - U_t^n(f, g) + o_n(1), \end{aligned}$$

where $o_n(1)$ vanishes as $n \rightarrow \infty$. Let us now focus on the term $V_t^n(f, g)$. The function v can be rewritten as

$$v(\omega) = \sum_{x \in \mathbb{Z}} \rho_1(x) \nabla(\omega_x^2) + \psi(\omega)$$

where

$$\psi(\omega) = \sum_{x \in \mathbb{Z}} \sum_{k \geq 2} \rho_k(x) \nabla \{ \omega_{x-1} \omega_{x+k} + \omega_x \omega_{x+k-1} \} + \sum_{x \in \mathbb{Z}} \rho_1(x) \{ \omega_x \omega_{x+2} - \omega_{x-1} \omega_{x+1} \}. \tag{4.9}$$

Then, accordingly to this decomposition, we write the term $V_t^n(f, g)$ as the sum of two terms

$$V_t^n(f, g) = K_t^n(f, g) + \Psi_t^n(f, g).$$

It turns out that

$$K_t^n(f, g) = -n^{a-1} \int_0^t \sigma_s^n(F, g) ds,$$

where the function F is defined on $\frac{1}{n}\mathbb{Z}$ by

$$\begin{aligned} F\left(\frac{w}{n}\right) &= \sum_{z \in \mathbb{Z}} \rho_1(z) \left[(\nabla_n f)\left(\frac{w-z}{n}\right) - (\nabla_n f)\left(\frac{w-z-1}{n}\right) \right] \\ &= \frac{1}{n} \sum_{z \in \mathbb{Z}} \rho_1(z) (\Delta_n f)\left(\frac{w-z}{n}\right), \quad w \in \mathbb{Z}. \end{aligned}$$

In the sequel we prove the following convergences:

- (i) If $b < 1$, then $\lim_{n \rightarrow +\infty} |U_t^n(f, g)| = 0$.
- (ii) If $a = 2 - b/2$ and $b < 2$, then $\lim_{n \rightarrow \infty} \left| K_t^n(f, g) - \int_0^t \sigma_s^n(\kappa f'', g) ds \right| = 0$.
- (iii) If $b < 2/3$, then $\lim_{n \rightarrow \infty} |\Psi_t^n(f, g)| = 0$.

One can easily check that these three points imply Theorem 3.1. Besides, we shall see in the proof of (ii) that the case $a < 2 - b/2$ corresponds to $|V_t^n(f, h)| \rightarrow 0$, as $n \rightarrow \infty$. In other words, there is no evolution up to the time scale $a = 2 - b/2$.

Remark 4.2 If $b > 2$ and $a = 3/2$, we can adapt the argument in the proof of (ii) and show that the limit results in a constant times the 3/4-fractional Laplacian of f (instead of a constant times the second derivative of f). However, this is not sufficient to prove that for $b > 2$ the limit of the energy fluctuation field is given by a fractional heat equation, because we do not know how to control the other terms (U_t^n and Ψ_t^n) for $b > 2$. And in fact we know from the results of Sect. 5 that the contribution of these terms is not trivial, since a drift term should appear.

4.3 Proofs of Convergence and Boltzmann–Gibbs Principle

In this section we prove the above three convergence results.

Lemma 4.3 (Fluctuation part) *If $b < 1$, then*

$$\lim_{n \rightarrow +\infty} |U_t^n(f, g)| = 0.$$

Proof For each $z \in \mathbb{Z}$ we have that

$$n^a \int_0^t (\mathcal{L}_n \tau_z u)(sn^a) ds = (\tau_z u)(tn^a) - (\tau_z u)(0) + \mathcal{N}_t^{z,n},$$

where $\mathcal{N}^{z,n}$ is a martingale equal to 0 at $t = 0$ so that for any $t \geq 0$,

$$\langle \mathcal{N}_t^{z,n} (\omega_0^2(0) - \beta^{-1}) \rangle_\beta = 0.$$

Notice that the expression for u only involves terms of the form $\omega_x \omega_y$ with $x \neq y$, so that $\langle \tau_z u(0) (\omega_0^2(0) - \beta^{-1}) \rangle_\beta = 0$. We use the translation invariance of the dynamics in order to write

$$\begin{aligned} U_t^n(f, h) &= \frac{1}{n^2} \sum_{y,z \in \mathbb{Z}} (\nabla_n f) \left(\frac{y+z}{n} \right) g \left(\frac{y}{n} \right) \left\langle [(\tau_z u)(tn^a) - (\tau_z u)(0)] (\omega_0^2(0) - \beta^{-1}) \right\rangle_\beta \\ &= \frac{1}{n} \left\langle \left[\frac{1}{\sqrt{n}} \sum_{y \in \mathbb{Z}} g \left(\frac{y}{n} \right) (\omega_y^2(0) - \beta^{-1}) \right] \left[\frac{1}{\sqrt{n}} \sum_{z \in \mathbb{Z}} (\nabla_n f) \left(\frac{z}{n} \right) (\tau_z u)(tn^a) \right] \right\rangle_\beta. \end{aligned}$$

Then, by using the Cauchy–Schwarz inequality and the stationarity of μ_β , there exist constants $C', C > 0$ independent of n such that

$$\begin{aligned} |U_t^n(f, g)| &\leq \frac{C'}{n} \left\| \frac{1}{\sqrt{n}} \sum_{y \in \mathbb{Z}} g \left(\frac{y}{n} \right) (\omega_y^2(0) - \beta^{-1}) \right\|_{\mathbf{L}^2(\mu_\beta)} \left\| \frac{1}{\sqrt{n}} \sum_{z \in \mathbb{Z}} (\nabla_n f) \left(\frac{z}{n} \right) \tau_z u \right\|_{\mathbf{L}^2(\mu_\beta)} \\ &\leq \frac{C}{n} \left\| \frac{1}{\sqrt{n}} \sum_{z \in \mathbb{Z}} (\nabla_n f) \left(\frac{z}{n} \right) \tau_z u \right\|_{\mathbf{L}^2(\mu_\beta)}. \end{aligned}$$

Using (4.2) and the Parseval–Plancherel’s relation, a simple computation shows that

$$\begin{aligned} &\left\| \frac{1}{\sqrt{n}} \sum_{z \in \mathbb{Z}} (\nabla_n f) \left(\frac{z}{n} \right) \tau_z u \right\|_{\mathbf{L}^2(\mu_\beta)}^2 \\ &= \frac{1}{\beta^2 n} \sum_{k \geq 1} \sum_{y,x \in \mathbb{Z}} (\nabla_n f) \left(\frac{y}{n} \right) (\nabla_n f) \left(\frac{y-x}{n} \right) \left\{ \sum_{z \in \mathbb{Z}} \rho_k(z) \rho_k(z-x) \right\} \\ &= \frac{1}{\beta^2 n} \sum_{k \geq 1} \int_{\mathbb{T}} |\widehat{\rho}_k(\theta)|^2 \left[\sum_{y,x \in \mathbb{Z}} e^{-2i\pi\theta x} (\nabla_n f) \left(\frac{y}{n} \right) (\nabla_n f) \left(\frac{y-x}{n} \right) \right] d\theta \\ &= \frac{n}{\beta^2} \sum_{k \geq 1} \int_{\mathbb{T}} |\widehat{\rho}_k(\theta)|^2 |\mathcal{F}_n(\nabla_n f)|^2(n\theta) d\theta \\ &= \frac{n}{\beta^2} \int_{\mathbb{T}} \frac{|\widehat{\rho}_1(\theta)|^2}{1 - |X(\theta)|^2} |\mathcal{F}_n(\nabla_n f)|^2(n\theta) d\theta. \end{aligned}$$

In Appendix 1, the Fourier transform of a function $g : \frac{1}{n}\mathbb{Z} \rightarrow \mathbb{R}$ is defined by

$$\mathcal{F}_n(g)(\xi) = \frac{1}{n} \sum_{x \in \mathbb{Z}} g \left(\frac{x}{n} \right) e^{2i\pi\xi x/n}, \quad \xi \in \mathbb{R}. \tag{4.10}$$

According to Lemma 4.1, there exists a constant $C > 0$ such that

$$\frac{|\widehat{\rho}_1(\theta)|^2}{1 - |X(\theta)|^2} \leq \frac{C}{\sqrt{\gamma_n} \left[\frac{\gamma_n}{\lambda} + \sin^2(\pi\theta) \right]}.$$

By Lemma 7.1, $|\mathcal{F}_n(\nabla_n f)|^2(n\theta)$ is bounded above by a constant $C > 0$ independent of n and θ , from where we deduce that

$$\left\| \frac{1}{\sqrt{n}} \sum_{z \in \mathbb{Z}} (\nabla_n f)\left(\frac{z}{n}\right) \tau_z u \right\|_{L^2(\mu_\beta)}^2 \leq \frac{Cn}{\sqrt{\gamma_n}} \int_{\mathbb{T}} \frac{1}{\left[\frac{\gamma_n}{\lambda} + \sin^2(\pi\theta)\right]} d\theta.$$

The right-hand side (RHS) of the last inequality is of order n/γ_n so that the lemma follows as soon as $\sqrt{n\gamma_n}$ diverges to ∞ , which is equivalent to the condition $b < 1$. \square

Now we deal with the term $K_t^n(f, g)$.

Lemma 4.4 (Diffusive behavior) *If $a = 2 - b/2$ and $b < 2$, then*

$$\lim_{n \rightarrow \infty} \left| K_t^n(f, g) - \int_0^t \sigma_s^n(\kappa f'', g) ds \right| = 0.$$

Proof First, let us write

$$\begin{aligned} & K_t^n(f, g) - \int_0^t \sigma_s^n(\kappa f'', g) ds \\ &= \int_0^t \frac{1}{n} \sum_{x, y \in \mathbb{Z}} \left[-n^{a-1} F\left(\frac{x}{n}\right) - \kappa f''\left(\frac{x}{n}\right) \right] g\left(\frac{y}{n}\right) \left\langle (\omega_x^2(s n^a) - \beta^{-1})(\omega_y^2(0) - \beta^{-1}) \right\rangle_\beta ds. \end{aligned}$$

Then, from the Cauchy–Schwarz inequality, there exists a constant $C > 0$ such that

$$\left| K_t^n(f, g) - \int_0^t \sigma_s^n(\kappa f'', g) ds \right|^2 \leq \frac{C}{n} \sum_{w \in \mathbb{Z}} \left[(n^{a-1} F)\left(\frac{w}{n}\right) + \kappa (f'')\left(\frac{w}{n}\right) \right]^2.$$

We are reduced to prove that the RHS vanishes, as $n \rightarrow \infty$. The proof relies on the Fourier transform. The discrete Fourier transform of $n^{a-1} F$ is given by

$$\begin{aligned} n^{a-1} \mathcal{F}_n(F)(\xi) &= n^{a-2} \sum_{y \in \mathbb{Z}} F\left(\frac{y}{n}\right) e^{2i\pi\xi y/n} = n^{a-2} \mathcal{F}_n(\Delta_n f)(\xi) \widehat{\rho}_1\left(\frac{\xi}{n}\right) \\ &= -n^a \frac{4 \sin^2\left(\pi \frac{\xi}{n}\right)}{\gamma_n + \lambda \sqrt{\left(1 + \frac{\gamma_n}{\lambda}\right)^2 - \cos^2\left(\pi \frac{\xi}{n}\right)}} \mathcal{F}_n(f)(\xi), \quad \xi \in \mathbb{R}. \end{aligned}$$

We denote

$$q_n\left(\frac{\xi}{n}\right) := -n^a \frac{4 \sin^2\left(\pi \frac{\xi}{n}\right)}{\gamma_n + \lambda \sqrt{\left(1 + \frac{\gamma_n}{\lambda}\right)^2 - \cos^2\left(\pi \frac{\xi}{n}\right)}}.$$

Recall that $\kappa = \frac{1}{\sqrt{2\lambda c}}$. By the Parseval–Plancherel’s relation it is equivalent to prove

$$\lim_{n \rightarrow \infty} I_n := \lim_{n \rightarrow \infty} \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi) - \frac{1}{\sqrt{2\lambda c}} \mathcal{F}_n(f'')(\xi) \right|^2 d\xi = 0.$$

Since

$$\lim_{n \rightarrow \infty} \int_{[-\frac{n}{2}, \frac{n}{2}]} |\mathcal{F}_n(g)(\xi) - (\mathcal{F}g)(\xi)|^2 d\xi = 0$$

for any $g \in \mathcal{S}(\mathbb{R})$, we can replace in I_n the term $\mathcal{F}_n(f'')(\xi)$ by

$$\mathcal{F}(f'')(\xi) = -4\pi^2|\xi|^2(\mathcal{F}f)(\xi),$$

where \mathcal{F} is the usual Fourier transform defined on $\mathcal{S}(\mathbb{R})$. Then we bound

$$\begin{aligned} & \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right)\mathcal{F}_n(f)(\xi) + \frac{4\pi^2\xi^2}{\sqrt{2\lambda c}}(\mathcal{F}f)(\xi) \right|^2 d\xi \\ & \leq 2 \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) + \frac{4\pi^2\xi^2}{\sqrt{2\lambda c}} \right|^2 |\mathcal{F}_n(f)(\xi)|^2 d\xi \\ & \quad + \frac{16\pi^4}{\lambda c} \int_{[-\frac{n}{2}, \frac{n}{2}]} \xi^4 |\mathcal{F}_n(f)(\xi) - (\mathcal{F}f)(\xi)|^2 d\xi. \end{aligned}$$

The last term of the RHS of the previous inequality goes to 0, as $n \rightarrow \infty$, since $f \in \mathcal{S}(\mathbb{R})$. Therefore are reduced to show that

$$\lim_{n \rightarrow \infty} \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) + \frac{4\pi^2\xi^2}{\sqrt{2\lambda c}} \right|^2 |\mathcal{F}_n(f)(\xi)|^2 d\xi = 0.$$

Since $f \in \mathcal{S}(\mathbb{R})$, a simple application of Lemma 7.1 shows that it is equivalent to prove that

$$\lim_{n \rightarrow \infty} \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) + \frac{4n^2}{\sqrt{2\lambda c}} \sin^2\left(\pi \frac{\xi}{n}\right) \right|^2 |\mathcal{F}_n(f)(\xi)|^2 d\xi = 0.$$

Observe now that

$$\begin{aligned} & \left| q_n\left(\frac{\xi}{n}\right) + \frac{4n^2}{\sqrt{2\lambda c}} \sin^2\left(\pi \frac{\xi}{n}\right) \right| \\ & = \frac{4n^2}{\sqrt{2\lambda c}} \sin^2\left(\pi \frac{\xi}{n}\right) \left| \frac{1}{\sqrt{1 + \frac{cn^{-b}}{2\lambda} + \frac{n^b\lambda}{2c} \sin^2\left(\pi \frac{\xi}{n}\right) + \sqrt{\frac{cn^{-b}}{2\lambda}}}} - 1 \right|. \end{aligned} \tag{4.11}$$

In particular, we have

$$\left| q_n\left(\frac{\xi}{n}\right) + \frac{4n^2}{\sqrt{2\lambda c}} \sin^2\left(\pi \frac{\xi}{n}\right) \right| \leq C|\xi|^2. \tag{4.12}$$

Observe that by Lemma 7.1

$$\limsup_{A \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{A \leq |\xi| \leq n/2} |\xi|^4 |\mathcal{F}_n(f)(\xi)|^2 d\xi = 0.$$

Thus, by (4.12), it is sufficient to prove that for any $A > 0$ fixed,

$$\lim_{n \rightarrow \infty} \int_{|\xi| \leq A} \left| q_n\left(\frac{\xi}{n}\right) + \frac{4n^2}{\sqrt{2\lambda c}} \sin^2\left(\pi \frac{\xi}{n}\right) \right|^2 |\mathcal{F}_n(f)(\xi)|^2 d\xi = 0.$$

By (4.11) we have

$$\begin{aligned} & \int_{|\xi| \leq A} \left| q_n\left(\frac{\xi}{n}\right) + \frac{4n^2}{\sqrt{2\lambda c}} \sin^2\left(\pi \frac{\xi}{n}\right) \right|^2 |\mathcal{F}_n(f)(\xi)|^2 d\xi \\ & \leq \varepsilon_n(A) \int_{|\xi| \leq A} |\mathcal{F}_n(f)(\xi)|^2 d\xi \leq \frac{\varepsilon_n(A)}{n} \sum_{x \in \mathbb{Z}} f^2\left(\frac{x}{n}\right), \end{aligned}$$

where

$$\varepsilon_n(A) = \sup_{|\xi| \leq A} \left\{ \frac{4n^2}{\sqrt{2\lambda c}} \sin^2 \left(\pi \frac{\xi}{n} \right) \left| \frac{1}{\sqrt{1 + \frac{cn^{-b}}{2\lambda} + \frac{n^b \lambda}{2c} \sin^2 \left(\pi \frac{\xi}{n} \right)} + \sqrt{\frac{cn^{-b}}{2\lambda}}} - 1 \right| \right\}$$

goes to 0 as $n \rightarrow \infty$ since $b < 2$. Since

$$\sup_n \frac{1}{n} \sum_{x \in \mathbb{Z}} f^2 \left(\frac{x}{n} \right) < \infty$$

the claim follows. □

Lemma 4.5 (Boltzman–Gibbs principle) *If $b < 2/3$, then*

$$\lim_{n \rightarrow \infty} |\Psi_t^n(f, g)| = 0.$$

Proof Recall (4.9). Performing a simple computation we rewrite ψ as

$$\psi(\omega) = \sum_{x \in \mathbb{Z}} \sum_{k \geq 1} (\rho_{k-1}(x+1) + \rho_{k+1}(x)) \nabla \omega_x \omega_{x+k},$$

with the convention that $\rho_0(x) = 0$ for any $x \in \mathbb{Z}$. Let us introduce ψ_k defined by

$$\psi_k(x) = \rho_{k-1}(x+1) + \rho_{k+1}(x).$$

By using the translation invariance of μ_β we get that

$$\Psi_t^n(f, g) = n^{a-2} \left\langle \left\{ \sum_{y \in \mathbb{Z}} g \left(\frac{y}{n} \right) \left(\omega_y^2(0) - \frac{1}{\beta} \right) \right\} \int_0^t \varphi(\omega(sn^a)) ds \right\rangle_\beta$$

where the function φ is given by

$$\varphi(\omega) = \sum_{x, z \in \mathbb{Z}} \sum_{k \geq 1} (\nabla_n f) \left(\frac{x}{n} \right) [\psi_k(x-1) - \psi_k(x)] (\omega_{x+z} \omega_{x+z+k}). \tag{4.13}$$

Thus, by the Cauchy–Schwarz inequality, it is sufficient to prove that the $\mathbf{L}^2(\mathbb{P}_{\mu_\beta})$ -norm of

$$n^{a-3/2} \int_0^t \varphi(\omega(sn^a)) ds$$

vanishes with n , where \mathbb{P}_{μ_β} denotes the law of the Markov process $\{\omega(tn^a)\}_{t \geq 0}$ starting from μ_β . We denote by \mathbb{E}_{μ_β} the corresponding expectation. By a general inequality for the variance of additive functionals of Markov processes (see [15], Lemma 2.4), we have

$$\begin{aligned} \mathbb{E}_{\mu_\beta} \left[\left(\int_0^t \varphi(\omega(sn^a)) ds \right)^2 \right] &\leq Ct \left\langle \varphi (t^{-1} - n^a S_n)^{-1} \varphi \right\rangle_\beta \\ &= Ctn^{-a} \left\langle \varphi ([tn^a]^{-1} - S_n)^{-1} \varphi \right\rangle_\beta. \end{aligned} \tag{4.14}$$

We can now use some ideas from [6] in order to have a very sharp estimate of (4.14). In Appendix 2, we prove that $\langle \varphi ([tn^a]^{-1} - S_n)^{-1} \varphi \rangle_\beta$ is bounded from above by Cn^{2b} . As a consequence, the Boltzmann–Gibbs principle holds if $a - 3 + 2b < 0$, and with the condition $a = 2 - b/2$ it implies $b < 2/3$. □

5 Superdiffusive Domain of the Energy

In this section we give the strategy of the proof of Theorem 3.2, which is the same as in [5]. In the whole section, $a = 3/2$, and $b > 1$. We also assume $\beta = 1$, the general case follows after an easy change of variables in the Markov process.

5.1 Weak Formulation

Two Coupled Differential Equations Let g be a fixed function in $\mathcal{S}(\mathbb{R})$. We define the process $\{\mathcal{S}_t^n; t \geq 0\}$ acting on functions $f \in \mathcal{S}(\mathbb{R})$ as

$$\mathcal{S}_t^n(f) := \frac{1}{2} \sigma_t^n(f, g). \tag{5.1}$$

for any $t \geq 0, n \in \mathbb{N}$. After arranging terms in a convenient way we have that

$$\mathcal{S}_t^n(f) = \frac{1}{2} \left\langle \left(\frac{1}{\sqrt{n}} \sum_{x \in \mathbb{Z}} g\left(\frac{x}{n}\right) (\omega_x^2(0) - 1) \right) \left(\frac{1}{\sqrt{n}} \sum_{y \in \mathbb{Z}} f\left(\frac{y}{n}\right) (\omega_y^2(tn^{3/2}) - 1) \right) \right\rangle_1.$$

For a function $h \in \mathcal{S}(\mathbb{R}^2)$ we define $\{Q_t^n(h); t \geq 0\}$ as

$$Q_t^n(h) = \frac{1}{2} \left\langle \left(\frac{1}{\sqrt{n}} \sum_{x \in \mathbb{Z}} g\left(\frac{x}{n}\right) (\omega_x^2(0) - 1) \right) \left(\frac{1}{n} \sum_{\substack{y, z \in \mathbb{Z} \\ y \neq z}} h\left(\frac{y}{n}, \frac{z}{n}\right) \omega_y(tn^{3/2}) \omega_z(tn^{3/2}) \right) \right\rangle_1.$$

Notice that $Q_t^n(h)$ depends only on the symmetric part of the function h . Therefore, we will always assume, without loss of generality, that $h(x, y) = h(y, x)$ for any $x, y \in \mathbb{Z}$. Notice as well that $Q_t^n(h)$ does not depend on the values of h at the diagonal $\{x = y\}$. Let us write now the differential equations for $\mathcal{S}_t^n(f)$ and $Q_t^n(h)$. We start by introducing some definitions.

Definition 5.1 (*Discrete approximations*) For $f \in \mathcal{S}(\mathbb{R})$ and $h \in \mathcal{S}(\mathbb{R}^2)$, we define the discrete approximations

- (i) $\Delta_n f : \mathbb{R} \rightarrow \mathbb{R}$ of the second derivative of f as

$$\Delta_n f\left(\frac{x}{n}\right) = n^2 \left(f\left(\frac{x+1}{n}\right) + f\left(\frac{x-1}{n}\right) - 2f\left(\frac{x}{n}\right) \right),$$

- (ii) $\nabla_n f \otimes \delta : \frac{1}{n} \mathbb{Z}^2 \rightarrow \mathbb{R}$ of the distribution $f'(x)\delta(x = y)$ as

$$(\nabla_n f \otimes \delta)\left(\frac{x}{n}, \frac{y}{n}\right) = \begin{cases} \frac{n^2}{2} (f(\frac{x+1}{n}) - f(\frac{x}{n})); & y = x + 1 \\ \frac{n^2}{2} (f(\frac{x}{n}) - f(\frac{x-1}{n})); & y = x - 1 \\ 0; & \text{otherwise,} \end{cases} \tag{5.2}$$

- (iii) $\Delta_n h : \mathbb{R}^2 \rightarrow \mathbb{R}$ of the Laplacian of h as

$$\Delta_n h\left(\frac{x}{n}, \frac{y}{n}\right) = n^2 \left(h\left(\frac{x+1}{n}, \frac{y}{n}\right) + h\left(\frac{x-1}{n}, \frac{y}{n}\right) + h\left(\frac{x}{n}, \frac{y+1}{n}\right) + h\left(\frac{x}{n}, \frac{y-1}{n}\right) - 4h\left(\frac{x}{n}, \frac{y}{n}\right) \right),$$

- (iv) $A_n h : \mathbb{R}^2 \rightarrow \mathbb{R}$ of the directional derivative $(-2, -2) \cdot \nabla h$ as

$$A_n h\left(\frac{x}{n}, \frac{y}{n}\right) = n \left(h\left(\frac{x}{n}, \frac{y-1}{n}\right) + h\left(\frac{x-1}{n}, \frac{y}{n}\right) - h\left(\frac{x}{n}, \frac{y+1}{n}\right) - h\left(\frac{x+1}{n}, \frac{y}{n}\right) \right),$$

(v) $\mathcal{D}_n h : \frac{1}{n}\mathbb{Z} \rightarrow \mathbb{R}$ of the directional derivative of h along the diagonal $\{x = y\}$ as

$$\mathcal{D}_n h\left(\frac{x}{n}\right) = n\left(h\left(\frac{x}{n}, \frac{x+1}{n}\right) - h\left(\frac{x-1}{n}, \frac{x}{n}\right)\right),$$

(vi) $\tilde{\mathcal{D}}_n h : \frac{1}{n}\mathbb{Z}^2 \rightarrow \mathbb{R}$ of the distribution $\partial_y h(x, x) \otimes \delta(x = y)$ as

$$\tilde{\mathcal{D}}_n h\left(\frac{x}{n}, \frac{y}{n}\right) = \begin{cases} n^2\left(h\left(\frac{x}{n}, \frac{x+1}{n}\right) - h\left(\frac{x}{n}, \frac{x}{n}\right)\right); & y = x + 1 \\ n^2\left(h\left(\frac{x-1}{n}, \frac{x}{n}\right) - h\left(\frac{x-1}{n}, \frac{x-1}{n}\right)\right); & y = x - 1 \\ 0; & \text{otherwise.} \end{cases}$$

The following proposition can be deduced after straightforward computations:

Proposition 5.2 For all functions $f \in \mathcal{S}(\mathbb{R})$ and $h \in \mathcal{S}(\mathbb{R}^2)$,

$$\frac{d}{dt} S_t^n(f) = -2Q_t^n(\nabla_n f \otimes \delta) + S_t^n(n^{-1/2} \Delta_n f), \tag{5.3}$$

$$\frac{d}{dt} Q_t^n(h) = Q_t^n(L_n h) - 2S_t^n(\mathcal{D}_n h) + 2Q_t^n(n^{-1/2} \tilde{\mathcal{D}}_n h), \tag{5.4}$$

where the operator L_n is defined by

$$L_n = \sqrt{n} A_n + \frac{1}{\sqrt{n}} \Delta_n - 4n^{3/2} \gamma_n \text{Id}. \tag{5.5}$$

A Priori Bounds For $f \in \mathcal{S}(\mathbb{R})$, define the weighted $\ell^2(\mathbb{Z})$ -norm as

$$\|f\|_{2,n} = \sqrt{\frac{1}{n} \sum_{x \in \mathbb{Z}} f\left(\frac{x}{n}\right)^2}.$$

By the Cauchy–Schwarz inequality we have the *a priori* bound

$$|S_t^n(f)| \leq \|g\|_{2,n} \|f\|_{2,n} \tag{5.6}$$

for any $t \geq 0$, any $n \in \mathbb{N}$ and any $f, g \in \mathcal{S}(\mathbb{R})$. Therefore, the term $S_t^n\left(\frac{1}{\sqrt{n}} \Delta_n f\right)$ is negligible, as $n \rightarrow \infty$. In (5.3), the term $Q_t^n(\nabla_n f \otimes \delta)$ is the relevant one. We also have the *a priori* bound

$$|Q_t^n(h)| \leq 2\|g\|_{2,n} \|\bar{h}\|_{2,n}, \tag{5.7}$$

where $\|\bar{h}\|_n$ is the weighted $\ell^2(\mathbb{Z}^2)$ -norm of \bar{h}

$$\|\bar{h}\|_{2,n} = \sqrt{\frac{1}{n^2} \sum_{x,y \in \mathbb{Z}} \bar{h}\left(\frac{x}{n}, \frac{y}{n}\right)^2}$$

and \bar{h} is defined by

$$\bar{h}\left(\frac{x}{n}, \frac{y}{n}\right) = h\left(\frac{x}{n}, \frac{y}{n}\right) \mathbf{1}_{x \neq y}.$$

In Eq. (5.3), both fields S_t^n and Q_t^n appear with non-negligible terms. Moreover, the term involving Q_t^n is quite singular, since it involves an approximation of a distribution. Let us recall the clever strategy well-detailed in [5]: given $f \in \mathcal{S}(\mathbb{R})$, if we choose h such that $L_n h = \nabla_n f \otimes \delta$, we can try to cancel the terms $Q_t^n(\nabla_n f \otimes \delta)$ and $Q_t^n(L_n h)$ out. Then the term $S_t^n(\mathcal{D}_n h)$ provides a non-trivial drift for the differential equation (5.3) and the term $Q_t^n(n^{-1/2} \tilde{\mathcal{D}}_n h)$ turns out to be negligible.

5.2 Sketch of the Proof

After giving the topological setting needed for Theorem 3.2, we sketch the main steps of its proof, which are detailed in [5].

Topological Setting Let us define the *Hermite polynomials* $H_\ell : \mathbb{R} \rightarrow \mathbb{R}$ as

$$H_\ell(x) = (-1)^\ell e^{\frac{x^2}{2}} \frac{d^\ell}{dx^\ell} \left[e^{-\frac{x^2}{2}} \right]$$

for any $\ell \in \mathbb{N}_0$ and any $x \in \mathbb{R}$. We define the *Hermite functions* $f_\ell : \mathbb{R} \rightarrow \mathbb{R}$ as

$$f_\ell(x) = \frac{1}{\sqrt{\ell! \sqrt{2\pi}}} H_\ell(x) e^{-\frac{x^2}{4}},$$

for any $\ell \in \mathbb{N}_0$ and any $x \in \mathbb{R}$. The Hermite functions $\{f_\ell; \ell \in \mathbb{N}_0\}$ form an orthonormal basis of $\mathbf{L}^2(\mathbb{R})$. Let us denote by $C_c^\infty(\mathbb{R})$ the space of smooth functions with compact support. For each $k \in \mathbb{R}$, we define the *Sobolev space* \mathcal{H}_k as the completion of $C_c^\infty(\mathbb{R})$ with respect to the norm $\|\cdot\|_{\mathcal{H}_k}$ defined as

$$\|g\|_{\mathcal{H}_k} = \sqrt{\sum_{\ell \in \mathbb{N}_0} (1 + \ell)^{2k} \langle f_\ell, g \rangle^2}$$

for any $g \in C_c^\infty(\mathbb{R})$. Here we use the notation $\langle f_\ell, g \rangle = \int g(x) f_\ell(x) dx$. Let us denote by $\mathcal{C}([0, T]; \mathcal{H}_k)$ the space of continuous functions from $[0, T]$ to \mathcal{H}_k .

Main Steps of the Proof First, we need to show tightness, and then to characterize the limit points of weakly converging subsequences.

(1) *Tightness* The same standard arguments exposed in [5] imply the following

Lemma 5.3 *For any $k > \frac{19}{24}$, the sequence $\{S_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ is weakly relatively compact in $\mathbf{L}^2([0, T]; \mathcal{H}_{-k})$. Moreover, for any $t \in [0, T]$ fixed, the sequence $\{S_t^n; n \in \mathbb{N}\}$ is sequentially, weakly relatively compact in \mathcal{H}_{-k} .*

(2) *Characterization of Limit Points* Fix $k > \frac{19}{24}$ and let us consider a limit point of $\{S_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ with respect to the weak topology of $\mathbf{L}^2([0, T]; \mathcal{H}_{-k})$. Without loss of generality we can denote by n the subsequence for which $\{S_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ converges to $\{S_t; t \in [0, T]\}$. The aim is to prove the following

Proposition 5.4 *Let $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function of compact support (in $C_c^\infty([0, T] \times \mathbb{R})$). Then,*

$$\mathcal{S}_T(f_T) = \mathcal{S}_0(f_0) + \int_0^T \mathcal{S}_t((\partial_t + \mathbb{L})f_t) dt, \tag{5.8}$$

where \mathbb{L} is defined in (3.1).

As a result, $\{S_t; t \in [0, T]\}$ is a *weak solution* of the fractional heat equation:

$$\partial_t u = -\frac{1}{\sqrt{2}} \{(-\Delta)^{3/4} + \nabla(-\Delta)^{1/4}\} u,$$

as defined in (2.1) of [13]. In Section 8.1 of [13], it is shown that there is a unique solution of (5.8) and therefore the limit process $\{S_t; t \in [0, T]\}$ is unique. Proposition 5.4 is the most challenging part of the proof, and the next section is devoted to it.

(3) *Conclusion* The proof of Theorem 3.2 is almost done: the first two points imply that the sequence $\{S_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ weakly converges in $\mathbf{L}^2([0, T]; \mathcal{H}_{-k})$ to a unique limit point,

denoted by $\{S_t; t \in [0, T]\}$. It can be proved that the convergence also holds for fixed times $t \in [0, T]_{\mathbb{Q}}$ with respect to the weak topology of \mathcal{H}_{-k} , where

$$[0, T]_{\mathbb{Q}} := \left\{ t \in [0, T]; \frac{t}{T} \in \mathbb{Q} \right\}.$$

Since T is arbitrary, this last convergence holds for any $t \in [0, \infty)$. In particular, $S_t^n(f)$ converges to $S_t(f)$, as $n \rightarrow \infty$, for any $f \in C_c^\infty(\mathbb{R})$.

The main differences between the model in [5] and ours rely on the additional velocity-flip noise, which is of intensity γ_n . This perturbation first appears in the definition of the operator L_n in (5.5). Then, some technical proofs have to be slightly modified. More precisely, rigorous convergence estimates lead to the condition: $b > 1$.

5.3 Convergence Estimates

Here we give the proof of Proposition 5.4. Recall that $b > 1$. Let us assume that $\{S_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ converges to $\{S_t; t \in [0, T]\}$ with respect to the weak topology of $L^2([0, T]; \mathcal{H}_{-k})$.

Let us fix $f \in \mathcal{S}(\mathbb{R})$ and let $h_n : \frac{1}{n}\mathbb{Z} \times \frac{1}{n}\mathbb{Z} \rightarrow \mathbb{R}$ be the solution of the equation

$$L_n h_n = \nabla_n f \otimes \delta. \tag{5.9}$$

The following properties of h_n are shown in Appendix 3, following [5]:

Lemma 5.5 *Let $f \in \mathcal{S}(\mathbb{R})$. The solution of (5.9) satisfies*

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{x, y \in \mathbb{Z}} h_n^2\left(\frac{x}{n}, \frac{y}{n}\right) = 0, \tag{5.10}$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{x \in \mathbb{Z}} \left| \mathcal{D}_n h_n\left(\frac{x}{n}\right) + \frac{1}{4} \mathbb{L}f\left(\frac{x}{n}\right) \right|^2 = 0. \tag{5.11}$$

In other words, $\|h_n\|_{2,n}$ and $\|\mathcal{D}_n h_n + \frac{1}{4} \mathbb{L}f\|_{2,n}$ converge to 0, as $n \rightarrow \infty$.

By (5.3) and (5.4), we see that

$$\begin{aligned} S_T^n(f) &= S_0^n(f) + \int_0^T S_t^n(-4\mathcal{D}_n h_n) dt + 2[Q_0^n(h_n) - Q_T^n(h_n)] \\ &\quad + \int_0^T S_t^n\left(\frac{1}{\sqrt{n}} \Delta_n f\right) dt + 4 \int_0^T Q_t^n\left(\frac{1}{\sqrt{n}} \tilde{\mathcal{D}}_n(h_n)\right) dt. \end{aligned}$$

Therefore, by the *a priori* bound (5.7) and Lemma 5.5, we have that

$$S_T^n(f) = S_0^n(f) + \int_0^T S_t^n(\mathbb{L}f) dt + 4 \int_0^T Q_t^n\left(\frac{1}{\sqrt{n}} \tilde{\mathcal{D}}_n(h_n)\right) dt \tag{5.12}$$

plus an error term which goes to 0, as $n \rightarrow \infty$. It turns out that the *a priori* bound (5.7) is not sufficient to show that the last term on the RHS of (5.12) goes to 0 with n since

$$\frac{1}{n^3} \sum_{x \in \mathbb{Z}} (\tilde{\mathcal{D}}_n h_n)^2\left(\frac{x}{n}, \frac{x+1}{n}\right) \tag{5.13}$$

is of order one. Therefore we use again (5.4) applied to $h = v_n$ where v_n is the solution of the Poisson equation

$$L_n v_n = n^{-1/2} \tilde{\mathcal{D}}_n h_n. \tag{5.14}$$

Then we have

$$\int_0^T Q_t^n \left(\frac{1}{\sqrt{n}} \tilde{D}_n h_n \right) dt = 2 \int_0^T S_t^n (\mathcal{D}_n v_n) dt - 2 \int_0^T Q_t^n \left(\frac{1}{\sqrt{n}} \tilde{D}_n v_n \right) dt + Q_T^n(v_n) - Q_0^n(v_n).$$

Now, we can use the *a priori* bounds (5.6) and (5.7). The following estimates on v_n are proved in Appendix 3.

Lemma 5.6 *The solution v_n of (5.14) satisfies*

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{x, y \in \mathbb{Z}} v_n^2 \left(\frac{x}{n}, \frac{y}{n} \right) = 0, \tag{5.15}$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{x \in \mathbb{Z}} (\mathcal{D}_n v_n)^2 \left(\frac{x}{n} \right) = 0, \tag{5.16}$$

$$\lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{x \in \mathbb{Z}} (\tilde{D}_n v_n)^2 \left(\frac{x}{n}, \frac{x+1}{n} \right) = 0. \tag{5.17}$$

In other words, $\|v_n\|_{2,n}$, $\|\mathcal{D}_n v_n\|_{2,n}$ and $\|\frac{1}{\sqrt{n}} \tilde{D}_n v_n\|_{2,n}$ converge to 0, as $n \rightarrow \infty$.

It follows that

$$S_T^n(f) = S_0^n(f) + \int_0^T S_t^n(\mathbb{L}f) dt \tag{5.18}$$

plus an error term which goes to 0, as $n \rightarrow \infty$. Recall that $\{S_t^n; t \in [0, T]\}_{n \in \mathbb{N}}$ weakly converges to $\{S_t; t \in [0, T]\}$. The main difficulty is that the operator \mathbb{L} is an integro-differential operator with heavy tails (in other words, even for $f \in C_c^\infty(\mathbb{R})$ the function $\mathbb{L}f$ has heavy tails). As a result, we cannot take the limit $n \rightarrow \infty$ in (5.18). Bernardin et al. in [5] achieved Proposition 5.4 after truncature considerations, together with some results about the Lipschitz property of the function $t \mapsto S_t(f)$. We refer the reader to their paper, and also to [11] and [13] for useful properties of the fractional Laplacian.

In Appendix 3, Lemma 5.6 and Lemma 5.5 are proved. Here, the computations are similar to [5], but we take into account the additional term due to the presence of the velocity-flip noise, and explain the needed assumption on the parameter b .

6 Volume Fluctuations

Recall that the volume $\sum_{x \in \mathbb{Z}} \omega_x$ is conserved if and only if $\gamma = 0$. In this section we give the behavior of the space-time correlation functions of the volume fluctuation field defined for $f, g \in \mathcal{S}(\mathbb{R})$ as in (3.4).

6.1 Explicit Computations and Fourier Transform

Contrary to the energy fluctuation field, here computations are explicit. Let us introduce the following notation:

$$\mathcal{V}(x, t) := \langle \omega_x(t) \omega_0(0) \rangle_\beta,$$

and notice that for all $x \in \mathbb{Z}$ and $t > 0$

$$\begin{aligned} \frac{d}{dt} \mathcal{V}(x, t) &= \mathcal{V}(x + 1, t) - \mathcal{V}(x - 1, t) - 2\gamma_n \mathcal{V}(x, t) \\ &\quad + \lambda (\mathcal{V}(x + 1, t) + \mathcal{V}(x - 1, t) - 2\mathcal{V}(x, t)). \end{aligned}$$

This infinite ODE’s system can be rewritten for the Fourier transform $\widehat{\mathcal{V}}$ as

$$\frac{d}{dt}\widehat{\mathcal{V}}(\theta, t) = [-2i \sin(2\pi\theta) - 2\gamma_n - 4\lambda \sin^2(\pi\theta)]\widehat{\mathcal{V}}(\theta, t), \quad \theta \in \mathbb{T}, t > 0.$$

Since $\widehat{\mathcal{V}}(\theta, 0) = \beta^{-1}$ for all $\theta \in \mathbb{T}$, we conclude that

$$\widehat{\mathcal{V}}(\theta, t) = \beta^{-1} e^{t[-2i \sin(2\pi\theta) - 2\gamma_n - 4\lambda \sin^2(\pi\theta)]}.$$

We assume from now on that $\beta = 1$. The inverse Fourier transform gives

$$\begin{aligned} \eta_t^n(f, g) &= \frac{1}{n} \sum_{y, z \in \mathbb{Z}} f\left(\frac{y+z}{n}\right) g\left(\frac{y}{n}\right) \int_{[-\frac{1}{2}, \frac{1}{2}]} \widehat{\mathcal{V}}(\theta, tn^a) e^{-2i\pi\theta z} d\theta \\ &= \frac{1}{n} \sum_{y, z \in \mathbb{Z}} f\left(\frac{y+z}{n}\right) g\left(\frac{y}{n}\right) \int_{[-\frac{1}{2}, \frac{1}{2}]} e^{tn^a[-2i \sin(2\pi\theta) - 2\gamma_n - 4\lambda \sin^2(\pi\theta)] - 2i\pi\theta z} d\theta \\ &= \frac{1}{n} \sum_{y \in \mathbb{Z}} \left\{ g\left(\frac{y}{n}\right) \frac{1}{n} \sum_{z \in \mathbb{Z}} f\left(\frac{y+z}{n}\right) \int_{[-\frac{n}{2}, \frac{n}{2}]} e^{tn^a[-2i \sin\left(2\pi\frac{\xi}{n}\right) - 2\gamma_n - 4\lambda \sin^2\left(\pi\frac{\xi}{n}\right)] - 2i\pi\frac{\xi}{n}z} d\xi \right\}. \end{aligned} \tag{6.1}$$

Case $a > 2$ or $b < a$. Now, we study the convergence of the quantity $\mathcal{J}_n(t) e^{-2tn^a\gamma_n}$ where

$$\mathcal{J}_n(t) := \frac{1}{n} \sum_{z \in \mathbb{Z}} f\left(\frac{y+z}{n}\right) \int_{[-\frac{n}{2}, \frac{n}{2}]} e^{tn^a[-2i \sin(2\pi\frac{\xi}{n}) - 4\lambda \sin^2(\pi\frac{\xi}{n})] - 2i\pi\frac{\xi}{n}z} d\xi.$$

The modulus of the integrand above equals $e^{-4\lambda tn^a \sin^2\left(\frac{\pi\xi}{n}\right)}$, and it can be easily proved that

$$\int_{[-\frac{n}{2}, \frac{n}{2}]} e^{-4\lambda tn^a \sin^2\left(\frac{\pi\xi}{n}\right)} d\xi \begin{cases} \xrightarrow[n \rightarrow \infty]{} 0; & \text{if } a > 2, \\ \xrightarrow[n \rightarrow \infty]{} \int_{\mathbb{R}} e^{-4\lambda t \pi^2 \xi^2} d\xi; & \text{if } a = 2, \\ \text{is bounded by } n; & \text{if } a < 2. \end{cases}$$

The first two statements are consequences of the dominated convergence theorem. Therefore, from the fact that $\sup_n \left\{ \frac{1}{n} \sum_{z \in \mathbb{Z}} |f\left(\frac{z}{n}\right)| \right\} < +\infty$, we conclude:

- (i) If $a > 2$, then $\eta_t^n(f, g)$ vanishes, as $n \rightarrow \infty$, for any b (because of the above integral),
- (ii) If $a \leq 2$ and $b < a$, then $\eta_t^n(f, g)$ vanishes as $n \rightarrow \infty$ (because of $e^{-2tn^a\gamma_n} = e^{-2tc n^{a-b}}$).

This proves the statements $A(iii)$, $B(v)$ and $C(v)$ of Theorem 3.3. In the following we distinguish the cases $a \leq 1$ and $a > 1$.

Case $a \leq 1$ and $b \geq a$. We introduce:

$$q_n(\theta) := n^a [-2i \sin(2\pi\theta) - 2\gamma_n - 4\lambda \sin^2(\pi\theta)]. \tag{6.2}$$

From (6.1), we get

$$\eta_t^n(f, g) = \int_{[-\frac{n}{2}, \frac{n}{2}]} \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \mathcal{F}_n(f)(\xi) \mathcal{F}_n(g)(\xi) d\xi \tag{6.3}$$

and

$$\begin{aligned}
 \frac{d}{dt} \eta_t^n(f, g) &= \frac{1}{n} \sum_{y, z \in \mathbb{Z}} f\left(\frac{y+z}{n}\right) g\left(\frac{y}{n}\right) \int_{[-\frac{1}{2}, \frac{1}{2}]} q_n(\theta) \widehat{\mathcal{V}}(\theta, tn^a) e^{-2i\pi\theta z} d\theta \\
 &= \frac{1}{n} \sum_{y \in \mathbb{Z}} \left\{ g\left(\frac{y}{n}\right) \frac{1}{n} \sum_{z \in \mathbb{Z}} f\left(\frac{y+z}{n}\right) \int_{[-\frac{n}{2}, \frac{n}{2}]} q_n\left(\frac{\xi}{n}\right) \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) e^{-2i\pi\frac{\xi}{n}z} d\xi \right\} \\
 &= \int_{[-\frac{n}{2}, \frac{n}{2}]} q_n\left(\frac{\xi}{n}\right) \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \mathcal{F}_n(f)(\xi) \mathcal{F}_n(g)(\xi) d\xi.
 \end{aligned} \tag{6.4}$$

Let us write down the lemma that we are going to prove.

Lemma 6.1 *Reformulation of A(ii). If $b \leq 1$ and $b = a$, then*

$$\int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi) - \mathcal{F}_n(\mathbf{1}_{b=1} 2f' - 2cf)(\xi) \right| \left| \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \mathcal{F}_n(g)(\xi) \right| d\xi \xrightarrow{n \rightarrow \infty} 0. \tag{6.5}$$

Reformulation of A(i), B(i) and C(i). If $a \in (0, 1)$ and $a < b$,

$$\left| \int_{[-\frac{n}{2}, \frac{n}{2}]} q_n\left(\frac{\xi}{n}\right) \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \mathcal{F}_n(f)(\xi) \mathcal{F}_n(g)(\xi) d\xi \right| \xrightarrow{n \rightarrow \infty} 0. \tag{6.6}$$

Reformulation of B(iii) and C(iii). If $a = 1$ and $b > 1$, then

$$\int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi) - \mathcal{F}_n(2f')(\xi) \right|^2 \left| \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \mathcal{F}_n(g)(\xi) \right|^2 d\xi \xrightarrow{n \rightarrow \infty} 0. \tag{6.7}$$

Proof We use some powerful results regarding the Fourier transform, and the argument will be repeated many times.

First, we need the following, easy-to-prove, fact: for any $a \leq 2$ and any $g \in \mathcal{S}(\mathbb{R})$, there exists a constant $C > 0$ such that, for any n and any $\xi \in [-\frac{n}{2}, \frac{n}{2}]$,

$$\left| \mathcal{F}_n(g)(\xi) \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \right| \leq C. \tag{6.8}$$

Let us prove for instance the first point C(iii) with $b = a = 1$. The other statements can be obtained with the same procedure. From (6.3) and (6.4), we can write

$$\begin{aligned}
 & \left| \partial_t \eta_t^n(f, g) - \eta_t^n(2f' - 2cf, h) \right| \\
 & \leq \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi) - \mathcal{F}_n(2f' - 2cf)(\xi) \right| \left| \mathcal{F}_n(g)(\xi) \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \right| d\xi \\
 & \leq C \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi) - \mathcal{F}_n(2f' - 2cf)(\xi) \right| d\xi.
 \end{aligned}$$

Then, the strategy is to replace $\mathcal{F}_n(2f' - 2cf)(\xi)$ with

$$\mathcal{F}(2f' - 2cf)(\xi) = (-4i\pi\xi - 2c)(\mathcal{F}f)(\xi),$$

and this can be done thanks to Corollary 7.2. We write then

$$\begin{aligned} & \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi) - (-4i\pi\xi - 2c) \mathcal{F}(f)(\xi) \right| d\xi \\ & \leq \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) + 4i\pi\xi + 2c \right| |\mathcal{F}_n(f)(\xi)| d\xi \\ & \quad + \int_{[-\frac{n}{2}, \frac{n}{2}]} (4\pi\xi + 2c) |\mathcal{F}_n(f)(\xi) - (\mathcal{F}f)(\xi)| d\xi. \end{aligned}$$

Again, from Corollary 7.2, the last term of the RHS goes to 0, as $n \rightarrow \infty$. We are reduced to show that

$$\lim_{n \rightarrow \infty} \int_{[-\frac{n}{2}, \frac{n}{2}]} \left| q_n\left(\frac{\xi}{n}\right) + 4i\pi\xi + 2c \right| |\mathcal{F}_n(f)(\xi)| d\xi = 0.$$

It is now easy to see that if $a = b = 1$, then there exists a constant $C > 0$ such that

$$\left| q_n\left(\frac{\xi}{n}\right) + 4i\pi\xi + 2c \right| = \left| 2i[2\pi\xi - n \sin(2\pi\frac{\xi}{n})] - 4\lambda n \sin^2(\pi\frac{\xi}{n}) \right| \leq C\xi^2.$$

Indeed, this is a consequence of the following fact: $\forall x \in [-\pi, \pi], |\sin(x) - x| \leq \frac{|x|^3}{3}$. We recover the same type of inequality as (4.12), and the conclusion comes straightforwardly. \square

Case $1 < a \leq 2$ and $b \geq a$. All the previous computations can be adapted to the translated field $\tilde{\eta}_t^n$, whose definition is recalled below:

$$\tilde{\eta}_t^n(f, g) = \frac{1}{n} \sum_{x, y \in \mathbb{Z}} f\left(\frac{x-2tn^a}{n}\right) g\left(\frac{y}{n}\right) \langle \omega_x(tn^a) \omega_y(0) \rangle_\beta.$$

The translation gives rise to a multiplicative term in the computation of the Fourier transform. In the same way, we define $\tilde{q}_n(\theta) := q_n(\theta) + n^a 4i\pi\theta$, where $q_n(\theta)$ was given in (6.2) and we can easily write

$$\begin{aligned} \tilde{\eta}_t^n(f, g) &= \int_{[-\frac{n}{2}, \frac{n}{2}]} \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) e^{4i\pi tn^a \frac{\xi}{n}} \mathcal{F}_n(f)(\xi) \mathcal{F}_n(g)(\xi) d\xi, \\ \frac{d}{dt} \tilde{\eta}_t^n(f, g) &= \int_{[-\frac{n}{2}, \frac{n}{2}]} \tilde{q}_n\left(\frac{\xi}{n}\right) \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) e^{4i\pi tn^a \frac{\xi}{n}} \mathcal{F}_n(f)(\xi) \mathcal{F}_n(g)(\xi) d\xi. \end{aligned}$$

Lemma 6.2 *Reformulation of B(ii) and C(ii). If $a \in (1, 2)$ and $a < b$,*

$$\left| \int_{[-\frac{n}{2}, \frac{n}{2}]} \tilde{q}_n\left(\frac{\xi}{n}\right) \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) e^{4i\pi tn^a \frac{\xi}{n}} \mathcal{F}_n(f)(\xi) \mathcal{F}_n(g)(\xi) d\xi \right| \xrightarrow[n \rightarrow \infty]{} 0. \tag{6.9}$$

Reformulation of B(iv). If $a \in (1, 2)$ and $a = b$,

$$\int_{[-\frac{n}{2}, \frac{n}{2}]} \left| \tilde{q}_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi) - \mathcal{F}_n(-2cf)(\xi) \right| \left| e^{4i\pi tn^a \frac{\xi}{n}} \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \mathcal{F}_n(g)(\xi) \right| d\xi \xrightarrow[n \rightarrow \infty]{} 0. \tag{6.10}$$

Reformulation of C(iii). If $a = 2$ and $b \geq 2$, then

$$\int_{[-\frac{n}{2}, \frac{n}{2}]} \left| \tilde{q}_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi) - \mathcal{F}_n(\lambda f'' - \mathbf{1}_{b=2} 2cf)(\xi) \right|^2 \left| e^{4i\pi tn^a \frac{\xi}{n}} \widehat{\mathcal{V}}\left(\frac{\xi}{n}, tn^a\right) \mathcal{F}_n(g)(\xi) \right|^2 d\xi \xrightarrow[n \rightarrow \infty]{} 0.$$

Proof We repeat the same arguments as before. First, for any $a \leq 2$ and any $g \in \mathcal{S}(\mathbb{R})$, there exists a constant $C > 0$ such that, for any n and any $\xi \in [-\frac{n}{2}, \frac{n}{2}]$,

$$|\mathcal{F}_n(g)(\xi)\widehat{\mathcal{V}}(\frac{\xi}{n}, tn^a)e^{4i\pi tn^a \frac{\xi}{n}}| \leq C. \tag{6.11}$$

Let us prove for instance the last point *C(iii)* with $b > 2$ and $a = 2$. The other statements can be obtained with the same procedure. From (6.3) and (6.4), we can write

$$\begin{aligned} & |\partial_t \widetilde{\eta}_t^n(f, g) - \lambda \widetilde{\eta}_t^n(f'', g)| \\ & \leq \int_{[-\frac{n}{2}, \frac{n}{2}]} |\widetilde{q}_n(\frac{\xi}{n})\mathcal{F}_n(f)(\xi) - \mathcal{F}_n(\lambda f'')(\xi)| |e^{4i\pi tn^a \frac{\xi}{n}} \mathcal{F}_n(g)(\xi)\widehat{\mathcal{V}}(\frac{\xi}{n}, tn^a)| d\xi \\ & \leq C \int_{[-\frac{n}{2}, \frac{n}{2}]} |\widetilde{q}_n(\frac{\xi}{n})\mathcal{F}_n(f)(\xi) - \mathcal{F}_n(\lambda f'')(\xi)| d\xi. \end{aligned}$$

As before, we are going to replace $\mathcal{F}_n(f'')(\xi)$ with $\mathcal{F}(f'')(\xi) = -4\pi^2|\xi|^2(\mathcal{F}f)(\xi)$, thanks to Corollary 7.2. We write then

$$\begin{aligned} & \int_{[-\frac{n}{2}, \frac{n}{2}]} |\widetilde{q}_n(\frac{\xi}{n})\mathcal{F}_n(f)(\xi) + 4\pi^2\lambda\xi^2\mathcal{F}(f)(\xi)| d\xi \\ & \leq \int_{[-\frac{n}{2}, \frac{n}{2}]} |\widetilde{q}_n(\frac{\xi}{n}) + 4\pi^2\lambda\xi^2| |\mathcal{F}_n(f)(\xi)| d\xi \\ & \quad + 4\pi^2\lambda \int_{[-\frac{n}{2}, \frac{n}{2}]} \xi^2 |\mathcal{F}_n(f)(\xi) - (\mathcal{F}f)(\xi)| d\xi. \end{aligned}$$

Again, from Corollary 7.2, the last term of the RHS goes to 0, as $n \rightarrow \infty$. We are reduced to show that

$$\lim_{n \rightarrow \infty} \int_{[-\frac{n}{2}, \frac{n}{2}]} |\widetilde{q}_n(\frac{\xi}{n}) + 4\pi^2\lambda\xi^2| |\mathcal{F}_n(f)(\xi)| d\xi = 0.$$

Now, if $a = 2$ and $b > 2$, then there exists a constant $C > 0$ such that $|\widetilde{q}_n(\frac{\xi}{n}) + 4\pi^2\lambda\xi^2| \leq C\xi^2$, and the conclusion follows. □

Acknowledgments PG thanks CNPq (Brazil) for support through the research project “Additive functionals of particle systems” 480431/2013-2. PG thanks CMAT for support by “FEDER” through the “Programa Operacional Factores de Competitividade COMPETE” and by FCT through the project PEst-C/MAT/UI0013/2011. This work has been supported by the Brazilian-French Network in Mathematics, the French Ministry of Education through the grant ANR (EDNHS) and the fellowship L’Oréal-France UNESCO “Pour les Femmes et la Science”.

Appendix 1: Fourier Analysis

(1) *Fourier transform of integrable functions*—If $f : \mathbb{R} \rightarrow \mathbb{R}$ is an integrable function, we define its Fourier transform $\mathcal{F}f : \mathbb{R} \rightarrow \mathbb{C}$ as

$$\mathcal{F}f(\xi) := \int_{\mathbb{R}} f(x)e^{2i\pi\xi x} dx, \quad \xi \in \mathbb{R}. \tag{7.1}$$

(2) *Fourier transform of square summable sequences*—If $h : \mathbb{Z} \rightarrow \mathbb{R}$ is square summable, we define its Fourier transform $\widehat{h} : \mathbb{T} \rightarrow \mathbb{C}$ in $\mathbf{L}^2(\mathbb{T})$ as

$$\widehat{h}(\theta) := \sum_{x \in \mathbb{Z}} h(x) e^{2i\pi\theta x}, \quad \theta \in \mathbb{T}. \quad (7.2)$$

(3) *Discrete Fourier transform of integrable functions*—If $g : \mathbb{R} \rightarrow \mathbb{R}$ is an integrable function, we define its Fourier transform $\mathcal{F}_n(g) : \mathbb{R} \rightarrow \mathbb{C}$ as

$$\mathcal{F}_n(g)(\xi) = \frac{1}{n} \sum_{x \in \mathbb{Z}} g\left(\frac{x}{n}\right) e^{2i\pi x \frac{\xi}{n}}, \quad \xi \in \mathbb{R}.$$

These definitions can easily be extended for d -dimensional spaces, $d \geq 1$. For each Fourier transform, we have the *Parseval–Plancherel’s identity* between suitable norms of the involved spaces, and we also can recover the initial functions for the knowledge of their Fourier transforms by the *inverse Fourier transform*.

For example, for the Fourier transform (3), the Parseval–Plancherel’s identity reads as

$$\|g\|_{2,n}^2 = \frac{1}{n} \sum_{x \in \mathbb{Z}} \left|g\left(\frac{x}{n}\right)\right|^2 = \int_{[-\frac{n}{2}, \frac{n}{2}]} |\mathcal{F}_n(g)(\xi)|^2 d\xi =: \|\mathcal{F}_n(g)\|^2.$$

The function g can be recovered from the knowledge of its Fourier transform by the inverse Fourier transform of $\mathcal{F}_n(g)$:

$$g\left(\frac{x}{n}\right) = \int_{[-\frac{n}{2}, \frac{n}{2}]} \mathcal{F}_n(g)(\xi) e^{-2i\pi x \frac{\xi}{n}} d\xi.$$

Now we give the properties that we need.

Lemma 7.1 *If $g \in \mathcal{S}(\mathbb{R})$, then for any $p \geq 1$, there exists a constant $C := C(p, f)$ such that for any $|y| \leq 1/2$,*

$$|\mathcal{F}_n(g)(ny)| \leq \frac{C}{1 + (n|y|)^p}.$$

Proof This lemma is entirely proved in [5]. □

The following result is an easy corollary of the previous lemma.

Corollary 7.2 *If $g \in \mathcal{S}(\mathbb{R})$, then, for any $p \geq 0$,*

$$\lim_{n \rightarrow \infty} \int_{[-\frac{n}{2}, \frac{n}{2}]} |\xi|^p |\mathcal{F}_n(g)(\xi) - (\mathcal{F}g)(\xi)|^2 d\xi = 0,$$

and there exists a constant $C > 0$ such that

$$\int_{[-\frac{n}{2}, \frac{n}{2}]} |\xi|^p |\mathcal{F}_n(g)(\xi)|^2 d\xi \leq C.$$

Appendix 2: Sharp Estimate of the Resolvent Norm

Hermite Polynomials

For simplicity, we assume $\beta = 1$ and we drop the index β from the notations. Let us decompose the generator S_n in the basis of multivariate Hermite polynomials. We denote by

$(h_n)_{n \geq 0}$ the sequence of standard Hermite polynomials, which are orthogonal with respect to the standard Gaussian probability measure on \mathbb{R} .

Let Σ be the set composed of configurations $\sigma = (\sigma_x)_{x \in \mathbb{Z}} \in \mathbb{N}^{\mathbb{Z}}$ such that $\sigma_x \neq 0$ only for a finite number of x . The number $\sum_{x \in \mathbb{Z}} \sigma_x$ is called the size of σ and is denoted by $|\sigma|$. Let $\Sigma_n = \{\sigma \in \Sigma ; |\sigma| = n\}$. On the set of n -tuples $\mathbf{x} := (x_1, \dots, x_n)$ of \mathbb{Z}^n , we introduce the equivalence relation $\mathbf{x} \sim \mathbf{y}$ if there exists a permutation p on $\{1, \dots, n\}$ such that $x_{p(i)} = y_i$ for all $i \in \{1, \dots, n\}$. The class of \mathbf{x} for the relation \sim is denoted by $[\mathbf{x}]$ and its cardinal by $c(\mathbf{x})$. Then the set of configurations of Σ_n can be identified with the set of n -tuples classes for \sim by the one-to-one application:

$$[\mathbf{x}] = [(x_1, \dots, x_n)] \in \mathbb{Z}^n / \sim \rightarrow \sigma^{[\mathbf{x}]} \in \Sigma_n$$

where for any $y \in \mathbb{Z}$, $(\sigma^{[\mathbf{x}]})_y = \sum_{i=1}^n \mathbf{1}_{y=x_i}$. We will identify $\sigma \in \Sigma_n$ with the occupation numbers of a configuration with n particles, and $[\mathbf{x}]$ will correspond to the positions of those n particles.

To any $\sigma \in \Sigma$, we associate the polynomial function h_σ given by

$$h_\sigma(\omega) = \prod_{x \in \mathbb{Z}} h_{\sigma_x}(\omega_x).$$

Then, the family $\{h_\sigma ; \sigma \in \Sigma\}$ forms an orthogonal basis of $\mathbf{L}^2(\mu_1)$ such that

$$\int h_\sigma(\omega) h_{\sigma'}(\omega) d\mu_1(\omega) = \delta_{\sigma=\sigma'}, \tag{7.3}$$

where δ denotes the Kronecker function, so that $\delta_{\sigma=\sigma'} = 1$ if $\sigma = \sigma'$, otherwise it is equal to zero.

A function $F : \Sigma \rightarrow \mathbb{R}$ such that $F(\sigma) = 0$ if $\sigma \notin \Sigma_n$ is called a degree n function. Thus, such a function is sometimes considered as a function defined only on Σ_n . A local function $f \in \mathbf{L}^2(\mu_1)$ whose decomposition on the orthogonal basis $\{h_\sigma ; \sigma \in \Sigma\}$ is given by $f = \sum_{\sigma} F(\sigma)h_\sigma$ is called of degree n if and only if F is of degree n . A function $F : \Sigma_n \rightarrow \mathbb{R}$ is nothing but a symmetric function $F : \mathbb{Z}^n \rightarrow \mathbb{R}$ through the identification of σ with $[\mathbf{x}]$. We denote² by $\langle \cdot, \cdot \rangle$ the scalar product on $\oplus \mathbf{L}^2(\Sigma_n)$, each Σ_n being equipped with the counting measure. Hence, if $F, G : \Sigma \rightarrow \mathbb{R}$, we have

$$\langle F, G \rangle = \sum_{n \geq 0} \sum_{\sigma \in \Sigma_n} F_n(\sigma)G_n(\sigma) = \sum_{n \geq 0} \sum_{\mathbf{x} \in \mathbb{Z}^n} \frac{1}{c(\mathbf{x})} F_n(\mathbf{x})G_n(\mathbf{x}),$$

with F_n, G_n the restrictions of F, G to Σ_n . We recall that $c(\mathbf{x})$ is the cardinal of $[\mathbf{x}]$.

If a local function $f \in \mathbf{L}^2(\mu_1)$ is written in the form $f = \sum_{\sigma \in \Sigma} F(\sigma)h_\sigma$ then we have

$$(\mathcal{S}_n f)(\omega) = \sum_{\sigma \in \Sigma} (\mathcal{S}_n F)(\sigma)h_\sigma(\omega)$$

with

$$(\mathcal{S} f)(\sigma) = \lambda \sum_{x \in \mathbb{Z}} (F(\sigma^{x,x+1}) - F(\sigma)) - \gamma_n \sum_{x \in \mathbb{Z}} W_x(\sigma)F(\sigma),$$

where $\sigma^{x,x+1}$ is obtained from σ by exchanging the occupation numbers σ_x and σ_{x+1} and $W_x(\sigma)$ is defined by $W_x(\sigma) = 1 - (-1)^{\sigma_x} \geq 0$. The formula for $W_x(\sigma)$ is a direct consequence of the fact that h_n is even (resp. odd) if n is even (resp. odd).

² With some abuse of notations.

$H_{1,z}$ and $H_{-1,z}$ Norms

For any $z > 0$ and any $f \in L^2(\mu_1)$ we define the $H_{\pm 1,z}$ -norm of f by

$$\|f\|_{\pm 1,z} = \left[\langle f(z - \mathcal{S}_n)^{\pm 1} f \rangle_1 \right]^{1/2}.$$

We have

$$\|f\|_{1,z}^2 = z \langle f^2 \rangle_1 + \mathcal{D}(f)$$

where $\mathcal{D}(f)$ is the Dirichlet form of f defined by $\mathcal{D}(f) = \langle f(-\mathcal{S}_n f) \rangle_1$.

If f has the decomposition $f = \sum_{\sigma \in \Sigma} F(\sigma)h_\sigma$ then

$$\mathcal{D}(f) = \frac{1}{2} \sum_{x \in \mathbb{Z}} \sum_{\sigma \in \Sigma} (F(\sigma^{x,x+1}) - F(\sigma))^2 + \gamma_n \sum_{\sigma \in \Sigma, x \in \mathbb{Z}} W_x(\sigma) F^2(\sigma). \tag{7.4}$$

Let $\Delta_+ = \{(x, y) \in \mathbb{Z}^2; y \geq x + 1\}$, $\Delta_- = \{(x, y) \in \mathbb{Z}^2; y \leq x - 1\}$ and $\Delta_0 = \{(x, x); x \in \mathbb{Z}\}$.

We denote by \mathbb{D} the Dirichlet form of a symmetric simple random walk on \mathbb{Z}^2 where jumps from Δ_\pm to Δ_0 and from Δ_0 to Δ_\pm have been suppressed and jumps from $(x, x) \in \Delta_0$ to $(x \pm 1, x \pm 1) \in \Delta_0$ have been added, i.e.

$$\mathbb{D}(f) = \frac{1}{2} \sum_{|\mathbf{e}|=1} \sum_{\mathbf{x} \in \Delta_\pm, \mathbf{x}+\mathbf{e} \in \Delta_\pm} (F(\mathbf{x} + \mathbf{e}) - F(\mathbf{x}))^2 + \frac{1}{2} \sum_{\mathbf{x} \in \Delta_0} (F(\mathbf{x} \pm (1, 1)) - F(\mathbf{x}))^2,$$

where $F : \mathbb{Z}^2 \rightarrow \mathbb{R}$ is a symmetric function such that $\sum_{\mathbf{x} \in \mathbb{Z}^2} F^2(\mathbf{x}) < \infty$.

Lemma 7.3 *Let $f = \sum_{\sigma \in \Sigma_2} F(\sigma)h_\sigma$ be a local function of degree 2. There exists a positive constant C , independent of f and n , such that*

$$C^{-1} \left[\mathbb{D}(f) + \gamma_n \sum_{\mathbf{x} \notin \Delta_0} F^2(\mathbf{x}) \right] \leq \mathcal{D}(f) \leq C \left[\mathbb{D}(f) + \gamma_n \sum_{\mathbf{x} \notin \Delta_0} F^2(\mathbf{x}) \right].$$

Proof Straightforward. □

We denote by Σ_2^0 the set of configurations σ of Σ_2 such that $\sigma = 2\delta_x, x \in \mathbb{Z}$, and Σ_2^\pm the complementary set of Σ_2^0 in Σ_2 , i.e. the set of configurations $\sigma \in \Sigma_2$ such that $\sigma = \delta_x + \delta_y, y \neq x \in \mathbb{Z}$.

Observe that the function φ defined by (4.13) is a function of degree 2 with a decomposition in the form $\varphi = \sum_{\sigma \in \Sigma_2} \Phi(\sigma)h_\sigma$ which satisfies $\Phi(\sigma) = 0$ if $\sigma \in \Sigma_2^0$. We have that

$$\left\langle \varphi(z - \mathcal{S}_n)^{-1} \varphi \right\rangle_1 = \sup_g \left\{ 2\langle \varphi g \rangle_1 - z\langle g^2 \rangle_1 - \mathcal{D}(g) \right\}$$

where the supremum is taken over local functions $g \in L^2(\mu_1)$. Decompose g appearing in this variational formula as $g = \sum_{\sigma} G(\sigma)h_\sigma$. Recall that $\{h_\sigma; \sigma \in \Sigma\}$ are orthogonal, φ is a degree 2 function such that $\Phi(\sigma) = 0$ for any $\sigma \notin \Sigma_2^\pm$ and recall (7.4) for the Dirichlet form $\mathcal{D}(g)$. Thus, we can restrict this supremum over degree 2 functions g such that $G(\sigma) = 0$ if $\sigma \in \Sigma_2^0$. Then, by Lemma 7.3, we have

$$\begin{aligned} \left\langle \varphi (z - \mathcal{S}_n)^{-1} \varphi \right\rangle_1 \leq & C \sup_G \left\{ \sum_{x \neq y} \Phi(x, y) G(x, y) - (z + \gamma_n) \sum_{\substack{(x, y) \in \mathbb{Z}^2 \\ x \neq y}} G^2(x, y) \right. \\ & \left. - C' \sum_{|\mathbf{e}|=1} \sum_{\substack{(x, y) \in \Delta^\pm \\ (x, y) + \mathbf{e} \in \Delta^\pm}} \left(G((x, y) + \mathbf{e}) - G(x, y) \right)^2 \right\} \end{aligned}$$

where C, C' are positive constants, $\Delta^\pm = \{(x, y) \in \mathbb{Z}^2; x \neq y\}$ and as usual we identify the functions defined on Σ_n with symmetric functions defined on \mathbb{Z}^n .

In order to get rid of the geometric constraints appearing in the last term of the variational formula, for any symmetric function G defined on the set Δ^\pm , we denote by \tilde{G} its extension to \mathbb{Z}^2 defined by

$$\tilde{G}(x, y) = G(x, y) \text{ if } x \neq y, \quad \tilde{G}(x, x) = \frac{1}{4} \sum_{|\mathbf{e}|=1} G((x, x) + \mathbf{e}).$$

It is trivial that there exists a constant $C > 0$ such that

$$\sum_{(x, y) \in \mathbb{Z}^2} \tilde{G}^2(x, y) \leq C \sum_{\substack{(x, y) \in \mathbb{Z}^2 \\ x \neq y}} G^2(x, y),$$

and

$$\sum_{|\mathbf{e}|=1} \sum_{(x, y) \in \mathbb{Z}^2} \left(\tilde{G}((x, y) + \mathbf{e}) - \tilde{G}(x, y) \right)^2 \leq C \sum_{|\mathbf{e}|=1} \sum_{\substack{(x, y) \in \Delta^\pm \\ (x, y) + \mathbf{e} \in \Delta^\pm}} \left(G((x, y) + \mathbf{e}) - G(x, y) \right)^2.$$

Thus, we have

$$\begin{aligned} \left\langle \varphi (z - \mathcal{S}_n)^{-1} \varphi \right\rangle_1 \leq & C_0 \sup_G \left\{ \sum_{(x, y) \in \mathbb{Z}^2} \Phi(x, y) G(x, y) - C_1 (z + \gamma_n) \sum_{(x, y) \in \mathbb{Z}^2} G^2(x, y) \right. \\ & \left. - C_2 \sum_{|\mathbf{e}|=1} \sum_{(x, y) \in \mathbb{Z}^2} \left(G((x, y) + \mathbf{e}) - G(x, y) \right)^2 \right\} \end{aligned}$$

where the supremum is now taken over all symmetric local functions $G : \mathbb{Z}^2 \rightarrow \mathbb{R}$. Notice that the last variational formula is equal to the resolvent norm, for a simple symmetric two dimensional random walk, of the function Φ .

Then by using Fourier transform, it is easy to see (see [6]) that the last supremum is equal to

$$\frac{C_0}{4} \int_{[0, 1]^2} \frac{|\widehat{\Phi}(\mathbf{k})|^2}{C_1 [z + \gamma_n] + 4C_2 \sum_{i=1}^2 \sin^2(\pi k_i)} d\mathbf{k} \tag{7.5}$$

where the Fourier transform $\widehat{\Phi}$ of Φ is given by

$$\widehat{\Phi}(\mathbf{k}) = \sum_{(x, y) \in \mathbb{Z}^2} \Phi(x, y) e^{2i\pi(k_1 x + k_2 y)}, \quad \mathbf{k} = (k_1, k_2) \in [0, 1]^2.$$

Thus we have reduced the problem to estimate the behavior w.r.t. n of the integral (7.5) with $z = 1/(tn^a)$. Since the constants t, C_0, C_1, C_2 do not play any role, we fix them equal to 1. The function φ can be rewritten as

$$\begin{aligned} \varphi(\omega) &= \sum_{v>u} \left\{ \sum_{z \in \mathbb{Z}} (\nabla_n f)\left(\frac{z}{n}\right) [\psi_{|v-u|}(u-z-1) - \psi_{|v-u|}(u-z)] \right\} \omega_u \omega_v \\ &= \sum_{u,v} \Phi(u, v) \omega_u \omega_v \end{aligned}$$

with the symmetric function Φ given by

$$\Phi(u, v) = \frac{1_{u \neq v}}{2} \sum_{z \in \mathbb{Z}} (\nabla_n f)\left(\frac{z}{n}\right) [\psi_{|v-u|}(u \wedge v - z - 1) - \psi_{|v-u|}(u \wedge v - z)].$$

Its Fourier transform is given by

$$\begin{aligned} \widehat{\Phi}(\mathbf{k}) &= \sum_{u \in \mathbb{Z}} \sum_{j=1}^{\infty} \Phi(u, u+j) e^{2i\pi(k_1 u + k_2(u+j))} + \sum_{u \in \mathbb{Z}} \sum_{j=1}^{\infty} \Phi(u+j, u) e^{2i\pi(k_1(u+j) + k_2 u)} \\ &= -\frac{1}{2} \mathcal{F}_n(\Delta_n f)(n(k_1 + k_2)) \sum_{j=1}^{\infty} \left(e^{2i\pi k_1 j} + e^{2i\pi k_2 j} \right) \widehat{\psi}_j(k_1 + k_2). \end{aligned}$$

Since $|\mathcal{F}_n(\Delta_n f)(\xi)| \leq C_0$ for a suitable constant $C_0 > 0$ independent of n and ξ , it follows that there exist constants $C, C' > 0$ such that

$$\begin{aligned} |\widehat{\Phi}(\mathbf{k})|^2 &\leq C \left| \sum_{j=1}^{\infty} \left(e^{2i\pi k_1 j} + e^{2i\pi k_2 j} \right) \widehat{\psi}_j(k_1 + k_2) \right|^2 \\ &= C \left| \sum_{j=2}^{\infty} \left\{ \left(e^{2i\pi k_1 j - 2i\pi k_2} + e^{2i\pi k_2 j - 2i\pi k_1} + e^{2i\pi k_1(j-1)} + e^{2i\pi k_2(j-1)} \right) \widehat{\rho}_j(k_1 + k_2) \right\} \right. \\ &\quad \left. + \left(e^{2i\pi(k_1 - k_2)} + e^{2i\pi(k_2 - k_1)} \right) \widehat{\rho}_1(k_1 + k_2) \right|^2 \\ &\leq C' \frac{|\widehat{\rho}_1(k_1 + k_2)|^2}{(1 - |X(k_1 + k_2)|)^2}. \end{aligned}$$

To get the last inequality, we used the explicit form of $\widehat{\rho}_j$ and the fact that

$$|1 - X(k_1 + k_2)w| \geq 1 - |X(k_1 + k_2)| > 0$$

for any complex number w of modulus one. Then, from Lemma 4.1 it follows that, for n sufficiently large,

$$\frac{|\widehat{\rho}_1(\theta)|^2}{(1 - |X(\theta)|)^2} \leq \frac{C}{\gamma_n [\gamma_n + \sin^2(\pi\theta)]}.$$

Estimate of an Integral

It remains to study the behavior of the integral

$$I_n := \int_{[0,1]^2} d\mathbf{k} \left\{ \frac{1}{(n^{-a} + \gamma_n) + \sin^2(\pi k_1) + \sin^2(\pi k_2)} \times \frac{1}{\gamma_n [\gamma_n + \sin^2(\pi(k_1 + k_2))]} \right\}.$$

The leading term in this integral is provided when

- (1) either \mathbf{k} is an extremal point of $[0, 1]^2$, i.e. $(0, 0), (0, 1), (1, 0), (1, 1)$,

(2) or \mathbf{k} belongs to the diagonal $\{k_1 + k_2 = 1\}$, and \mathbf{k} is not one of the previous four points.

We first consider the case (1). By periodicity, we can assume that the extremal point is $(0, 0)$. We also have that $a > b$, and then $\gamma_n \gg n^{-a}$. We perform a Taylor expansion and forget about the constants. We are reduced to compute the order, as $n \rightarrow \infty$, of

$$\int_{[0,1]^2} \frac{d\mathbf{k}}{\gamma_n [\gamma_n + k_1^2 + k_2^2] [\gamma_n + (k_1 + k_2)^2]} \approx \int_0^1 \frac{r dr}{\gamma_n [\gamma_n + r^2]^2} \approx n^{2b}.$$

The first estimate comes after a change into polar coordinates, and the last one is deduced from another change of variables:

$$\int_0^1 \frac{r dr}{\gamma_n [\gamma_n + r^2]^2} = \int_0^{1/\sqrt{\gamma_n}} \frac{du}{\gamma_n^2 [1 + u^2]^2} \approx n^{2b}.$$

In the case (2), by the same argument we are reduced to investigate the behavior of

$$\int_0^1 \frac{r dr}{\gamma_n^2 [\gamma_n + r^2]}$$

which is of the same order n^{2b} .

When \mathbf{k} is not close to one of these points, we can bound by below $\sin^2(\pi k_1) + \sin^2(\pi k_2)$ and $\sin^2(\pi(k_1 + k_2))$ by a strictly positive constant independent of n and then show that the corresponding integral gives a smaller contribution. Finally, I_n is of order n^{2b} .

Appendix 3: Computations in the Case $b > 1$

Computations Involving the Generator

In this subsection, we explain how to obtain the differential equations in Proposition 5.2. Let $f : \mathbb{Z} \rightarrow \mathbb{R}$ be a function of finite support and let $\mathcal{E}(f) : \Omega \rightarrow \mathbb{R}$ be defined as

$$\mathcal{E}(f) = \sum_{x \in \mathbb{Z}} f(x) \omega_x^2.$$

A simple computation shows that

$$\mathcal{S}_n \mathcal{E}(f) = \sum_{x \in \mathbb{Z}} \Delta f(x) \omega_x^2,$$

where $\Delta f(x) = f(x + 1) + f(x - 1) - 2f(x)$ is the discrete Laplacian on \mathbb{Z} . On the other hand

$$\mathcal{A} \mathcal{E}(f) = -2 \sum_{x \in \mathbb{Z}} \nabla f(x) \omega_x \omega_{x+1},$$

where $\nabla f(x) = f(x + 1) - f(x)$ is the discrete right-derivative in \mathbb{Z} . Let $f : \mathbb{Z}^2 \rightarrow \mathbb{R}$ be a symmetric function of finite support, and let $Q(f) : \Omega \rightarrow \mathbb{R}$ be defined as

$$Q(f) = \sum_{\substack{x, y \in \mathbb{Z} \\ x \neq y}} f(x, y) \omega_x \omega_y.$$

Define $\Delta f : \mathbb{Z}^2 \rightarrow \mathbb{R}$ as

$$\Delta f(x, y) = f(x + 1, y) + f(x - 1, y) + f(x, y + 1) + f(x, y - 1) - 4f(x, y)$$

for any $x, y \in \mathbb{Z}$ and $Af : \mathbb{Z}^2 \rightarrow \mathbb{R}$ by

$$Af(x, y) = f(x - 1, y) + f(x, y - 1) - f(x + 1, y) - f(x, y + 1)$$

for any $x, y \in \mathbb{Z}$. Notice that Δf is the discrete Laplacian on the lattice \mathbb{Z}^2 and Af is a possible definition of the discrete derivative of f in the direction $(-2, -2)$. We are using the same symbol Δ for the one-dimensional and two-dimensional, discrete Laplacian, but from the context it will be clear which operator is used. We have that

$$\begin{aligned} \mathcal{S}_n Q(f) &= \sum_{\substack{x, y \in \mathbb{Z} \\ x \neq y}} [(\Delta f)(x, y) - 4\gamma_n f(x, y)] \omega_x \omega_y \\ &\quad + 2 \sum_{x \in \mathbb{Z}} \left\{ [f(x, x + 1) - f(x, x)] + [f(x, x + 1) - f(x + 1, x + 1)] \right\} \omega_x \omega_{x+1} \\ &= Q(\Delta f - 2\gamma_n \text{Id}) \\ &\quad + 2 \sum_{x \in \mathbb{Z}} \left\{ [f(x, x + 1) - f(x, x)] + [f(x, x + 1) - f(x + 1, x + 1)] \right\} \omega_x \omega_{x+1}. \end{aligned}$$

Similarly, we have that

$$\begin{aligned} A Q(f) &= Q(Af) + 2 \sum_{x \in \mathbb{Z}} \left\{ [f(x - 1, x) - f(x, x + 1)] \omega_x^2 \right. \\ &\quad \left. - [f(x, x) - f(x + 1, x + 1)] \omega_x \omega_{x+1} \right\}. \end{aligned}$$

It follows that

$$\mathcal{L}_n Q(f) = Q((\Delta + A - 4\gamma_n \text{Id})f) + D(f), \tag{7.6}$$

where the diagonal term $D(f)$ is given by

$$\begin{aligned} D(f) &= 2 \sum_{x \in \mathbb{Z}} (\omega_x^2 - \beta^{-1}) (f(x - 1, x) - f(x, x + 1)) \\ &\quad + 4 \sum_{x \in \mathbb{Z}} (f(x, x + 1) - f(x, x)) \omega_x \omega_{x+1}. \end{aligned}$$

The normalization constant β^{-1} can be added for free because $f(x, x + 1) - f(x - 1, x)$ is a mean-zero function. The diagonal term will be of capital importance, in particular the one involving ω_x^2 . Notice that the operators $f \mapsto Q(f)$, $f \mapsto \mathcal{L}_n Q(f)$ are continuous maps from $\ell^2(\mathbb{Z}^2)$ to $L^2(\mu_\beta)$. Therefore, an approximation procedure shows that the identities above hold for any $f \in \ell^2(\mathbb{Z}^2)$.

Recalling the definition of \mathcal{S}_t^n , and assuming $\beta = 1$, we deduce (5.3), that is

$$\begin{aligned} \frac{d}{dt} \mathcal{S}_t^n(f) &= \frac{1}{2n} \mathbb{E} \left[\sum_{x \in \mathbb{Z}} g\left(\frac{x}{n}\right) (\omega_x^2(0) - 1) n^{3/2} \mathcal{L}_n(\mathcal{E}(f)) \right] \\ &= -2Q_t^n(\nabla_n f \otimes \delta) + \mathcal{S}_t^n\left(\frac{1}{\sqrt{n}} \Delta_n f\right). \end{aligned}$$

Estimates on the Poisson Equation

Let us define the functions

$$\Lambda(x, y) = 4 [\sin^2(\pi x) + \sin^2(\pi y)], \quad \Omega(x, y) = 2 [\sin(2\pi x) + \sin(2\pi y)]. \quad (7.7)$$

Several times we will use the following change of variable property proved in [5].

Lemma 7.4 *Let $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ be a n -periodic function in each direction of \mathbb{R}^2 . Then we have*

$$\int \int_{[-\frac{n}{2}, \frac{n}{2}]^2} f(k, \ell) dk d\ell = \int \int_{[-\frac{n}{2}, \frac{n}{2}]^2} f(\xi - \ell, \ell) d\xi d\ell.$$

Proof of Lemma 5.5

The proof is similar to the one given in [5]. The Fourier transform of h_n is given by

$$\mathcal{F}_n(h_n)(k, \ell) = \frac{1}{2\sqrt{n}} \frac{i\Omega(\frac{k}{n}, \frac{\ell}{n})\mathcal{F}_n(f)(k + \ell)}{(\Lambda + 4\gamma_n - i\Omega)(\frac{k}{n}, \frac{\ell}{n})}, \quad (7.8)$$

where Λ and Ω are defined in (7.7). Observe first that

$$i\Omega\left(\frac{\xi - \ell}{n}, \frac{\ell}{n}\right) = e^{\frac{2i\pi\ell}{n}} \left(1 - e^{-\frac{2i\pi\xi}{n}}\right) - e^{-\frac{2i\pi\ell}{n}} \left(1 - e^{\frac{2i\pi\xi}{n}}\right)$$

so that

$$\left| i\Omega\left(\frac{\xi - \ell}{n}, \frac{\ell}{n}\right) \right|^2 \leq 4 \left| 1 - e^{\frac{2i\pi\xi}{n}} \right|^2 = 16 \sin^2\left(\frac{\pi\xi}{n}\right). \quad (7.9)$$

Then, by the Parseval–Plancherel’s relation and by Lemma 7.4 we have

$$\begin{aligned} \|h_n\|_{2,n}^2 &= \iint_{[-\frac{n}{2}, \frac{n}{2}]^2} |\mathcal{F}_n(h_n)(k, \ell)|^2 dk d\ell \\ &= \frac{1}{4n} \iint_{[-\frac{n}{2}, \frac{n}{2}]^2} \frac{\Omega^2(\frac{k}{n}, \frac{\ell}{n}) |\mathcal{F}_n(f)(k + \ell)|^2}{(\Lambda(\frac{k}{n}, \frac{\ell}{n}) + 4\gamma_n)^2 + \Omega^2(\frac{k}{n}, \frac{\ell}{n})} dk d\ell \\ &\leq \frac{1}{n} \int_{-n/2}^{n/2} \left| 1 - e^{\frac{2i\pi\xi}{n}} \right|^2 |\mathcal{F}_n(f)(\xi)|^2 \left[\int_{-n/2}^{n/2} \frac{d\ell}{\Lambda^2(\frac{\xi - \ell}{n}, \frac{\ell}{n}) + \Omega^2(\frac{\xi - \ell}{n}, \frac{\ell}{n})} \right] d\xi \\ &= 4n \int_{-1/2}^{1/2} \sin^2(\pi y) |\mathcal{F}_n(f)(ny)|^2 W(y) dy, \end{aligned}$$

where for the last equality we performed the changes of variables $y = \frac{\xi}{n}$ and $x = \frac{\ell}{n}$ and forget the positive term $4\gamma_n$. The function W is defined by

$$W(y) = \int_{-1/2}^{1/2} \frac{dx}{\Lambda^2(y - x, x) + \Omega^2(y - x, x)}. \quad (7.10)$$

It is proved in [5, Lemma F.5] that $W(y) \leq C|y|^{-3/2}$ on $[-\frac{1}{2}, \frac{1}{2}]$. Hence, we get, by using the second part of Lemma 7.1 with $p = 3$ and the elementary inequality $\sin^2(\pi y) \leq (\pi y)^2$, that

$$\iint_{[-\frac{n}{2}, \frac{n}{2}]^2} |\mathcal{F}_n(h_n)(k, \ell)|^2 dk d\ell \leq C'n \int_{-1/2}^{1/2} \frac{|y|^{1/2}}{1 + (n|y|)^3} dy = O(n^{-1/2}).$$

We have proved the first part of Lemma 5.5, that is (5.10). We turn now to (5.11). We denote by G_n the 1-periodic function defined by

$$G_n(y) = \frac{1}{4} \int_{-1/2}^{1/2} \frac{\Omega^2(y - z, z)}{4\gamma_n + \Lambda(y - z, z) - i \Omega(y - z, z)} dz. \tag{7.11}$$

As $y \rightarrow 0$, the function G_n is close (in a sense defined below) to the function G_0 given by

$$G_0(y) = \frac{1}{2} |\pi y|^{3/2} (1 + i \operatorname{sgn}(y)). \tag{7.12}$$

In fact we show in Lemma 7.6 that there exists one constant $C > 0$ such that for any $|y| \leq 1/2$ and for all $n \in \mathbb{N}_0$,

$$|G_n(y) - G_0(y)| \leq C [\sin^2(\pi y) + \gamma_n^2 |\sin(\pi y)|^{-1/2} + \gamma_n |\sin(\pi y)|^{1/2}]. \tag{7.13}$$

Recall $\mathcal{F}f$ the (continuous) Fourier transform of f from (7.1) and denote by $q := q(f) : \mathbb{R} \rightarrow \mathbb{R}$ the function defined by

$$q(x) = \int_{-\infty}^{\infty} e^{-2i\pi xy} G_0(y) \mathcal{F}f(y) dy$$

which coincides with $-\frac{1}{4} \mathbb{L}f(x)$. Let $q_n : \frac{1}{n}\mathbb{Z} \rightarrow \mathbb{R}$ the function defined by $q_n(\frac{x}{n}) = \mathcal{D}_n h_n(\frac{x}{n})$. Then, the proof of (5.11) reduces to the following

Lemma 7.5

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{x \in \mathbb{Z}} [q(\frac{x}{n}) - q_n(\frac{x}{n})]^2 = 0.$$

Proof Since $\mathcal{F}_n(h_n)$ is a symmetric function we can easily see that

$$\mathcal{F}_n(q_n)(\xi) = -\frac{i}{2} \sum_{x \in \mathbb{Z}} e^{\frac{2i\pi \xi x}{n}} \iint_{[-\frac{n}{2}, \frac{n}{2}]^2} e^{-\frac{2i\pi(k+\ell)x}{n}} \Omega(\frac{k}{n}, \frac{\ell}{n}) \mathcal{F}_n(h_n)(k, \ell) dk d\ell.$$

We use now Lemma 7.4 and the inverse Fourier transform relation to get

$$\mathcal{F}_n(q_n)(\xi) = -\frac{in}{2} \int_{-n/2}^{n/2} \Omega(\frac{\xi-\ell}{n}, \frac{\ell}{n}) \mathcal{F}_n(h_n)(\xi - \ell, \ell) d\ell.$$

By the explicit expression (7.8) of $\mathcal{F}_n(h_n)$ we obtain that

$$\mathcal{F}_n(q_n)(\xi) = \frac{\sqrt{n}}{4} \left[\int_{-n/2}^{n/2} \frac{\Omega^2(\frac{\xi-\ell}{n}, \frac{\ell}{n})}{4\gamma_n + \Lambda(\frac{\xi-\ell}{n}, \frac{\ell}{n}) - i \Omega(\frac{\xi-\ell}{n}, \frac{\ell}{n})} d\ell \right] \mathcal{F}_n(f)(\xi).$$

Again by the inverse Fourier transform we get that

$$q_n(\frac{x}{n}) = \int_{-n/2}^{n/2} e^{-\frac{2i\pi \xi x}{n}} n^{3/2} G_n(\frac{\xi}{n}) \mathcal{F}_n(f)(\xi) d\xi.$$

Then we have

$$\begin{aligned} q(\frac{x}{n}) - q_n(\frac{x}{n}) &= \int_{|\xi| \geq n/2} e^{-\frac{2i\pi \xi x}{n}} G_0(\xi) \mathcal{F}f(\xi) d\xi \\ &+ \int_{|\xi| \leq n/2} e^{-\frac{2i\pi \xi x}{n}} G_0(\xi) [\mathcal{F}f(\xi) - \mathcal{F}_n(f)(\xi)] d\xi \\ &+ n^{3/2} \int_{|\xi| \leq n/2} e^{-\frac{2i\pi \xi x}{n}} (G_0 - G_n)(\frac{\xi}{n}) \mathcal{F}_n(f)(\xi) d\xi. \end{aligned}$$

Above we have used the fact that $n^{3/2}G_0(\frac{\xi}{n}) = G_0(\xi)$. Then we use the triangular inequality and the Parseval–Plancherel’s Theorem in the two last terms of the RHS to write

$$\begin{aligned} \frac{1}{n} \sum_{x \in \mathbb{Z}} [q(\frac{x}{n}) - q_n(\frac{x}{n})]^2 &\leq \frac{1}{n} \sum_{x \in \mathbb{Z}} \left| \int_{|\xi| \geq n/2} e^{-\frac{2i\pi\xi x}{n}} G_0(\xi) (\mathcal{F}f)(\xi) d\xi \right|^2 \\ &\quad + \int_{|\xi| \leq n/2} |G_0(\xi)[\mathcal{F}f(\xi) - \mathcal{F}_n(f)(\xi)]|^2 d\xi \\ &\quad + n^3 \int_{|\xi| \leq n/2} |(G_0 - G_n)(\frac{\xi}{n})\mathcal{F}_n(f)(\xi)|^2 d\xi \\ &= (I) + (II) + (III). \end{aligned} \tag{7.14}$$

We refer to [5] for a proof of the following fact: both terms (I) and (II) give a trivial contribution in (7.14). The first one is estimated by performing an integration by parts and using the fact that for $f \in \mathcal{S}(\mathbb{R})$ its Fourier transform $\mathcal{F}f$ is in $\mathcal{S}(\mathbb{R})$, and that G_0 and G'_0 grow polynomially. For the second one, a change of variables $\xi = \frac{y}{n}$ and the fact that $f \in \mathcal{S}(\mathbb{R})$ together with Lemma 7.1 permit to conclude.

The contribution of (III) is estimated by using (7.20) and is different from [5]. Recall the trivial inequality $|\sin(\pi y)| \leq |\pi y|$, for $|y| \leq 1/2$. The first term of the RHS of (7.20) gives the upper bound

$$\frac{C}{n} \int_{|\xi| \leq n/2} |\xi|^4 |\mathcal{F}_n(f)(\xi)|^2 d\xi = C \int_{-1/2}^{1/2} n^4 |z|^4 |\mathcal{F}_n(f)(nz)|^2 dz$$

which goes to 0, as $n \rightarrow \infty$, by Lemma 7.1 applied with $p = 2$. We now deal with the last term of the RHS of (7.20), which gives the upper bound

$$Cn^2 \gamma_n^2 \int_{|\xi| \leq n/2} |\xi| |\mathcal{F}_n(f)(\xi)|^2 d\xi = C (\gamma_n^2 n^3) \int_{-1/2}^{1/2} n|z| |\mathcal{F}_n(f)(nz)|^2 dz,$$

which goes to 0, as $n \rightarrow \infty$. Indeed, this is a consequence of Lemma 7.1, and the fact that $\gamma_n^2 n^2$ goes to 0 (as long as $b > 1$). The second term gives

$$Cn^3 \gamma_n^4 \int_{|\xi| \leq n/2} \frac{|\mathcal{F}_n(f)(\xi)|^2}{|\sin(\frac{\pi\xi}{n})|} d\xi = Cn^4 \gamma_n^4 \int_{-1/2}^{1/2} \frac{|\mathcal{F}_n(f)(nz)|^2}{|\sin(\pi z)|} dz.$$

Again, since $z \mapsto |\sin(\pi z)|^{-1}$ is integrable on $[-\frac{1}{2}, \frac{1}{2}]$, and thanks to Lemma 7.1, this bound goes to 0 as soon as $\gamma_n^4 n^4$ goes to 0, which is automatically satisfied if $b > 1$. \square

Proof of Lemma 5.6

The proof is similar to the proof given in [5]. Let $w_n : \frac{1}{n}\mathbb{Z} \rightarrow \mathbb{R}$ be defined by

$$w_n(\frac{x}{n}) = h_n(\frac{x}{n}, \frac{x+1}{n}) - h_n(\frac{x}{n}, \frac{x}{n})$$

and observe that

$$\frac{1}{\sqrt{n}} \widetilde{\mathcal{D}}_n h_n(\frac{x}{n}, \frac{y}{n}) = n^{3/2} \begin{cases} w_n(\frac{x}{n}); & y = x + 1, \\ w_n(\frac{x-1}{n}); & y = x - 1, \\ 0; & \text{otherwise.} \end{cases}$$

The Fourier transform $\mathcal{F}_n(v_n)$ is thus given by

$$\mathcal{F}_n(v_n)(k, \ell) = -\frac{1}{n} \frac{e^{2i\pi \frac{k}{n}} + e^{2i\pi \frac{\ell}{n}}}{(\Lambda + 4\gamma_n - i\Omega)\left(\frac{k}{n}, \frac{\ell}{n}\right)} \mathcal{F}_n(w_n)(k + \ell). \quad (7.15)$$

By using Lemma 7.4, we have that the Fourier transform of w_n is given by

$$\mathcal{F}_n(w_n)(\xi) = \int_{-n/2}^{n/2} \mathcal{F}_n(h_n)(\xi - \ell, \ell) \left\{ e^{-\frac{2i\pi \ell}{n}} - 1 \right\} d\ell.$$

By (7.8) we get

$$\mathcal{F}_n(w_n)(\xi) = -\frac{\sqrt{n}}{2} I_n\left(\frac{\xi}{n}\right) \mathcal{F}_n(f)(\xi), \quad (7.16)$$

where the function I_n is defined by

$$I_n(y) = \int_{-1/2}^{1/2} \left(\frac{i\Omega R}{\Lambda + 4\gamma_n - i\Omega} \right) (y - x, x) dx. \quad (7.17)$$

Proof of (5.15). As in [5], we can easily get

$$\|v_n\|_{2,n}^2 \leq Cn \int_{-1/2}^{1/2} |\mathcal{F}_n(f)(ny)|^2 |I_n(y)|^2 W(y) dy$$

by using Lemma 7.1. Then, from Lemma 7.7 and since $W(y) \leq C|y|^{-3/2}$ (see [5], Lemma F.5), we get that

$$\begin{aligned} \|v_n\|_{2,n}^2 &\leq Cn \int_{-1/2}^{1/2} |\mathcal{F}_n(f)(ny)|^2 |\sin(\pi y)|^{3/2} dy \leq C \int_{-1/2}^{1/2} \frac{|y|^{3/2}}{1 + |ny|^p} dy \\ &= \frac{C}{n^{3/2}} \int_{-n/2}^{n/2} \frac{|z|^{3/2}}{1 + |z|^p} dz, \end{aligned}$$

which goes to 0, as soon as p is chosen bigger than 3.

Proof of (5.16). Following [5], straightforward computations lead to

$$\mathcal{F}_n(\mathcal{D}_n v_n)(\xi) = -n \left(1 - e^{\frac{2i\pi \xi}{n}} \right) \mathcal{F}_n(w_n)(\xi) J_n\left(\frac{\xi}{n}\right),$$

where J_n is given by

$$J_n(y) = \int_{-1/2}^{1/2} \frac{1 + e^{2i\pi(y-2x)}}{(\Lambda + 4\gamma_n - i\Omega)(y - x, x)} dx. \quad (7.18)$$

Now, by using (7.16) we finally get that

$$\mathcal{F}_n(\mathcal{D}_n v_n)(\xi) = \frac{n^{3/2}}{2} \left(1 - e^{\frac{2i\pi \xi}{n}} \right) \mathcal{F}_n(f)(\xi) I_n\left(\frac{\xi}{n}\right) J_n\left(\frac{\xi}{n}\right),$$

where I_n is defined by (7.17).

By the Parseval–Plancherel’s relation we have to prove that

$$\begin{aligned} & n^3 \int_{-n/2}^{n/2} \sin^2\left(\pi \frac{\xi}{n}\right) |\mathcal{F}_n(f)(\xi)|^2 |I_n\left(\frac{\xi}{n}\right)|^2 |J_n\left(\frac{\xi}{n}\right)|^2 d\xi \\ &= n^4 \int_{-1/2}^{1/2} \sin^2(\pi y) |I_n(y)|^2 |J_n(y)|^2 |\mathcal{F}_n(f)(ny)|^2 dy \end{aligned}$$

vanishes, as $n \rightarrow \infty$. By Lemmas 7.1 and 7.7, this is equivalent to show that the following term goes to 0, as $n \rightarrow \infty$:

$$n^4 \int_{-1/2}^{1/2} \frac{|y|^4}{1 + |ny|^p} dy = \frac{1}{n} \int_{-n/2}^{n/2} \frac{|z|^4}{1 + |z|^p} dz.$$

But, for p bigger than 5, this term goes to 0, as $n \rightarrow \infty$.

Proof of (5.17).

Let $\theta_n : \frac{1}{n}\mathbb{Z} \rightarrow \mathbb{R}$ be defined by

$$\theta_n\left(\frac{x}{n}\right) = v_n\left(\frac{x}{n}, \frac{x+1}{n}\right) - v_n\left(\frac{x}{n}, \frac{x}{n}\right)$$

and observe that

$$\frac{1}{\sqrt{n}} \tilde{\mathcal{D}}_n v_n\left(\frac{x}{n}, \frac{y}{n}\right) = n^{3/2} \begin{cases} \theta_n\left(\frac{x}{n}\right); & y = x + 1, \\ \theta_n\left(\frac{x-1}{n}\right); & y = x - 1, \\ 0; & \text{otherwise.} \end{cases}$$

We have to show that

$$\lim_{n \rightarrow \infty} n \sum_{x \in \mathbb{Z}} \theta_n^2(x) = 0$$

which is equivalent, by the Parseval–Plancherel’s, relation to show that

$$\lim_{n \rightarrow +\infty} n^2 \int_{-n/2}^{n/2} \mathcal{F}_n(\theta_n)(\xi) d\xi = 0.$$

By using Lemma 7.4, we have that the Fourier transform of θ_n is given by

$$\mathcal{F}_n(\theta_n)(\xi) = \int_{-n/2}^{n/2} \mathcal{F}_n(v_n)(\xi - \ell, \ell) \left\{ e^{-\frac{2i\pi\ell}{n}} - 1 \right\} d\ell.$$

Performing a change of variables and using (7.15) and (7.16), we get that

$$\mathcal{F}_n(\theta_n)(\xi) = \frac{\sqrt{n}}{2} \mathcal{F}_n(f)(\xi) I_n\left(\frac{\xi}{n}\right) K_n\left(\frac{\xi}{n}\right),$$

where I_n is defined by (7.17) and K_n is given by

$$K_n(y) = \int_{-1/2}^{1/2} \frac{(e^{2i\pi(y-x)} + e^{2i\pi x})(e^{-2i\pi x} - 1)}{(\Lambda + 4\gamma_n - i\Omega)(y - x, x)} dx. \tag{7.19}$$

We need to show that $\lim_{n \rightarrow \infty} n^2 \|\theta_n\|_{2,n}^2 = 0$. By the Parseval–Plancherel’s relation, this is equivalent to prove that

$$\lim_{n \rightarrow \infty} n^3 \int_{-n/2}^{n/2} |\mathcal{F}_n(f)(\xi)|^2 |I_n\left(\frac{\xi}{n}\right)|^2 |K_n\left(\frac{\xi}{n}\right)|^2 d\xi = 0.$$

By using the change of variables $y = \xi/n$, Lemmas 7.1 and 7.7, we have

$$n^3 \int_{-n/2}^{n/2} |\mathcal{F}_n(f)(\xi)|^2 |I_n(\frac{\xi}{n})|^2 |K_n(\frac{\xi}{n})|^2 d\xi \leq Cn^4 \int_{-1/2}^{1/2} \frac{|y|^4}{1+|ny|^p} dy = \frac{C}{n} \int_{-n/2}^{n/2} \frac{|z|^4}{1+|z|^p} dz$$

which goes to 0, as $n \rightarrow \infty$, for p bigger than 5.

Few Integral Estimates

The following lemma is the new technical estimate which takes into account the extra-term coming from the velocity-flip noise of intensity γ_n .

Lemma 7.6 *Recall that G_n and G_0 are defined by (7.11) and (7.12) respectively. There exists a constant $C > 0$ such that for any $|y| \leq 1/2$ and for all $n \in \mathbb{N}_0$,*

$$|G_n(y) - G_0(y)| \leq C [\sin^2(\pi y) + \gamma_n^2 |\sin(\pi y)|^{-1/2} + \gamma_n |\sin(\pi y)|^{1/2}]. \tag{7.20}$$

Proof We compute the function G_n thanks to the residue theorem. For any $y \in [-1/2, 1/2]$ we denote by $w := w(y)$ the complex number $w = e^{2i\pi y}$. By denoting $z = e^{2i\pi x}$, for $x \in [-1/2, 1/2]$, we have that

$$\begin{aligned} \Lambda(y - x, x) &= 4 - z(w^{-1} + 1) - z^{-1}(w + 1), \\ i \Omega(y - x, x) &= z(1 - w^{-1}) + z^{-1}(w - 1). \end{aligned}$$

We denote by \mathcal{C} the unit circle positively oriented. Then, we have

$$G_n(y) = \frac{1}{16i\pi} \oint_{\mathcal{C}} f_w(z) dz$$

where the meromorphic function f_w is defined by

$$f_w(z) = \frac{[(w - 1) + z^2(1 - w^{-1})]^2}{z^2(z^2 - 2z(1 + \gamma_n) + w)}.$$

The poles of f_w are 0 and z_-, z_+ , the latter being the two solutions of $z^2 - 2z(1 + \gamma_n) + w$. We can check that

$$z_{\pm} = (1 + \gamma_n) \pm \sqrt{\alpha} e^{i\theta/2}, \tag{7.21}$$

where

$$\begin{aligned} \alpha^2 &= 4(1 + \gamma_n)^2 \sin^2(\pi y) + [(1 + \gamma_n)^2 - 1]^2, \\ \theta &= \arctan\left(\frac{(1 + \gamma_n)^2 - \cos(2\pi y)}{\sin(2\pi y)}\right) - \frac{\pi}{2} \operatorname{sgn}(y). \end{aligned}$$

Observe that $|z_-| < 1$ and $|z_+| > 1$. Indeed, since

$$\frac{(1 + \gamma_n)^2 - \cos(2\pi y)}{\sin(2\pi y)} = \frac{\gamma_n^2 + 2\gamma_n + 2 \sin^2(\pi y)}{\sin(2\pi y)}$$

we have $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$ (distinguish the case $y > 0$ and the case $y < 0$). Moreover $\gamma_n > 0$ so that a simple geometric argument permits to conclude that $|z_+| > 1$. Since $|z_- z_+| = 1$, we have $|z_-| < 1$.

By the residue theorem,

$$\oint_{\mathcal{C}} f_w(z) dz = 2\pi i [\operatorname{Res}(f_w, 0) + \operatorname{Res}(f_w, z_-)]$$

where $\text{Res}(f_w, a)$ denotes the value of the residue of f_w at the pole a . An elementary computation shows that

$$\begin{aligned} \text{Res}(f_w, 0) &= \frac{2(w-1)^2}{w^2}(1+\gamma_n), \\ \text{Res}(f_w, z_-) &= \lim_{z \rightarrow z_-} (z-z_-)f_w(z) = \frac{1}{z_- - z_+} \frac{[(w-1) + (1-w^{-1})z_-^2]^2}{z_-^2}. \end{aligned}$$

By using the fact that $z_-^2 = 2z_-(1+\gamma_n) - w$, we obtain that

$$\begin{aligned} G_n(y) &= \frac{1}{8} [\text{Res}(f_w, 0) + \text{Res}(f_w, z_-)] = \frac{(w-1)^2}{4w^2}(1+\gamma_n) \left[1 + \frac{2(1+\gamma_n)}{z_- - z_+} \right] \\ &= \frac{(w-1)^2}{4w^2}(1+\gamma_n) \left[1 - \frac{(1+\gamma_n)}{\sqrt{\alpha}} e^{-i\theta/2} \right] \\ &= \frac{(w-1)^2}{4w^2}(1+\gamma_n) - \frac{(w-1)^2}{4w^2} \frac{(1+\gamma_n)^{3/2}}{\sqrt{2}|\sin(\pi y)|} \left[1 + \frac{(2+\gamma_n)^2}{4(1+\gamma_n)^2} \frac{\gamma_n^2}{\sin^2(\pi y)} \right]^{-1/4} e^{-i\theta/2} \\ &= \left[\frac{(w-1)^2}{4w^2} - \frac{(w-1)^2}{4w^2} \frac{(1+\gamma_n)^{1/2}}{\sqrt{2}|\sin(\pi y)|} e^{-i\theta/2} \right] (1+\gamma_n) + \varepsilon_n(y), \end{aligned}$$

where

$$\varepsilon_n(y) = \frac{(w-1)^2}{4w^2} \frac{(1+\gamma_n)^{3/2}}{\sqrt{2}|\sin(\pi y)|} e^{-i\theta/2} \left\{ 1 - \left[1 + \frac{(2+\gamma_n)^2}{4(1+\gamma_n)^2} \frac{\gamma_n^2}{\sin^2(\pi y)} \right]^{-1/4} \right\}.$$

We deal separately with the two terms. First,

$$\left| \frac{(w-1)^2}{4w^2} \right| = \sin^2(\pi y).$$

Second, after noticing that

$$\frac{1}{4} \leq \frac{(2+\gamma_n)^2}{4(1+\gamma_n)^2} \leq 1,$$

we get

$$\left| 1 - \left[1 + \frac{(2+\gamma_n)^2}{4(1+\gamma_n)^2} \frac{\gamma_n^2}{\sin^2(\pi y)} \right]^{-1/4} \right| \leq C \left| \left[1 + C' \frac{\gamma_n^2}{\sin^2(\pi y)} \right]^{1/4} - 1 \right| \leq C'' \frac{\gamma_n^2}{\sin^2(\pi y)}.$$

Then, we get easily that

$$|\varepsilon_n(y)| \leq C \gamma_n^2 |\sin(\pi y)|^{-1/2}.$$

Moreover, we have that

$$|(1+\gamma_n)^{1/2} - 1| \leq C \gamma_n.$$

Since the derivative of arctan is bounded above by a constant, we have also that

$$\begin{aligned} & \left| e^{-i\theta/2} - \frac{1}{\sqrt{2}}(1 + i \operatorname{sgn}(y))e^{-i\pi y/2} \right| \\ &= \left| \frac{1 + i \operatorname{sgn}(y)}{\sqrt{2}} \right| \left| \exp \left\{ -\frac{i}{2} \left[\arctan \left(\tan(\pi y) + \frac{\gamma_n(2 + \gamma_n)}{\sin(2\pi y)} \right) \right] \right\} - \exp \left\{ -\frac{i\pi y}{2} \right\} \right| \\ &\leq C \left| \arctan \left(\tan(\pi y) + \frac{\gamma_n(2 + \gamma_n)}{\sin(2\pi y)} \right) - \arctan(\tan(\pi y)) \right| \leq C \frac{\gamma_n}{|\sin(\pi y)|}. \end{aligned}$$

We deduce the result

$$|G_n(y) - G_0(y)| \leq C \left[\sin^2(\pi y) + \gamma_n^2 |\sin(\pi y)|^{-1/2} + \gamma_n |\sin(\pi y)|^{1/2} \right].$$

□

The last lemma below is widely inspired from [5].

Lemma 7.7 *The functions I_n , J_n , and K_n , respectively defined by (7.17), (7.18) and (7.19), satisfy for any $y \in \mathbb{R}$ and $n \in \mathbb{N}_0$,*

$$\begin{aligned} |I_n(y)| &\leq C |\sin(\pi y)|^{3/2} \\ |J_n(y)| &\leq C |\sin(\pi y)|^{-1/2} \\ |K_n(y)| &\leq C |\sin(\pi y)|^{1/2} \end{aligned}$$

where C is a positive constant which does not depend on y nor on n .

Proof We divide the proof into three parts, corresponding to the three inequalities

- (1) As previously, we compute I_n by using the residue theorem. For any $y \in [-\frac{1}{2}, \frac{1}{2}]$ we denote by $w := w(y)$ the complex number $w = e^{2i\pi y}$. Then we have

$$I_n(y) = -\frac{1}{4i\pi} \frac{w-1}{w} \oint_{\mathcal{C}} f_w(z) dz, \tag{7.22}$$

where the meromorphic function f_w is defined by

$$f_w(z) = \frac{(z-1)(z^2+w)}{z^2(z-z_+)(z-z_-)}$$

with z_{\pm} defined by (7.21). We recall that $|z_-| < 1$ and $|z_+| > 1$ so that by the residue theorem we have

$$I_n(y) = -\frac{w-1}{2w} [\operatorname{Res}(f_w, 0) + \operatorname{Res}(f_w, z_-)].$$

A simple computation shows that

$$\operatorname{Res}(f_w, 0) = 1 - 2/w, \quad \operatorname{Res}(f_w, z_-) = 1/z_-.$$

It follows that

$$I_n(y) = -\frac{w-1}{2w} \left[\frac{1}{z_-} + 1 - \frac{2}{w} \right].$$

Replacing w and z_- by their explicit values we get the result.

(2) Then we also compute J_n by using the residue theorem:

$$J_n(y) = -\frac{1}{4i\pi} \oint_C f_w(z) dz$$

where the meromorphic function f_w is defined by

$$f_w(z) = \frac{(z^2 + w)}{z^2(z - z_+)(z - z_-)}$$

with z_{\pm} defined by (7.21). By the residue theorem, we get

$$J_n(y) = -\frac{1}{2} (\text{Res}(f_w, 0) + \text{Res}(f_w, z_-)).$$

A simple computation shows that

$$\text{Res}(f_w, 0) = -\frac{w}{2}, \quad \text{Res}(f_w, z_-) = \frac{2}{z_-(z_- - z_+)}.$$

By using the explicit expressions for w , z_{\pm} , we get the result.

(3) Finally, it is not difficult to see that

$$K_n(y) = -\frac{w}{w-1} I_n(y),$$

and by the first estimate the result follows. □

References

1. Basile, G., Olla, S.: Energy diffusion in harmonic system with conservative noise. *J. Stat. Phys.* **155**(6), 1126–1142 (2014)
2. Basile, G., Bernardin, C., Olla, S.: Thermal conductivity for a momentum conserving model. *Commun. Math. Phys.* **287**(1), 67–98 (2009)
3. Bernardin, C.: Superdiffusion of energy in Hamiltonian systems perturbed by a conservative noise in *From Particle Systems to Partial Differential Equations*. In: Springer Proceedings in Mathematics & Statistics, vol. 75 (2014)
4. Bernardin, C., Olla, S.: Fourier’s law for a microscopic model of heat conduction. *J. Stat. Phys.* **121**, 271–289 (2005)
5. Bernardin, C., Gonçalves, P., Jara, M.: Fractional superdiffusion in a system of harmonic oscillators perturbed by a conservative noise. *Arch. Rational Mech. Anal.* (accepted)
6. Bernardin, C., Gonçalves, P.: Anomalous fluctuations for a perturbed hamiltonian system with exponential interactions. *Commun. Math. Phys.* **325**, 291–332 (2014)
7. Bernardin, C., Stoltz, G.: Anomalous diffusion for a class of systems with two conserved quantities. *Nonlinearity* **25**(4), 1099–1133 (2012)
8. Bonetto, F., Lebowitz, J.L., Lukkarinen, J.: Fourier’s law for a harmonic crystal with self-consistent stochastic reservoirs. *J. Stat. Phys.* **116**, 783–813 (2004)
9. Brox, T., Rost, H.: Equilibrium fluctuations of stochastic particle systems: the role of conserved quantities. *Ann. Probab.* **12**, 742–759 (1984)
10. Dhar, A.: Heat transport in low-dimensional systems. *Adv. Phys.* **57**, 457 (2008)
11. Dawson, D.A., Gorostiza, L.G.: Generalized solutions of a class of nuclear-space-valued stochastic evolution equations. *Appl. Math. Optim.* **22**, 241–263 (1990)
12. Fritz, J., Funaki, T., Lebowitz, J.L.: Stationary states of random hamiltonian systems. *Probab. Theory Relat. Fields* **99**, 211–236 (1994)
13. Jara, M.: Current and density fluctuations for interacting particle systems with anomalous diffusive behavior. [arXiv:0901.0229](https://arxiv.org/abs/0901.0229) (2014)
14. Kipnis, C., Landim, C.: *Scaling limits of interacting particle systems*. Springer, Berlin (1999)

15. Komorowski, T., Landim, C. Olla, S.: Fluctuations in Markov processes, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Time symmetry and martingale approximation, vol. 345. Springer, Heidelberg (2012)
16. Lanford, O.E., Lebowitz, J.L., Lieb, E.H.: Time evolution of infinite anharmonic systems. *J. Stat. Phys.* **16**(6), 453–461 (1977)
17. Lepri, S., Livi, R., Politi, A.: Thermal conduction in classical low-dimensional lattices. *Phys. Rep.* **377**, 1–80 (2003)
18. Simon, M.: Hydrodynamic limit for the velocity-flip model. *Stoch. Process. Appl.* **123**, 3623–3662 (2013)
19. Spohn, H.: Nonlinear fluctuating hydrodynamics for anharmonic chains. *J. Stat. Phys.* **154**, 11911–71227 (2014)