

GLUING RESIDUATED LATTICES

Sara Ugolini

University of Denver, Department of Mathematics

sara.ugolini@du.edu

Ongoing joint work with Nick Galatos

TACL 2019

A **residuated lattice**, or RL, is a structure $\mathbf{A} = (A, \cdot, \backslash, /, \wedge, \vee, 1)$ where:

- (A, \wedge, \vee) is a lattice,
- $(A, \cdot, 1)$ is a monoid,
- $x \cdot y \leq z$ iff $y \leq x \backslash z$ iff $x \leq z / y$.

RLs constitute a variety, \mathbb{RL} .

A **residuated lattice**, or RL, is a structure $\mathbf{A} = (A, \cdot, \backslash, /, \wedge, \vee, 1)$ where:

- (A, \wedge, \vee) is a lattice,
- $(A, \cdot, 1)$ is a monoid,
- $x \cdot y \leq z$ iff $y \leq x \backslash z$ iff $x \leq z / y$.

RLs constitute a variety, \mathbb{RL} .

We call a residuated lattice **integral** if 1 is the top element of the lattice, and **bounded** if there is an extra constant 0 in the signature that is the least element of the lattice.

The RL is **commutative** if the monoid is commutative, and we write $x \rightarrow y$ for $x \backslash y = y / x$.

A **residuated lattice**, or RL, is a structure $\mathbf{A} = (A, \cdot, \backslash, /, \wedge, \vee, 1)$ where:

- (A, \wedge, \vee) is a lattice,
- $(A, \cdot, 1)$ is a monoid,
- $x \cdot y \leq z$ iff $y \leq x \backslash z$ iff $x \leq z / y$.

RLs constitute a variety, \mathbb{RL} .

We call a residuated lattice **integral** if 1 is the top element of the lattice, and **bounded** if there is an extra constant 0 in the signature that is the least element of the lattice.

The RL is **commutative** if the monoid is commutative, and we write $x \rightarrow y$ for $x \backslash y = y / x$.

Residuated lattices are the equivalent algebraic semantics of substructural logics, that include: classical logic, intuitionistic logic, linear logic, relevance logic, fuzzy logics...

Examples: Boolean algebras, Heyting algebras, MV-algebras, lattice ordered groups...

- Residuated lattices do not satisfy any special purely lattice-theoretic or monoid-theoretic property
- R. Belohlavek, V. Vychodil, *Residuated Lattices of Size ≤ 12* , *Order*, 27:147–161, 2010 :

	1	2	3	4	5	6	7	8	9	10	11	12
Lattices	1	1	1	2	5	15	53	222	1,078	5,994	37,622	262,776
Residuated lattices	1	1	2	7	26	129	723	4,712	34,698	290,565	2,779,183	30,653,419
Linear res. lattices	1	1	2	6	22	94	451	2,386	13,775	86,417	590,489	4,446,029


- Large classes of residuated lattices currently lack a structural understanding.

ORDINAL SUM

An example of a construction that puts together two integral residuated (\wedge -semi)lattices to give a new one is the **ordinal sum**, first introduced by Ferreirim.


ORDINAL SUM

An example of a construction that puts together two integral residuated (\wedge -semi)lattices to give a new one is the **ordinal sum**, first introduced by Ferreirim.




ORDINAL SUM

An example of a construction that puts together two integral residuated (\wedge -semi)lattices to give a new one is the **ordinal sum**, first introduced by Ferreirim.



ORDINAL SUM

An example of a construction that puts together two integral residuated (\wedge -semi)lattices to give a new one is the **ordinal sum**, first introduced by Ferreirim.



ORDINAL SUM

The ordinal sum has proved to be a powerful construction, in particular in the realm of the algebraic semantics of fuzzy logics.

Aglianò and Montagna in 2003 prove the following characterization:
Every totally ordered basic hoop (or BL algebra) is the ordinal sum of a family of Wajsberg hoops (whose first component is bounded).

ORDINAL SUM

The ordinal sum has proved to be a powerful construction, in particular in the realm of the algebraic semantics of fuzzy logics.

Aglianò and Montagna in 2003 prove the following characterization:
Every totally ordered basic hoop (or BL algebra) is the ordinal sum of a family of Wajsberg hoops (whose first component is bounded).

BL-algebras are the algebraic semantics of Hájek's Basic Logic, the logic of continuous t-norms, and are semilinear (subdirect products of chains) commutative integral residuated lattices satisfying divisibility: $x \wedge y = x \cdot (x \rightarrow y)$, while basic hoops are their 0-free reducts.

ORDINAL SUM

The ordinal sum has proved to be a powerful construction, in particular in the realm of the algebraic semantics of fuzzy logics.


Aglianò and Montagna in 2003 prove the following characterization:
Every totally ordered basic hoop (or BL algebra) is the ordinal sum of a family of Wajsberg hoops (whose first component is bounded).

BL-algebras are the algebraic semantics of Hájek's Basic Logic, the logic of continuous t-norms, and are semilinear (subdirect products of chains) commutative integral residuated lattices satisfying divisibility: $x \wedge y = x \cdot (x \rightarrow y)$, while basic hoops are their 0-free reducts.

The ordinal sum construction intuitively stacks one (semi)lattice on top of the other gluing together the top elements.


What if we glue more? For example, a filter.

Let $\mathbf{B} = (B, \cdot, \backslash, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \backslash, /, \wedge, 1)$ be integral residuated \wedge -semilattices, that intersect in a principal lattice filter generated by an element a that is **conical** (comparable with all other elements) and **idempotent** ($a \cdot a = a$).




If \mathbf{B} and \mathbf{C} are residuated **lattices**, to obtain a lattice structure we need \mathbf{C} to have a lower bound or a to be join-irreducible in \mathbf{B} .

Let $\mathbf{B} = (B, \cdot, \backslash, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \backslash, /, \wedge, 1)$ be integral residuated \wedge -semilattices, that intersect in a principal lattice filter generated by an element a that is **conical** (comparable with all other elements) and **idempotent** ($a \cdot a = a$).



If \mathbf{B} and \mathbf{C} are residuated **lattices**, to obtain a lattice structure we need \mathbf{C} to have a lower bound or a to be join-irreducible in \mathbf{B} .

Let $\mathbf{B} = (B, \cdot, \backslash, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \backslash, /, \wedge, 1)$ be integral residuated \wedge -semilattices, that intersect in a principal lattice filter generated by an element a that is **conical** (comparable with all other elements) and **idempotent** ($a \cdot a = a$).



If \mathbf{B} and \mathbf{C} are residuated **lattices**, to obtain a lattice structure we need \mathbf{C} to have a lower bound or a to be join-irreducible in \mathbf{B} .

PROPERTIES

The **gluing** of **B** and **C** with respect to a , $\mathbf{B} \oplus^a \mathbf{C}$ is an integral residuated (\wedge -semi)lattice.

PROPERTIES

The **gluing** of **B** and **C** with respect to a , $\mathbf{B} \oplus^a \mathbf{C}$ is an integral residuated (\wedge -semi)lattice.

The ordinal sum is a gluing where $a = 1$, $\oplus = \oplus_1$.

PROPERTIES

The **gluing** of **B** and **C** with respect to a , $\mathbf{B} \oplus^a \mathbf{C}$ is an integral residuated (\wedge -semi)lattice.

The ordinal sum is a gluing where $a = 1$, $\oplus = \oplus_1$.

The gluing construction preserves:

- all unary equations without join (e.g. it preserves n -potency, $x^n = x^{n+1}$, for every $n \geq 1$)

PROPERTIES

The **gluing** of **B** and **C** with respect to a , $\mathbf{B} \oplus^a \mathbf{C}$ is an integral residuated (\wedge -semi)lattice.

The ordinal sum is a gluing where $a = 1$, $\oplus = \oplus_1$.

The gluing construction preserves:

- all unary equations without join (e.g. it preserves n -potency, $x^n = x^{n+1}$, for every $n \geq 1$)
- If a is join irreducible in **B**, any gluing of the kind $\mathbf{B} \oplus_a \mathbf{C}$ preserves all unary equations satisfied by both **B** and **C**.

PROPERTIES

The **gluing** of **B** and **C** with respect to a , $\mathbf{B} \oplus^a \mathbf{C}$ is an integral residuated (\wedge -semi)lattice.

The ordinal sum is a gluing where $a = 1$, $\oplus = \oplus_1$.

The gluing construction preserves:

- all unary equations without join (e.g. it preserves n -potency, $x^n = x^{n+1}$, for every $n \geq 1$)
- If a is join irreducible in **B**, any gluing of the kind $\mathbf{B} \oplus_a \mathbf{C}$ preserves all unary equations satisfied by both **B** and **C**.
- commutativity

PROPERTIES

The **gluing** of **B** and **C** with respect to a , $\mathbf{B} \oplus^a \mathbf{C}$ is an integral residuated (\wedge -semi)lattice.

The ordinal sum is a gluing where $a = 1$, $\oplus = \oplus_1$.

The gluing construction preserves:

- all unary equations without join (e.g. it preserves n -potency, $x^n = x^{n+1}$, for every $n \geq 1$)
- If a is join irreducible in **B**, any gluing of the kind $\mathbf{B} \oplus_a \mathbf{C}$ preserves all unary equations satisfied by both **B** and **C**.
- commutativity
- divisibility $x \cdot (x \setminus y) = x \wedge y = (y / x) \cdot x$

PROPERTIES

The **gluing** of **B** and **C** with respect to a , $\mathbf{B} \oplus^a \mathbf{C}$ is an integral residuated (\wedge -semi)lattice.

The ordinal sum is a gluing where $a = 1$, $\oplus = \oplus_1$.


The gluing construction preserves:

- all unary equations without join (e.g. it preserves n -potency, $x^n = x^{n+1}$, for every $n \geq 1$)
- If a is join irreducible in **B**, any gluing of the kind $\mathbf{B} \oplus_a \mathbf{C}$ preserves all unary equations satisfied by both **B** and **C**.
- commutativity
- divisibility $x \cdot (x \setminus y) = x \wedge y = (y / x) \cdot x$
- when not an ordinal sum, prelinearity $(x \setminus y) \vee (y \setminus x) = 1$

If we glue **commutative** integral residuated lattices, the gluing construction is reflected in the deductive filters lattice.

If we glue **commutative** integral residuated lattices, the gluing construction is reflected in the deductive filters lattice.


$\text{Fil}(\mathbf{B})$



$\text{Fil}(\mathbf{C})$




$\text{Fil}(\mathbf{B} \oplus^a \mathbf{C})$



If we glue **commutative** integral residuated lattices, the gluing construction is reflected in the deductive filters lattice.


$\text{Fil}(\mathbf{B})$



$\text{Fil}(\mathbf{C})$



$\text{Fil}(\mathbf{B} \oplus^a \mathbf{C})$



Since a coatom in the lattice of deductive filters would need to be above $\langle a \rangle$, $\mathbf{B} \oplus^a \mathbf{C}$ is subdirectly irreducible iff \mathbf{B} and \mathbf{C} are subdirectly irreducible.

In the non-commutative case, it is slightly more complicated.

Given an integral RL \mathbf{A} , $F \subseteq A$ is a *deductive filter* if it is a lattice filter closed under products and under conjugates: if $x \in F$, then $yx/y, y \setminus xy \in F$ for every $y \in A$.

In the non-commutative case, it is slightly more complicated.

Given an integral RL \mathbf{A} , $F \subseteq A$ is a *deductive filter* if it is a lattice filter closed under products and under conjugates: if $x \in F$, then $yx/y, y \setminus xy \in F$ for every $y \in A$.

- If $\uparrow a$ is a deductive filter (e.g. a is a central element) in \mathbf{B} , we get the same picture of the commutative case.

In the non-commutative case, it is slightly more complicated.

Given an integral RL \mathbf{A} , $F \subseteq A$ is a *deductive filter* if it is a lattice filter closed under products and under conjugates: if $x \in F$, then $yx/y, y \setminus xy \in F$ for every $y \in A$.

- If $\uparrow a$ is a deductive filter (e.g. a is a central element) in \mathbf{B} , we get the same picture of the commutative case.
- Otherwise, $\text{Fil}(\mathbf{B} \oplus^a \mathbf{C})$ is isomorphic to $\text{Fil}(\mathbf{B})$.


In the non-commutative case, it is slightly more complicated.

Given an integral RL \mathbf{A} , $F \subseteq A$ is a *deductive filter* if it is a lattice filter closed under products and under conjugates: if $x \in F$, then $yx/y, y \setminus xy \in F$ for every $y \in A$.

- If $\uparrow a$ is a deductive filter (e.g. a is a central element) in \mathbf{B} , we get the same picture of the commutative case.
- Otherwise, $\text{Fil}(\mathbf{B} \oplus^a \mathbf{C})$ is isomorphic to $\text{Fil}(\mathbf{B})$.
- Thus, in any case, $\mathbf{B} \oplus^a \mathbf{C}$ is subdirectly irreducible iff \mathbf{B} is subdirectly irreducible.


What if we want to glue a filter and an ideal?

Let now $\mathbf{B} = (B, \cdot, \backslash, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \backslash, /, \wedge, 1)$ be integral residuated \wedge -semilattices, $B \cap C = \uparrow a \cup \downarrow d$ with a, d conical and idempotent.




If \mathbf{B} and \mathbf{C} are lattices, we need $\mathbf{C} \setminus \{\downarrow d\}$ with a lower bound or a \vee -irreducible in \mathbf{B} to have a lattice structure.

Let now $\mathbf{B} = (B, \cdot, \backslash, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \backslash, /, \wedge, 1)$ be integral residuated \wedge -semilattices, $B \cap C = \uparrow a \cup \downarrow d$ with a, d conical and idempotent.




If \mathbf{B} and \mathbf{C} are lattices, we need $\mathbf{C} \setminus \{\downarrow d\}$ with a lower bound or a \vee -irreducible in \mathbf{B} to have a lattice structure.

Let now $\mathbf{B} = (B, \cdot, \backslash, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \backslash, /, \wedge, 1)$ be integral residuated \wedge -semilattices, $B \cap C = \uparrow a \cup \downarrow d$ with a, d conical and idempotent.




If \mathbf{B} and \mathbf{C} are lattices, we need $\mathbf{C} \setminus \{\downarrow d\}$ with a lower bound or a \vee -irreducible in \mathbf{B} to have a lattice structure.

Let now $\mathbf{B} = (B, \cdot, \setminus, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \setminus, /, \wedge, 1)$ be integral residuated \wedge -semilattices, $B \cap C = \uparrow a \cup \downarrow d$ with a, d conical and idempotent.




If \mathbf{B} and \mathbf{C} are lattices, we need $\mathbf{C} \setminus \{\downarrow d\}$ with a lower bound or a \vee -irreducible in \mathbf{B} to have a lattice structure.

Let now $\mathbf{B} = (B, \cdot, \setminus, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \setminus, /, \wedge, 1)$ be integral residuated \wedge -semilattices, $B \cap C = \uparrow a \cup \downarrow d$ with a, d conical and idempotent.




If \mathbf{B} and \mathbf{C} are lattices, we need $\mathbf{C} \setminus \{\downarrow d\}$ with a lower bound or a \vee -irreducible in \mathbf{B} to have a lattice structure.

Let now $\mathbf{B} = (B, \cdot, \backslash, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \backslash, /, \wedge, 1)$ be integral residuated \wedge -semilattices, $B \cap C = \uparrow a \cup \downarrow d$ with a, d conical and idempotent.



If \mathbf{B} and \mathbf{C} are lattices, we need $\mathbf{C} \setminus \{\downarrow d\}$ with a lower bound or a \vee -irreducible in \mathbf{B} to have a lattice structure.

Let now $\mathbf{B} = (B, \cdot, \backslash, /, \wedge, 1)$, $\mathbf{C} = (C, \cdot, \backslash, /, \wedge, 1)$ be integral residuated \wedge -semilattices, $B \cap C = \uparrow a \cup \downarrow d$ with a, d conical and idempotent.



If \mathbf{B} and \mathbf{C} are lattices, we need $\mathbf{C} \setminus \{\downarrow d\}$ with a lower bound or a \vee -irreducible in \mathbf{B} to have a lattice structure.

The **gluing** of \mathbf{B} and \mathbf{C} with respect to a and d , $\mathbf{B} \oplus_d^a \mathbf{C}$ is an integral residuated (\wedge -semi)lattice.

PROPERTIES

This new gluing construction still preserves commutativity and prelinearity (if $a \neq 1$), but it does not preserve divisibility.

In the commutative case, since a is again an idempotent conical element, gluings of subdirectly irreducible are subdirectly irreducible.

EXAMPLES

Let us consider bounded totally ordered CIRLs that are 2-potent, i.e. that satisfy $x^2 = x^3$, and such that for every x, y

$$x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

EXAMPLES

Let us consider bounded totally ordered CIRLs that are 2-potent, i.e. that satisfy $x^2 = x^3$, and such that for every x, y

$$x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

This is a positive universal first-order formula, these structures generate a variety of MTL-algebras (bounded semilinear CIRLs) that satisfy $x^2 = x^3$ and

$$x \vee ((x \cdot (x \wedge y)) \setminus (x \wedge y)^2) = 1.$$

This variety has the FMP (finite model property, i.e. it is generated by its finite members).

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

.1

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

- 1
- x_1

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

- 1
- x_1

$$\bullet x_1^2 = x_1^3$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

$$\begin{array}{l} \bullet 1 \\ \vdots \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 = x_1^3 \end{array}$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

$$\begin{array}{l} \bullet 1 \\ \vdots \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 = x_1^3 \\ \bullet y_1 \end{array}$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

$$\begin{array}{l} \bullet 1 \\ \vdots \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 = x_1^3 \\ \bullet y_1 \\ \\ \bullet y_1^2 = y_1^3 \end{array}$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

$$\begin{array}{l} \bullet 1 \\ \vdots \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 = x_1^3 \\ \vdots \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = y_1^3 \end{array}$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$


The finite subdirectly irreducible will be chains:

$$\begin{array}{c} \bullet 1 \\ \vdots \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 = x_1^3 \\ \vdots \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = y_1^3 \\ \vdots \\ \bullet \\ \vdots \\ \bullet \\ \vdots \\ \bullet 0 \end{array}$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:




$$x_i \cdot x_j = x_i^2 = x_j^2 = x_1^2$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:



$$x_i \cdot x_j = x_i^2 = x_j^2 = x_1^2$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

$$\begin{array}{l}
 \bullet 1 \\
 \bullet x_1 \\
 \vdots \\
 x_i \vdots \\
 x_j \vdots \\
 \bullet x_1^2 = x_1^3 \\
 \bullet y_1 \\
 y_k \vdots \\
 \bullet y_1^2 = y_1^3 \\
 \vdots \\
 \bullet 0
 \end{array}
 \quad
 \begin{array}{l}
 \\
 \\
 \\
 x_i \cdot x_j = x_i^2 = x_j^2 = x_1^2 \\
 \\
 x_i \cdot y_k \leq y_k^2 = y_1^2 \Rightarrow x_i \cdot y_k = y_1^2 = x_1^2 \cdot y_k \\
 \\
 \\
 \\
 \\
 \end{array}$$

EXAMPLE

$$x^2 = x^3, \quad x = 1 \text{ or } x \cdot (x \wedge y) \leq (x \wedge y)^2$$

The finite subdirectly irreducible will be chains:

$$\begin{array}{l}
 \bullet 1 \\
 \bullet x_1 \\
 x_i \vdots \\
 x_j \vdots \\
 \bullet x_1^2 = x_1^3 \\
 \bullet y_1 \\
 y_k \vdots \\
 \bullet y_1^2 = y_1^3 \\
 \vdots \\
 \vdots \\
 \bullet 0
 \end{array}
 \quad
 \begin{array}{l}
 \\
 \\
 x_i \cdot x_j = x_i^2 = x_j^2 = x_1^2 \\
 \\
 x_i \cdot y_k \leq y_k^2 = y_1^2 \Rightarrow x_i \cdot y_k = y_1^2 = x_1^2 \cdot y_k \\
 \\
 \\
 \\
 \\
 \\
 \end{array}$$

We can generate these chains as gluings of finite simple 2-potent MTL-chains.

• 1

• 0

$$\begin{array}{ccc} \bullet 1 & & \bullet 1 \\ \bullet 0 & \oplus 1 & \bullet x_1 \\ & & \vdots \\ & & \bullet 0 = x_1^2 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet 0 \end{array} \oplus^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet 0 = x_1^2 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet 0 \end{array} \oplus^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet 0 = x_1^2 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} \quad \begin{array}{c} \bullet 1 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet 0 \end{array} \oplus^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet 0 = x_1^2 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} \qquad \begin{array}{c} \bullet 1 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array} \oplus_0^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet 0 \end{array} \oplus^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet 0 = x_1^2 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} \qquad \begin{array}{c} \bullet 1 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array} \oplus_0^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet 0 \end{array} \oplus^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet 0 = x_1^2 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} \qquad \begin{array}{c} \bullet 1 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array} \oplus_0^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet 0 \end{array} \oplus^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet 0 = x_1^2 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} \qquad \begin{array}{c} \bullet 1 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array} \oplus_0^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array}$$


$$\begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array} \oplus^{x_1^2} \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet z_1 \\ \vdots \\ \bullet z_1^2 = 0 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet 0 \end{array} \oplus^1 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet 0 = x_1^2 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} \qquad \begin{array}{c} \bullet 1 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array} \oplus^1_0 \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet 0 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array}$$

$$\begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array} \oplus^{x_1^2} \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet z_1 \\ \vdots \\ \bullet z_1^2 = 0 \end{array} = \begin{array}{c} \bullet 1 \\ \bullet x_1 \\ \vdots \\ \bullet x_1^2 \\ \bullet z_1 \\ \vdots \\ \bullet z_1^2 \\ \bullet y_1 \\ \vdots \\ \bullet y_1^2 = 0 \end{array}$$

EXAMPLES


Given any bounded commutative residuated lattice \mathbf{A} , we can glue its disconnected rotation and its n -lifting: $\mathbf{L}_n \oplus \mathbf{A}$



What we obtain has been introduced as generalized n -rotation of \mathbf{A} in [Busaniche, Marcos, U. - 2019].

EXAMPLES


Given any bounded commutative residuated lattice \mathbf{A} , we can glue its disconnected rotation and its n -lifting: $\mathbf{L}_n \oplus \mathbf{A}$



What we obtain has been introduced as generalized n -rotation of \mathbf{A} in [Busaniche, Marcos, U. - 2019].

EXAMPLES

Given any bounded commutative residuated lattice \mathbf{A} , we can glue its disconnected rotation and its n -lifting: $\mathbf{L}_n \oplus \mathbf{A}$



What we obtain has been introduced as generalized n -rotation of \mathbf{A} in [Busaniche, Marcos, U. - 2019].

Thank you!