

Norm complete Abelian l-groups: equational axiomatization

Marco Abbadini

Dipartimento di Matematica *Federigo Enriques*

University of Milan, Italy

`marco.abbadini@unimi.it`

Joint work with Vincenzo Marra and Luca Spada

TACL 2019

Nice, France

June 20, 2019

Main question

Is the category of norm-complete ℓ -groups equivalent to a variety of (possibly infinitary) algebras?

Main question

Is the category of norm-complete ℓ -groups equivalent to a variety of (possibly infinitary) algebras?

Definition

Variety of algebras := category of \mathcal{L} -algebras (where \mathcal{L} is a set of function symbols) satisfying a certain set of \mathcal{L} -equations.

$$\forall \underline{x} \quad \gamma(\underline{x}) = \eta(\underline{x}).$$

(We admit operations of infinite arity.)

EXAMPLE OF NORM-COMPLETE ℓ -GROUP

Let X be a compact Hausdorff space, and, for every $x \in X$, let us assign a set A_x such that either $A_x := \frac{1}{n}\mathbb{Z}$ for some $n \in \mathbb{N}_{>0}$, or $A_x = \mathbb{R}$. We can encode $(A_x)_{x \in X}$ via a function $\zeta: X \rightarrow \mathbb{N}$.

$$\mathcal{C}_\zeta(X) :=$$

$$\{f: X \rightarrow \mathbb{R} \mid f \text{ continuous, } \forall x \in X \ f(x) \in A_x\} =$$

$$\{f: X \rightarrow \mathbb{R} \mid f \text{ continuous, } \forall x \in X \ \text{den}(f(x)) \text{ divides } \zeta(x)\}.$$

$\mathcal{C}_\zeta(X)$, endowed with pointwise operations $+$, \vee , \wedge , $-$, 0 , 1 , is an *Abelian lattice-ordered group* (*l-group*, for short):

1. $\langle \mathcal{C}_\zeta(X), 0, +, - \rangle$ is an Abelian group;
2. $\langle \mathcal{C}_\zeta(X), \vee, \wedge \rangle$ is a lattice;
3. the order is translation invariant:

$$\forall f, g, h \in \mathcal{C}_\zeta(X) \quad f \leq g \Rightarrow f + h \leq g + h.$$

- ▶ 1 is a *strong unit*:
for all $f \in \mathcal{C}_\zeta(X)$, there exists $n \in \mathbb{N}$ s.t. $(-n)1 \leq f \leq (n)1$,
- ▶ $\mathcal{C}_\zeta(X)$ is *Archimedean*:
for all $f, g \in \mathcal{C}_\zeta(X)$ such that $f \geq 0$ and $g \geq 0$ we have:
if, for all $n \in \mathbb{N}$, $(n)f \leq g$, then $f = 0$.
- ▶ $\mathcal{C}_\zeta(X)$ is *norm-complete*, i.e., complete in the metric induced by the supremum norm

$$\|f\| := \inf \left\{ \frac{p}{q} \in \mathbb{Q} \mid p, q \in \mathbb{N}, q \neq 0, (q)|f| \leq (p)1 \right\}.$$

Norm-complete ℓ -group := ℓ -group with strong unit, which is Archimedean and norm-complete.

$\mathcal{C}_\zeta(X)$ is a norm-complete ℓ -group, and, viceversa, every norm-complete ℓ -group is of this form, for some choice of X and ζ .

Morphisms of norm-complete ℓ -groups: functions that preserve $+, \vee, \wedge, -, 0, 1$.

Main question

Is the category of norm complete ℓ -groups equivalent to a variety of (possibly infinitary) algebras?

Main question

Is the category of norm complete ℓ -groups equivalent to a variety of (possibly infinitary) algebras?

Answer (main result)

Yes.

Main question

Is the category of norm complete ℓ -groups equivalent to a variety of (possibly infinitary) algebras?

Answer (main result)

Yes.

In the following: we provide an explicit finite equational axiomatization of this infinitary variety.

Is the class of norm-complete ℓ -groups closed (in the class of $\{+, \vee, \wedge, -, 0, 1\}$ -algebras) under...

1. ... **products?**

No, 1 is a *strong unit* of \mathbb{R} , but not of $\mathbb{R}^{\mathbb{N}}$. **X**

2. ... **subalgebras?**

No, \mathbb{R} is *norm-complete*, but $\mathbb{Q} \subseteq \mathbb{R}$ is not. **X**

3. ... **homomorphic images?**

No, the image of a norm-complete ℓ -group might fail to be *Archimedean*. **X**

Is the class of norm-complete ℓ -groups closed (in the class of $\{+, \vee, \wedge, -, 0, 1\}$ -algebras) under...

1. ... **products?**

No, 1 is a *strong unit* of \mathbb{R} , but not of $\mathbb{R}^{\mathbb{N}}$. \times

2. ... **subalgebras?**

No, \mathbb{R} is *norm-complete*, but $\mathbb{Q} \subseteq \mathbb{R}$ is not. \times

3. ... **homomorphic images?**

No, the image of a norm-complete ℓ -group might fail to be *Archimedean*. \times

Idea: introduce some additional operations together with new axioms regulating them.

This might solve 2 and 3. But not 1.

To solve the problem given by the *strong unit*, we use the theory of MV-algebras.

Given an ℓ -group G with strong unit (*ul*-group, for short),

$$\Gamma(G) := \{x \in G \mid 0 \leq x \leq 1\}.$$

For $x, y \in \Gamma(G)$,

$$x \oplus y := (x + y) \wedge 1;$$

$$\neg x := 1 - x.$$

An MV-algebra is a structure $(A, \oplus, \neg, 0)$ such that

$$(A, \oplus, 0) \text{ is a commutative monoid.} \quad (\text{MV 1})$$

$$x \oplus \neg 0 = \neg 0. \quad (\text{MV 2})$$

$$\neg(\neg x) = x. \quad (\text{MV 3})$$

$$\neg(\neg x \oplus y) \oplus y = \neg(\neg y \oplus x) \oplus x. \quad (\text{MV 4})$$

Mundici showed that Γ establishes an equivalence between the category of *ul*-groups and the category of MV-algebras.

Idea

In addition to the operations of ul -groups, consider an operation γ ($\simeq \lim$) of countably infinite arity, together with some new axioms, so that

$$\gamma(x_1, x_2, x_3, \dots) = \lim_{n \rightarrow \infty} x_n$$

for 'enough' Cauchy sequences (x_1, x_2, x_3, \dots) .

Definition

A sequence (x_1, x_2, x_3, \dots) in a metric space (X, d) is called *super-Cauchy* if, for every $n \geq 2$,

$$d(x_n, x_{n-1}) \leq \frac{1}{2^n}.$$

Every super-Cauchy sequence is Cauchy.

Lemma

(X, d) is complete if, and only if, every super-Cauchy sequence converges.

Intended interpretation of γ on a norm-complete ℓ -group:

$$\gamma(x_1, x_2, x_3, \dots) = \lim_{n \rightarrow \infty} \rho_n(x_1, \dots, x_n)$$

where ρ_n is a term in the language of $u\ell$ -groups—yet to be defined—such that





1. if (x_1, x_2, x_3, \dots) is a super-Cauchy sequence, then, for all n ,

$$\rho_n(x_1, \dots, x_n) = x_n;$$

2. for any (x_1, x_2, x_3, \dots) , the sequence $(\rho_n(x_1, \dots, x_n))_{n \geq 1}$ is super-Cauchy.

For $n \in \mathbb{N}_{>0}$, set

$$\theta_n(x) := \{ \{ [0 \wedge (-(2^{n+1} - 1)x - 1)] \vee x \} \wedge (-(2^{n+1} - 1)x + 1) \} \vee 0.$$



Let us define ρ_n as follows.

$$\rho_1(x_1) := x_1;$$

$$\rho_2(x_1, x_2) := \rho_1(x_1) + \theta_1(x_2 - x_1);$$

$$\rho_3(x_1, x_2, x_3) := \rho_2(x_1, x_2) + \theta_2(x_3 - x_2);$$

\vdots

$$\rho_n(x_1, \dots, x_n) := \rho_{n-1}(x_1, \dots, x_{n-1}) + \theta_{n-1}(x_n - x_{n-1}).$$

For every n , ρ_n is a term of ul -groups.

1. If $(x_n)_{n \in \mathbb{N}_{>0}}$ is a super-Cauchy sequence, then, for all n , $\rho_n(x_1, \dots, x_n) = x_n$.
2. For any $(x_n)_{n \in \mathbb{N}_{>0}}$ the sequence $(\rho_n(x_1, \dots, x_n))_{n \in \mathbb{N}_{>0}}$ is super-Cauchy.

Let us define ρ_n as follows.

$$\begin{aligned}\rho_1(x_1) &:= x_1; \\ \rho_2(x_1, x_2) &:= \rho_1(x_1) + \theta_1(x_2 - x_1); \\ \rho_3(x_1, x_2, x_3) &:= \rho_2(x_1, x_2) + \theta_2(x_3 - x_2); \\ &\vdots \\ \rho_n(x_1, \dots, x_n) &:= \rho_{n-1}(x_1, \dots, x_{n-1}) + \theta_{n-1}(x_n - x_{n-1}).\end{aligned}$$

For every n , ρ_n is a term of ul -groups.

1. If $(x_n)_{n \in \mathbb{N}_{>0}}$ is a super-Cauchy sequence, then, for all n , $\rho_n(x_1, \dots, x_n) = x_n$.
2. For any $(x_n)_{n \in \mathbb{N}_{>0}}$ the sequence $(\rho_n(x_1, \dots, x_n))_{n \in \mathbb{N}_{>0}}$ is super-Cauchy.

Then, in any norm-complete l -group, we can define

$$\gamma(x_1, x_2, x_3, \dots) := \lim_{n \rightarrow \infty} \rho_n(x_1, \dots, x_n)$$

and γ maps super-Cauchy sequences to their limit.

Operations

Operations of ul -group, together with an operation γ of countably infinite arity.

Axioms

0. Axioms of ℓ -groups.
1. The element 1 is a strong unit.
2. $\gamma(x, x, x, \dots) = x$.
3. $\gamma(\theta_1(x), \theta_2(x), \theta_3(x), \dots) = 0$.
4. For each $n \in \mathbb{N}_{>0}$

$$d(\gamma(x_1, x_2, x_3, \dots), \rho_n(x_1, \dots, x_n)) \leq \frac{1}{2^n},$$

i.e.

$$((2^n)|\gamma(x_1, x_2, x_3, \dots) - \rho_n(x_1, \dots, x_n)|) \vee 1 = 1.$$

Every norm-complete ℓ -group satisfies the axioms, with

$$\gamma(x_1, x_2, x_3, \dots) := \lim_{n \rightarrow \infty} \rho_n(x_1, \dots, x_n).$$

Lemma

The axioms

2. $\gamma(x, x, x, \dots) = x;$

3. $\gamma(\theta_1(x), \theta_2(x), \theta_3(x), \dots) = 0.$


imply the Archimedean property.

Proof.

Let x be infinitesimal. Then

$$x \stackrel{2.}{=} \gamma(x, x, x, \dots) = \gamma(\theta_1(x), \theta_2(x), \theta_3(x), \dots) \stackrel{3.}{=} 0.$$

□



The scheme of axioms

4. for each $n \in \mathbb{N}_{>0}$

$$d(\gamma(x_1, x_2, x_3, \dots), \rho_n(x_1, \dots, x_n)) \leq \frac{1}{2^n}$$

'defines' $\gamma(x_1, x_2, x_3, \dots)$ as the limit of $(\rho_n(x_1, \dots, x_n))_{n \in \mathbb{N}_{>0}}$ and implies norm-completeness.

Let G_γ be the category of $\{+, \vee, \wedge, -, 0, 1, \gamma\}$ -algebras satisfying Axioms 0, 1, 2, 3, 4.

Let G be the category of *ul*-groups.

Let $U: G_\gamma \rightarrow G$ be the forgetful functor (that forgets γ).

Theorem

The functor U is injective, full and faithful, and the objects in the image are precisely the norm-complete ℓ -groups.

Corollary

The category of norm-complete ℓ -groups is isomorphic to G_γ .

CONCLUSION

Theorem (Main result)

Up to an equivalence, the category of norm-complete ℓ -groups is a variety of infinitary algebras. Moreover, we have an explicit finite equational axiomatization of this variety.

Thank you.