

# Holomorphic symplectic geometry: a problem list

Arnaud Beauville

**Abstract** The usual structures of symplectic geometry (symplectic, contact, Poisson) make sense for complex manifolds; they turn out to be quite interesting on projective, or compact Kähler, manifolds. In these notes we review some of the recent results on the subject, with emphasis on the open problems and conjectures.

## Introduction

Though symplectic geometry is usually done on real manifolds, the main definitions (symplectic or contact structures, Poisson bracket) make perfect sense in the holomorphic setting. What is less obvious is that these structures are indeed quite interesting in this set-up, in particular on *global* objects – meaning compact, or projective, manifolds. The study of these objects has been much developed in the last 30 years – an exhaustive survey would require at least a book. The aim of these notes is much more modest: we would like to give a (very partial) overview of the subject by presenting some of the open problems which are currently investigated.

Most of the paper is devoted to holomorphic symplectic (= hyperkähler) manifolds, a subject which has been blossoming in recent years. Two short chapters are devoted to contact and Poisson structures : in the former we

---

Arnaud Beauville  
Laboratoire J.-A. Dieudonné (UMR 6621 du CNRS), Université de Nice, Parc Valrose,  
F-06108 Nice cedex 2, France  
e-mail: arnaud.beauville@unice.fr

discuss the conjectural classification of projective contact manifolds, and in the latter an intriguing conjecture of Bondal on the rank of the Poisson tensor.

## 1 Compact hyperkähler manifolds

### 1.1 Basic definitions

The interest for holomorphic symplectic manifolds comes from the following result, stated by Bogomolov in [8] :

**Theorem 1 (Decomposition theorem).** *Let  $X$  be a compact, simply-connected Kähler manifold with trivial canonical bundle. Then  $X$  is a product of manifolds of the following two types:*

- *projective manifolds  $Y$  of dimension  $\geq 3$ , with  $H^0(Y, \Omega_Y^*) = \mathbb{C} \oplus \mathbb{C}\omega$ , where  $\omega$  is a generator of  $K_Y$ ;*
- *compact Kähler manifolds  $Z$  with  $H^0(Z, \Omega_Z^*) = \mathbb{C}[\sigma]$ , where  $\sigma \in H^0(Z, \Omega_Z^2)$  is everywhere non-degenerate.*

This theorem has an important interpretation (and a proof) in terms of Riemannian geometry<sup>1</sup>. By the fundamental theorem of Yau [42], an  $n$ -dimensional compact Kähler manifold  $X$  with trivial canonical bundle admits a Kähler metric with holonomy group contained in  $SU(n)$  (this is equivalent to the vanishing of the Ricci curvature). By the Berger and de Rham theorems,  $X$  is a product of manifolds with holonomy  $SU(m)$  or  $Sp(r)$ ; this corresponds to the first and second case of the decomposition theorem.

We will call the manifolds of the first type *Calabi-Yau* manifolds, and those of the second type *hyperkähler* manifolds (they are also known as *irreducible holomorphic symplectic*).

### 1.2 Examples

For Calabi-Yau manifolds we know a huge quantity of examples (in dimension 3, the number of known families approaches 10 000), but relatively little general theory. In contrast, we have much information on hyperkähler manifolds, their period map, their cohomology (see below); what is lacking severely is

---

<sup>1</sup> See [5] for a more detailed exposition.

examples. In fact, at this time we know two families in each dimension [2], and two isolated families in dimension 6 and 10 [32], [33] :

a) Let  $S$  be a K3 surface. The symmetric product  $S^{(r)} := S^r/\mathfrak{S}_r$  parametrizes subsets of  $r$  points in  $S$ , counted with multiplicities; it is smooth on the open subset  $S_o^{(r)}$  consisting of subsets with  $r$  distinct points, but singular otherwise. If we replace “subset” by (analytic) “subspace”, we obtain a smooth compact manifold, the *Hilbert scheme*  $S^{[r]}$ ; the natural map  $S^{[r]} \rightarrow S^{(r)}$  is an isomorphism above  $S_o^{(r)}$ , but it resolves the singularities of  $S^{(r)}$ .

Let  $\omega$  be a non-zero holomorphic 2-form on  $S$ . The form  $\text{pr}_1^*\omega + \dots + \text{pr}_r^*\omega$  descends to a non-degenerate 2-form on  $S_o^{(r)}$ ; it is easy to check that this 2-form extends to a symplectic structure on  $S^{[r]}$ .

b) Let  $T$  be a 2-dimensional complex torus. The Hilbert scheme  $T^{[r]}$  has the same properties as  $S^{[r]}$ , but it is not simply connected. This is fixed by considering the composite map  $\sigma : T^{[r+1]} \rightarrow T^{(r+1)} \xrightarrow{s} T$ , where  $s(t_1, \dots, t_r) = t_1 + \dots + t_r$ ; the fibre  $K_r(T) := \sigma^{-1}(0)$  is a hyperkähler manifold of dimension  $2r$  (“generalized Kummer manifold”).

c) Let again  $S$  be a K3 surface, and  $\mathcal{M}$  the moduli space of stable rank 2 vector bundles on  $S$ , with Chern classes  $c_1 = 0$ ,  $c_2 = 4$ . According to Mukai [30], this space has a holomorphic symplectic structure. It admits a natural compactification  $\overline{\mathcal{M}}$ , obtained by adding classes of semi-stable torsion free sheaves; it is singular along the boundary, but O’Grady constructs a desingularization of  $\overline{\mathcal{M}}$  which is a new hyperkähler manifold, of dimension 10.

d) The analogous construction can be done starting from rank 2 bundles with  $c_1 = 0$ ,  $c_2 = 2$  on a 2-dimensional complex torus, and taking again some fibre to ensure the simple connectedness. The upshot is a new hyperkähler manifold of dimension 6.

In the two last examples it would seem simpler to start with a moduli space  $\mathcal{M}$  for which the natural compactification  $\overline{\mathcal{M}}$  is smooth; in that case  $\overline{\mathcal{M}}$  is a hyperkähler manifold [30], but it turns out that it is a deformation of  $S^{[r]}$  or  $K_r(T)$  (Göttsche-Huybrechts, O’Grady, Yoshioka ...). On the other hand, when  $\overline{\mathcal{M}}$  is singular, it admits a hyperkähler desingularization only in the two cases considered by O’Grady [23].

Thus it seems that a new idea is required to answer our first problem:

**Question 1.** *Find new examples of hyperkähler manifolds.*

### 1.3 The period map

In dimension 2 the only hyperkähler manifolds are K3 surfaces; we know them very well thanks to the *period map*, which associates to a K3 surface  $S$  the Hodge decomposition

$$H^2(S, \mathbb{C}) = H^{2,0} \oplus H^{1,1} \oplus H^{0,2} .$$

This is determined by the position of the line  $H^{2,0}$  in  $H^2(S, \mathbb{C})$  : indeed we have  $H^{0,2} = \overline{H^{2,0}}$ , and  $H^{1,1}$  is the orthogonal of  $H^{2,0} \oplus H^{0,2}$  with respect to the intersection form. Note that any non-zero element  $\sigma$  of  $H^{2,0}$  (that is, the class of a non-zero holomorphic 2-form) satisfies  $\sigma^2 = 0$  and  $\sigma \cdot \bar{\sigma} > 0$ .

To compare the Hodge structures of different K3 surfaces, we consider *marked* surfaces  $(S, \lambda)$ , where  $\lambda$  is an isometry of  $H^2(S, \mathbb{Z})$  onto a fixed lattice  $L$ , the unique even unimodular lattice  $L$  of signature  $(3, 19)$ . Then the data of the Hodge structure on  $H^2(S, \mathbb{Z})$  is equivalent to that of the *period point*  $\wp(S, \lambda) := \lambda_{\mathbb{C}}(H^{2,0}) \in \mathbb{P}(L_{\mathbb{C}})$ . By the above remark this point lies in the domain  $\Omega \subset \mathbb{P}(L_{\mathbb{C}})$  defined by the conditions  $x^2 = 0$ ,  $x \cdot \bar{x} > 0$ . There is a moduli space  $\mathcal{M}_L$  for marked K3 surfaces, which is a *non-Hausdorff* complex manifold; the *period map*  $\wp : \mathcal{M}_L \rightarrow \Omega_L$  is holomorphic. We know a lot about that map, thanks to the work of many people (Piatetski-Shapiro, Shafarevich, Todorov, Siu, ...):

**Theorem 2.** 1) (“local Torelli”)  $\wp$  is a local isomorphism.

2) (“global Torelli”) If  $\wp(S, \lambda) = \wp(S', \lambda')$ ,  $S$  and  $S'$  are isomorphic;

3) (“surjectivity”) Every point of  $\Omega$  is the period of some marked K3 surface.

Another way of stating 2) is that  $S$  and  $S'$  are isomorphic if and only if there is a *Hodge isometry*  $H^2(S, \mathbb{Z}) \xrightarrow{\sim} H^2(S', \mathbb{Z})$  (that is, an isometry inducing an isomorphism of Hodge structures). There is in fact a more precise statement, see e.g. [1].

There is a very analogous picture for higher-dimensional hyperkähler manifolds. The intersection form is replaced by a canonical quadratic form  $q : H^2(X, \mathbb{Z}) \rightarrow \mathbb{Z}$ , primitive<sup>2</sup>, of signature  $(3, b_2 - 3)$  [2]. The easiest way to define it is through the Fujiki relation

$$\int_X \alpha^{2r} = f_X q(\alpha)^r \quad \text{for each } \alpha \in H^2(X, \mathbb{Z}) , \quad \text{where } \dim(X) = 2r ;$$

---

<sup>2</sup> This means that the associated bilinear form is integral and not divisible by an integer  $> 1$ .

this relation determines  $f_X$  (the *Fujiki constant* of  $X$ ) and the form  $q$ ; they depend only on the topological type of  $X$ .

Let  $X$  be a hyperkähler manifold, and  $L$  a lattice. A *marking* of type  $L$  of  $X$  is an isometry  $\lambda : (H^2(X, \mathbb{Z}), q) \xrightarrow{\sim} L$ . The period of  $(X, \lambda)$  is the point  $\lambda_{\mathbb{C}}(H^{2,0}) \in \mathbb{P}(L_{\mathbb{C}})$ ; as above it belongs to the period domain

$$\Omega_L := \{[x] \in \mathbb{P}(L_{\mathbb{C}}) \mid x^2 = 0, x \cdot \bar{x} > 0\} .$$

Again we have a non-Hausdorff complex manifold  $\mathcal{M}_L$  parametrizing hyperkähler manifolds of a given dimension with a marking of type  $L$ ; the period map  $\varphi : \mathcal{M}_L \rightarrow \Omega_L$  is holomorphic. We have:

**Theorem 3.** 1) *The period map  $\varphi : \mathcal{M}_L \rightarrow \Omega_L$  is a local isomorphism.*

2) *The restriction of  $\varphi$  to any connected component of  $\mathcal{M}_L$  is surjective.*

1) is proved in [2], and 2) in [17]. What is missing is the analogue of the global Torelli theorem. It has long been known that it cannot hold in the form given in Theorem 2; in fact, it follows from the results of [17] that any birational map  $X \dashrightarrow X'$  induces a Hodge isometry  $H^2(X, \mathbb{Z}) \xrightarrow{\sim} H^2(X', \mathbb{Z})$ . This is not the only obstruction: Namikawa observed [31] that if  $T$  is a 2-dimensional complex torus, and  $T^*$  its dual torus, the Kummer manifolds  $K_2(T)$  and  $K_2(T^*)$  (1.2.b) have the same period (with appropriate markings), but are not bimeromorphic in general. Thus we can only ask:

**Question 2.** *Let  $X, X'$  be two hyperkähler manifolds of the same dimension. If there is a Hodge isometry  $\lambda : H^2(X, \mathbb{Z}) \xrightarrow{\sim} H^2(X', \mathbb{Z})$ , what can we say of  $X$  and  $X'$ ? Can we conclude that  $X$  and  $X'$  are isomorphic by imposing that  $\lambda$  preserves some extra structure?*

One can formulate analogous questions for *polarized* hyperkähler manifolds, requiring that  $\lambda$  preserves the polarization classes. Here again Namikawa's construction provides a counter-example. Nevertheless the recent work of Markman [27] and Verbitsky [39] gives a partial answer to question 2 and its "polarized" analogue, in particular for the case of example 1.2.a); see also the discussion in [15], Question 2.6, and the paper of E. Markman in these Proceedings.

## 1.4 Cohomology

Let  $X$  be a hyperkähler manifold. Since the quadratic form  $q$  plays such an important role, it is natural to expect that it determines most of the cohomology of  $X$ . This was indeed shown by Bogomolov [10] :

**Proposition 1.** *Let  $X$  be a hyperkähler manifold, of dimension  $2r$ , and let  $\mathcal{H}$  be the subalgebra of  $H^*(X, \mathbb{C})$  spanned by  $H^2(X, \mathbb{C})$ .*

1)  $\mathcal{H}$  is the quotient of  $\text{Sym}^* H^2(X, \mathbb{C})$  by the ideal spanned by the classes  $\alpha^{r+1}$  for  $\alpha \in H^2(X, \mathbb{C})$ ,  $q_{\mathbb{C}}(\alpha) = 0$ .

2)  $H^*(X, \mathbb{C}) = \mathcal{H} \oplus \mathcal{H}^{\perp}$ , where  $\mathcal{H}^{\perp}$  is the orthogonal of  $\mathcal{H}$  with respect to the cup-product.

Thus the subalgebra  $\mathcal{H}$  is completely determined by the form  $q$  and the dimension of  $X$ . In contrast, not much is known about the  $\mathcal{H}$ -module  $\mathcal{H}^{\perp}$ . Note that it is nonzero for the examples a) and b) of 1.2, with the exception of  $S^{[2]}$  for a K3 surface  $S$ .

We do not know much about the quadratic form  $q$  either. For the two infinite series of (1.2) we have lattice isomorphisms [2]

$$H^2(S^{[r]}, \mathbb{Z}) = H^2(S, \mathbb{Z}) \oplus^{\perp} \langle 2-2r \rangle \quad H^2(K_r(T), \mathbb{Z}) = H^2(T, \mathbb{Z}) \oplus^{\perp} \langle -2-2r \rangle ;$$

The lattices of O'Grady's two examples are computed in [36]; they are also even.

**Question 3.** *Is the quadratic form  $q$  always even? More generally, what are the possibilities for  $q$ ? What are the possibilities for the Fujiki index  $f_X$  (see 1.3)?*

## 1.5 Boundedness

Having so few examples leads naturally to the following question:

**Conjecture 1.** *There are finitely many hyperkähler manifolds (up to deformation) in each dimension.*

Note that the same question can be asked for Calabi-Yau manifolds, but there it seems completely out of reach.

Huybrechts observes that there are finitely many deformation types of hyperkähler manifolds  $X$  of dimension  $2r$  such that there exists  $\alpha \in H^2(X, \mathbb{Z})$

with  $q(\alpha) > 0$  and  $\int_X \alpha^{2r}$  bounded [18]. As a corollary, given a real number  $M$ , there are finitely many deformation types of hyperkähler manifolds with

$$f_X \leq M \quad , \quad \min\{q(\alpha) \mid \alpha \in H^2(X, \mathbb{Z}) \text{ , } q(\alpha) > 0\} \leq M \text{ .}$$

A first approximation to finiteness would be to bound the Betti numbers  $b_i$  of  $X$ , and in particular  $b_2$ . Here we have some more information in the case of fourfolds [16] :

**Proposition 2.** *Let  $X$  be a hyperkähler fourfold. Then either  $b_2 = 23$ , or  $3 \leq b_2 \leq 8$ .*

Note that  $b_2$  is 23 for  $S^{[2]}$  and 7 for  $K_2(T)$  (1.2). [16] contains some more information on the other Betti numbers.

**Question 4.** *Can we exclude some more cases, in particular  $b_2 = 3$ ? If  $b_2 = 23$ , can we conclude that  $X$  is deformation equivalent to  $S^{[2]}$ ?*

## 1.6 Lagrangian fibrations

Let  $(X, \sigma)$  be a holomorphic symplectic manifold (not necessarily compact), of dimension  $2r$ . A *Lagrangian fibration* is a proper map  $h : X \rightarrow B$  onto a manifold  $B$  such that the general fibre  $F$  of  $h$  is Lagrangian, that is,  $F$  is connected, of dimension  $r$ , and  $\sigma|_F = 0$ . This implies that the smooth fibres of  $h$  are complex tori (Arnold-Liouville theorem).

Suppose  $B = \mathbb{C}^r$ , so that  $h = (h_1, \dots, h_r)$ . The functions  $h_i$  define what is called in classical mechanics an *algebraically completely integrable hamiltonian system* : the Poisson brackets  $\{h_i, h_j\}$  vanish, the hamiltonian vector fields  $X_{h_i}$  commute with each other, they are tangent to the fibres of  $h$  and their restriction to a smooth fibre is a linear vector field on this complex torus (see for instance [4]).

The analogue of this notion when  $X$  is compact (hence hyperkähler) is a Lagrangian fibration  $X \rightarrow \mathbb{P}^r$ . There are many examples of such fibrations (see a sample below); moreover they turn out to be the only non-trivial morphisms from a hyperkähler manifold to a manifold of smaller dimension :

**Theorem 4.** *Let  $X$  be a hyperkähler manifold, of dimension  $2r$ ,  $B$  a Kähler manifold with  $0 < \dim B < 2r$ , and  $f : X \rightarrow B$  a surjective morphism with connected fibres. Then:*

- 1)  *$f$  is a Lagrangian fibration;*
- 2) *If  $X$  is projective,  $B \cong \mathbb{P}^r$ .*

1) is due to Matsushita (see [19], Prop. 24.8), and 2) to Hwang [20]. It is expected that 2) holds without the projectivity assumption on  $X$  (see the discussion in the introduction of [20]).

How do we detect the existence of a Lagrangian fibration on a given hyperkähler manifold? In dimension 2 there is a simple answer; a Lagrangian fibration on a K3 surface  $S$  is an elliptic fibration, and we have :

**Proposition 3.** a) *Let  $L$  be a nontrivial nef line bundle on  $S$  with  $L^2 = 0$ . There exists an elliptic fibration  $f : S \rightarrow \mathbb{P}^1$  such that  $L = f^* \mathcal{O}_{\mathbb{P}^1}(k)$  for some  $k \geq 1$ .*

b)  *$S$  admits an elliptic fibration if and only if it admits a line bundle  $L \neq \mathcal{O}_S$  with  $L^2 = 0$ .*

The proof of a) is straightforward. b) is reduced to a) by proving that some isometry  $w$  of  $\text{Pic}(S)$  maps  $L$  to a nef line bundle; see for instance [1], VIII, Lemma 17.4.

Proposition 3 has a natural (conjectural) generalization to higher-dimensional hyperkähler manifolds<sup>3</sup> :

**Conjecture 2.** a) *Let  $L$  be a nontrivial nef line bundle on  $X$  with  $q(L) = 0$ . There exists a Lagrangian fibration  $f : X \rightarrow \mathbb{P}^r$  such that  $L = f^* \mathcal{O}_{\mathbb{P}^r}(k)$  for some  $k \geq 1$ .*

b) *There exists a hyperkähler manifold  $X'$  bimeromorphic to  $X$  and a Lagrangian fibration  $X' \rightarrow \mathbb{P}^r$  if and only if  $X$  admits a line bundle  $L \neq \mathcal{O}_S$  with  $q(L) = 0$ .*

Note that it is not clear whether one of the statements implies the other.

There is some evidence in favor of the conjecture. Let  $S$  be a “general” K3 surface of genus  $g$  – that is,  $\text{Pic}(S) = \mathbb{Z}[L]$  with  $L^2 = 2g - 2$ . Then  $\text{Pic}(S^{[r]})$  is a rank 2 lattice with an orthonormal basis  $(h, e)$  satisfying  $q(h) = 2g - 2$ ,  $q(e) = -(2r - 2)$  [2]. Taking  $r = g$  we find  $q(h \pm e) = 0$ . The corresponding Lagrangian fibration is studied in [3]:  $S^{[g]}$  is birational to the relative compactified Jacobian  $\mathcal{J}^g \rightarrow |L|$ , whose fibre above a curve  $C \in |L|$  is the compactified Jacobian  $\overline{J^g C}$ .  $\mathcal{J}^g$  is hyperkähler by [30], and the fibration  $\mathcal{J}^g \rightarrow |L|$  is Lagrangian. The rational map  $S^{[g]} \dashrightarrow |L|$  associates to a general set of  $g$  points in  $S$  the unique curve of  $|L|$  passing through these points.

More generally, suppose that  $2g - 2 = (2r - 2)m^2$  for some integer  $m$ . Then  $q(h \pm me) = 0$ , and indeed  $S^{[r]}$  admits a birational model with a Lagrangian

---

<sup>3</sup> The conjecture has been known to experts for a long time; see the introduction of [38] for a discussion of its history.

fibration. This fibration has been constructed independently in [28] and [37];  $\mathcal{J}^g$  is replaced by a moduli space of *twisted sheaves* on  $S$ .

Another argument in favor of the conjecture has been given by Matsushita [29], who proved that a) holds “locally”, in the following sense. Let  $X$  be a hyperkähler manifold, with a Lagrangian fibration  $f : X \rightarrow \mathbb{P}^r$ , and let  $\text{Def}(X)$  be the local deformation space of  $X$ . Then the Lagrangian fibration deforms along a hypersurface in  $\text{Def}(X)$ . Thus any small deformation of  $X$  such that the cohomology class of  $f^*\mathcal{O}_{\mathbb{P}^r}(1)$  remains algebraic carries a Lagrangian fibration.

A related question, which comes from mathematical physics, is :

**Question 5.** *Does every hyperkähler manifold admit a deformation with a Lagrangian fibration?*

If Conjecture 2 holds, the answer is positive if and only if the quadratic form  $q$  is indefinite. I do not know any serious argument either in favor or against this.

**Question 6.** *Let  $X$  be a hyperkähler manifold, and  $T \subset X$  a Lagrangian submanifold which is a complex torus. Is it the fibre of a Lagrangian fibration  $X \rightarrow \mathbb{P}^r$ ?*

(A less optimistic version would ask only for a bimeromorphic Lagrangian fibration.)

## 1.7 Projective families

Deformation theory shows that when the K3 surface  $S$  varies, the manifolds  $S^{[r]}$  form a hypersurface in their deformation space; thus a general deformation of  $S^{[r]}$  is not the Hilbert scheme of a K3 – and we do not know how to describe it. This is not particularly surprising: after all, we do not know either how to describe a general K3 surface. On the other hand, if we start from the family of *polarized* K3 surfaces  $S$  of genus  $g$ , the projective manifolds  $S^{[r]}$  are polarized (in various ways)<sup>4</sup>, and the same argument tells us that they form again a hypersurface in their (polarized) deformation space; we should be able to describe a (locally) complete family of projective hyperkähler manifolds which specializes to  $S^{[r]}$  in codimension 1.

<sup>4</sup> For  $S$  general we have  $\text{Pic}(S^{[r]}) = \mathbb{Z}h \oplus^{\perp} \mathbb{Z}e$  (1.6); the polarizations on  $S^{[r]}$  are of the form  $ah - be$  with  $a, b > 0$ .

For  $r = 2$  there are indeed a few cases where we can describe the general deformation of  $S^{[2]}$  with an appropriate polarization :

1. The Fano variety of lines contained in a cubic fourfold ([7];  $g = 8$ )
2. The “variety of sum of powers” associated to a cubic fourfold ([21];  $g = 8$ )
3. The double cover of certain sextic hypersurfaces in  $\mathbb{P}^5$  ([34];  $g = 6$ )
4. The subspace of the Grassmannian  $\mathbb{G}(6, V)$  consisting of 6-planes  $L$  such that  $\sigma|_L = 0$ , where  $\sigma : \wedge^3 \mathbb{C}^{10} \rightarrow \mathbb{C}$  is a sufficiently general 3-form ([12];  $g = 12$ ).

Note that K3 surfaces of genus 8 appear in both cases 1) and 2); what happens is that the corresponding polarizations on  $S^{[2]}$  are different [22]<sup>5</sup>.

**Question 7.** *Describe the general projective deformation of  $S^{[2]}$ , for  $S$  a polarized K3 surface of genus 1, 2, 3, ... (and for some choice of polarization on  $S^{[2]}$ ); or at least find more examples of locally complete projective families. Same question with  $S^{[r]}$  for  $r \geq 3$ .*

(With the notation of footnote 4, a natural choice of polarization for  $g \geq 3$  is  $h - e$ .)

A different issue concerns the *Chow ring* of a projective hyperkähler manifold. In [6] and [40] the following conjecture is proposed :

**Conjecture 3.** *Let  $D_1, \dots, D_k$  in  $\text{Pic}(X)$ , and let  $z \in CH(X)$  be a class which is a polynomial in  $D_1, \dots, D_k$  and the Chern classes  $c_i(X)$ . If  $z = 0$  in  $H^*(X, \mathbb{Z})$ , then  $z = 0$ .*

This would follow from a much more general (and completely out of reach) conjecture, for which we refer to the introduction of [6]. Conjecture 3 is proved in [40] for the Hilbert scheme  $S^{[n]}$  of a K3 surface for  $n \leq 8$ , and for the Fano variety of lines on a cubic fourfold.

## 2 Compact Poisson manifolds

Since hyperkähler manifolds are so rare, it is natural to turn to a more flexible notion. Symplectic geometry provides a natural candidate, *Poisson manifolds*. Recall that a (holomorphic) Poisson structure on a complex manifold  $X$  is a bivector field  $\tau \in H^0(X, \wedge^2 T_X)$ , such that the bracket  $\{f, g\} := \langle \tau, df \wedge dg \rangle$

<sup>5</sup> The Corollary in [22] is slightly misleading: the moduli spaces of *polarized* hyperkähler manifolds of type 1) and of type 2) are disjoint.

defines a Lie algebra structure on  $\mathcal{O}_X$ . A Poisson structure defines a skew-symmetric map  $\tau^\sharp : \Omega_X^1 \rightarrow T_X$ ; the *rank* of  $\tau$  at a point  $x \in X$  is the rank of  $\tau^\sharp(x)$ . It is even (because  $\tau^\sharp$  is skew-symmetric). The data of a Poisson structure of rank  $\dim X$  is equivalent to that of a (holomorphic) symplectic structure. In general, we have a partition

$$X = \coprod_{s \text{ even}} X_s \quad \text{where } X_s := \{x \in X \mid \text{rk } \tau(x) = s\} .$$

The following conjecture is due to Bondal ([11], see also [35]):

**Conjecture 4.** *If  $X$  is Fano and  $s$  even,  $X_{\leq s} := \coprod_{k \leq s} X_k$  contains a component of dimension  $> s$ .*

This is much larger than one would expect from a naive dimension count. It implies for instance that a Poisson field which vanishes at some point must vanish along a curve.

The condition “ $X$  Fano” is probably far too strong. In fact an optimistic modification would be :

**Conjecture 5.** *If  $X_s$  is non-empty, it contains a component of dimension  $> s$ .*

Here are some arguments in favor of this conjecture:

**Proposition 4.** *Let  $(X, \tau)$  be a compact Poisson manifold.*

- 1) *Every component of  $X_s$  has dimension  $\geq s$ .*
- 2) *Let  $r$  be the generic rank of  $\tau$  ( $r$  even); assume that  $c_1(X)^q \neq 0$  in  $H^q(X, \Omega_X^q)$ , where  $q = \dim X - r + 1$ . Then the degeneracy locus  $X - X_r$  of  $\tau$  has a component of dimension  $> r - 2$ .*
- 3) *Assume that  $X$  is a projective threefold. If  $X_0$  is non-empty, it contains a curve.*

*Sketch of proof :* 1) Let  $Z$  be a component of  $X_s$  (with its reduced structure). It is not difficult to prove that  $Z$  is a Poisson subvariety of  $X$  (see [35]); this means that at a smooth point  $x$  of  $Z$ , the tensor  $\tau(x)$  lives in  $\Lambda^2 T_x(Z) \subset \Lambda^2 T_x(X)$ . But this implies  $s \leq \dim T_x(Z) = \dim Z$ .

2) is proved in [35], §9, under the extra hypothesis  $\dim X = r + 1$ . The proof extends easily to the slightly more general situation considered here.

3) is proved in [14] by a case-by-case analysis (leading to a complete classification of those Poisson threefolds for which  $X_0 = \emptyset$ ). It would be interesting to have a more conceptual proof.

The paper [35] contains many interesting results on Poisson manifolds; in particular, a complete classification of the Poisson structures on  $\mathbb{P}^3$  for which the zero locus contains a smooth curve.

### 3 Compact contact manifolds

Let  $X$  be a complex manifold, of odd dimension  $2r+1$ . A *contact structure* on  $X$  is a one-form  $\theta$  with values in a line bundle  $L$  on  $X$ , such that  $\theta \wedge (d\theta)^r \neq 0$  at each point of  $X$  (though  $\theta$  is a twisted 1-form, it is easy to check that  $\theta \wedge (d\theta)^r$  makes sense as a section of  $K_X \otimes L^{r+1}$ ; in particular, the condition on  $\theta$  implies  $K_X = L^{-r-1}$ ).

There are only two classes of compact holomorphic contact manifolds known so far:

a) The projective cotangent bundle  $\mathbb{P}T_M^*$ , where  $M$  is any compact complex manifold;

b) Let  $\mathfrak{g}$  be a simple complex Lie algebra. The action of the adjoint group on  $\mathbb{P}(\mathfrak{g})$  has a unique closed orbit  $X_{\mathfrak{g}}$ : every other orbit contains  $X_{\mathfrak{g}}$  in its closure.  $X_{\mathfrak{g}}$  is a contact Fano manifold.

The following conjecture is folklore:

**Conjecture 6.** *Any projective contact manifold is of type a) or b).*

Half of this conjecture is now proved, thanks to [24] and [13]: a contact projective manifold is either Fano with  $b_2 = 1$ , or of type a). It is easily seen that a homogeneous Fano contact manifold is of type b), so we can rephrase Conjecture 6 as :

**Conjecture 7.** *A contact Fano manifold is homogeneous.*

I refer to [4] for some evidence in favor of this conjecture, and to [5] for its application to differential geometry, more specifically to quaternion-Kähler manifolds. These are Riemannian manifolds with holonomy  $\mathrm{Sp}(1)\mathrm{Sp}(r)$ ; they are Einstein manifolds, and in particular they have constant scalar curvature. Thanks to work of Salamon and LeBrun [25, 26], a positive answer to Conjecture 7 would imply:

**Conjecture 8.** *The only compact quaternion-Kähler manifolds with positive scalar curvature are homogeneous.*

These positive homogeneous quaternion-Kähler manifolds have been classified by Wolf [41] : there is one,  $M_{\mathfrak{g}}$ , for each simple complex Lie algebra  $\mathfrak{g}$ .

The link between Conjectures 7 and 8 is provided by the twistor space construction. To any quaternion-Kähler manifold  $M$  is associated a  $\mathbb{S}^2$ -bundle  $X \rightarrow M$ , the twistor space, which carries a natural complex structure; when  $M$  has positive scalar curvature it turns out that  $X$  is a contact Fano manifold

– for instance the twistor space of  $M_{\mathfrak{g}}$  is  $X_{\mathfrak{g}}$ . Conjecture 7 implies that  $X$  is isomorphic to  $X_{\mathfrak{g}}$  for some simple Lie algebra  $\mathfrak{g}$ ; this in turn implies that  $M$  is isometric to  $M_{\mathfrak{g}}$  and therefore homogeneous.

## References

1. W. Barth, K. Hulek, C. Peters, A. Van de Ven: *Compact complex surfaces*. 2nd edition. Ergebnisse der Mathematik und ihrer Grenzgebiete, **4**. Springer-Verlag, Berlin (2004).
2. A. Beauville : *Variétés kählériennes dont la première classe de Chern est nulle*. J. of Diff. Geometry **18**, 755–782 (1983).
3. A. Beauville : *Systèmes hamiltoniens complètement intégrables associés aux surfaces K3*. Problems in the theory of surfaces and their classification (Cortona, 1988), 25–31, Sympos. Math. **32**, Academic Press, London (1991).
4. A. Beauville : *Fano contact manifolds and nilpotent orbits*. Comment. Math. Helv. **73**, 566–583 (1998).
5. A. Beauville : *Riemannian holonomy and algebraic geometry*. Enseign. Math. (2) **53** (2007), no. 1-2, 97–126.
6. A. Beauville : *On the splitting of the Bloch-Beilinson filtration*. Algebraic cycles and motives (vol. 2), London Math. Soc. Lecture Notes **344**, 38–53; Cambridge University Press (2007).
7. A. Beauville, R. Donagi : *La variété des droites d’une hypersurface cubique de dimension 4*. C.R. Acad. Sc. Paris **301**, 703–706 (1985).
8. F. Bogomolov : *The decomposition of Kähler manifolds with a trivial canonical class*. Mat. Sbornik (N.S.) **93(135)** (1974), 573–575, 630.
9. F. Bogomolov : *Hamiltonian Kählerian manifolds*. Dokl. Akad. Nauk SSSR **243** (1978), no. 5, 1101–1104.
10. F. Bogomolov : *On the cohomology ring of a simple hyper-Kähler manifold (on the results of Verbitsky)*. Geom. Funct. Anal. **6** (1996), no. 4, 612–618.
11. A. Bondal : *Noncommutative deformations and Poisson brackets on projective spaces*. Preprint MPI/93-67.
12. O. Debarre, C. Voisin : *Hyper-Kähler fourfolds and Grassmann geometry*. Preprint arXiv:0904.3974.
13. J.-P. Demailly : *On the Frobenius integrability of certain holomorphic  $p$ -forms*. Complex geometry (Göttingen, 2000), 93–98, Springer, Berlin (2002).
14. S. Druel : *Structures de Poisson sur les variétés algébriques de dimension 3*. Bull. Soc. Math. France **127** (1999), no. 2, 229–253.
15. V. Gritsenko, K. Hulek, G. Sankaran : *Moduli spaces of irreducible symplectic manifolds*. Compositio Math. **146**, no. 2, 404–434.
16. D. Guan : *On the Betti numbers of irreducible compact hyperkähler manifolds of complex dimension four*. Math. Res. Lett. **8** (2001), no. 5-6, 663–669.
17. D. Huybrechts : *Compact hyper-Kähler manifolds: basic results*. Invent. Math. **135** (1999), no. 1, 63–113. Erratum: Invent. Math. **152** (2003), no. 1, 209–212.
18. D. Huybrechts : *Finiteness results for compact hyperkähler manifolds*. J. Reine Angew. Math. **558** (2003), 15–22.

19. D. Huybrechts : *Compact hyperkähler manifolds*. Calabi-Yau manifolds and related geometries (Nordfjordeid, 2001), 161–225, Universitext, Springer, Berlin (2003).
20. J.-M. Hwang : *Base manifolds for fibrations of projective irreducible symplectic manifolds*. Invent. Math. **174** (2008), no. 3, 625–644.
21. A. Iliev, K. Ranestad : *K3 surfaces of genus 8 and varieties of sums of powers of cubic fourfolds*. Trans. Amer. Math. Soc. **353** (2001), 1455–1468.
22. A. Iliev, K. Ranestad : *Addendum to “K3 surfaces of genus 8 and varieties of sums of powers of cubic fourfolds”*. C. R. Acad. Bulgare Sci. **60** (2007), 1265–1270.
23. D. Kaledin, M. Lehn, C. Sorger : *Singular symplectic moduli spaces*. Invent. Math. **164** (2006), no. 3, 591–614.
24. S. Kebekus, T. Peternell, A. Sommese, J. Wiśniewski : *Projective contact manifolds*. Invent. Math. **142** (2000), no. 1, 1–15.
25. C. LeBrun : *Fano manifolds, contact structures, and quaternionic geometry*. Int. J. of Math. **6** (1995), 419–437.
26. C. LeBrun, S. Salamon : *Strong rigidity of quaternion-Kähler manifolds*. Invent. math. **118** (1994), 109–132.
27. E. Markman : *Integral constraints on the monodromy group of the hyperkähler resolution of a symmetric product of a K3 surface*. Preprint arXiv:0601304. To appear in Internat. J. of Math.
28. D. Markushevich : *Rational Lagrangian fibrations on punctual Hilbert schemes of K3 surfaces*. Manuscripta Math. **120** (2006), no. 2, 131–150.
29. D. Matsushita : *On deformations of Lagrangian fibrations*. Preprint arXiv:0903.2098.
30. S. Mukai : *Symplectic structure of the moduli space of sheaves on an abelian or K3 surface*. Invent. Math. **77** (1984), no. 1, 101–116.
31. Y. Namikawa : *Counter-example to global Torelli problem for irreducible symplectic manifolds*. Math. Ann. **324** (2002), no. 4, 841–845.
32. K. O’Grady : *Desingularized moduli spaces of sheaves on a K3*. J. Reine Angew. Math. **512** (1999), 49–117.
33. K. O’Grady : *A new six-dimensional irreducible symplectic variety*. J. Algebraic Geom. **12** (2003), no. 3, 435–505.
34. K. O’Grady : *Irreducible symplectic 4-folds and Eisenbud-Popescu-Walter sextics*. Duke Math. J. **134** (2006), no. 1, 99–137.
35. A. Polishchuk : *Algebraic geometry of Poisson brackets*. J. Math. Sci. (New York) **84** (1997), 1413–1444.
36. A. Rapagnetta : *On the Beauville form of the known irreducible symplectic varieties*. Math. Ann. **340** (2008), no. 1, 77–95.
37. J. Sawon : *Lagrangian fibrations on Hilbert schemes of points on K3 surfaces*. J. Algebraic Geom. **16** (2007), no. 3, 477–497.
38. M. Verbitsky : *Hyperkähler SYZ conjecture and semipositive line bundles*. Preprint arXiv:0811.0639.
39. M. Verbitsky : *A global Torelli theorem for hyperkähler manifolds*. Preprint arXiv:0908.4121.
40. C. Voisin : *On the Chow ring of certain algebraic hyper-Kähler manifolds*. Pure Appl. Math. Q. **4** (2008), no. 3, part 2, 613–649.
41. J. Wolf : *Complex homogeneous contact manifolds and quaternionic symmetric spaces*. J. Math. Mech. **14** (1965), 1033–1047.

42. S.-T. Yau : *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation* I. *Comm. Pure and Appl. Math.* **31** (1978), 339–411.