

Superdiffusivity of asymmetric energy model in dimensions 1 and 2

Cédric Bernardin^{a)}

Université de Lyon, CNRS (UMPA) Ecole Normale Supérieure de Lyon, 46, allée d'Italie, 69364 Lyon Cedex 07, France

(Received 28 May 2008; accepted 25 September 2008; published online 22 October 2008)

We discuss an asymmetric energy model introduced by Giardina *et al.* [J. Math. Phys. **48**, 033301 (2007)]. This model is expected to belong to the Kardar–Parisi–Zhang (KPZ) class. We obtain lower bounds for the diffusion coefficient. In particular, the diffusion coefficient is diverging in dimensions one and two as it is expected in the KPZ picture. © 2008 American Institute of Physics.
[DOI: [10.1063/1.3000580](https://doi.org/10.1063/1.3000580)]

I. INTRODUCTION

In their well-known paper, Kardar *et al.*¹² introduced a model for the evolution of the profile of a growing interface,

$$\partial_t h = \nabla^2 h + \frac{1}{2}(\nabla h)^2 + \eta(x, t). \quad (1)$$

Here $h(x, t) \in \mathbb{R}$ is the height of the interface at location $x \in \mathbb{R}^d$ and at time $t > 0$ and $\eta(x, t)$ is a space-time white noise. Starting from a flat state at time $t=0$, they are interested in the evolution of the fluctuations

$$\xi(x, t) = \langle [h(x, t) - \langle h(x, t) \rangle]^2 \rangle^{1/2}.$$

The intuitive picture is that the width grows with time as a power law up to a saturation time that scales with the substrate size L as L^z , where z is the scaling exponent. In other words, we expect that

$$\xi(L, t) = L^{2-z} f(t/L^z),$$

where the scaling function $f(x)$ saturates at large x and $f(x) \sim x^{(2-z)/z}$ for $x \sim 0$.¹⁴ By dynamic renormalization-group techniques, Kardar *et al.* show that in dimension 1, the dynamic scaling exponents are $z=3/2$ and $z=2$ in dimension $d \geq 3$. The dimension $d=2$ is the critical one and their numerical studies indicate $1/z=0.62004$. As noticed by Kardar *et al.*, Eq. (1) can be mapped to the Burgers equation for a vorticity-free velocity field,

$$\partial_t v + v \cdot \nabla v = \nabla^2 v - \nabla \eta(x, t), \quad (2)$$

with $v = -\nabla h$ and $\eta(x, t)$ a space-time white noise.

Burgers equation is also closely connected to driven diffusive systems. Consider a diffusive system under constant uniform driving force described by a nonlinear Langevin equation. Then a quadratic order expansion of the density gives the Burgers equation (2). In Ref. 4, van Beijern *et al.* investigated the steady-state scattering function for driven diffusive systems with a single conserved density. Mode-coupling arguments predict that in dimension $d=1$ ($d=2$), density fluctuations spread as $t^{2/3}$ [$t(\log t)^{2/3}$], whereas they are of order $t^{1/2}$ (ordinary diffusion law) in dimension $d \geq 3$. Guided by ideas of universality, we expect that a large class of microscopic

^{a)}Electronic mail: cbernard@umpa.ens-lyon.fr.

models whose evolutions are in a suitable coarse time and length scale an approximation of the Kardar–Parisi–Zhang (KPZ) equation (1) (or the noisy Burgers equation) have a universal scaling exponent z and a universal scaling limit.

Asymmetric simple exclusion process (ASEP) is a natural discretization (see Ref. 21) of the stochastic Burgers equation. The dynamics are given by asymmetric random walks on \mathbb{Z}^d with a drift in some direction such that jumps of particles to occupied sites are forbidden (exclusion rule). It can be reinterpreted as a growth model which is a natural discretization of the KPZ equation. During the last decade, a lot of work has been accomplished to test the validity of universality predictions for ASEP. In the one dimensional case the value of the dynamical exponent $z=3/2$ has been confirmed. Not only the exponent but also the scaling function was obtained.⁸ Moreover the limit is the same as the largest eigenvalue distribution in the random matrix theory (see Ref. 6 and references therein for more informations). Nevertheless an important fact has to be mentioned: the methods used to obtain dynamical scaling exponent and limit distribution are very dependent of the specific properties of ASEP. Indeed the main results are valid and proven in the one dimensional case and for the totally ASEP (TASEP), which corresponds to nearest neighbor jumps in the right direction. These results cannot be carried for general ASEP. For example, Prähofer and Spohn¹⁹ computed the current fluctuations for the TASEP, but their proof does not work for other ASEP models. A more robust method has been introduced by Balázs and Seppäläinen² but is restricted to attractive systems. Even if they do not obtain the scaling limit function, they are able to establish the order of current for the nearest neighbor ASEP but not for general ASEP (obtained in Ref. 21 by generalized duality techniques). Recently this method has been developed in the context of the asymmetric zero range process (AZRP) (see Ref. 1).

To my knowledge, the only models belonging to the KPZ class and for which a rigorous proof of the scaling order has been obtained are ASEP, AZRP, polynuclear growth model, and related models.^{1,2,7,20} In dimension 2, the class is even more restrictive and the only rigorous result is the scaling order obtained by Yau for general ASEP.²⁶ Hence, the class of microscopic models for which one can rigorously prove they belong to the KPZ class is very small and it is hence of extreme importance to have simple models for which one can rigorously prove they are in the KPZ class.

The aim of this paper is to study a *nonattractive* model introduced by Giardinà *et al.* in Ref. 11 and to show that it presents anomalous behavior in low dimension as it is expected in the KPZ picture. In Ref. 11 Giardinà *et al.* considered *symmetric energy model* (AEM) (heat conduction model in their terminology) and show that the system has a dual process.

They also introduce an asymmetric generalization of the model that we call the AEM. AEM should belong to the KPZ universality class. It presents several analogies with ASEP but also differences.¹¹ In this paper, we develop generalized duality properties for AEM and obtain lower bounds for the bulk diffusion coefficient $D(t)$, i.e., the variance of the two point correlation function [see (3)]. KPZ approach predicts large time behavior of $D(t)$. In particular, $D(t)$ is expected to be divergent in dimensions 1 and 2 and finite in dimension $d \geq 3$. The goal of this article is to obtain lower bounds for $D(t)$ consistent with this (Theorem 1). The proof of this result is based on generalized duality techniques introduced by Landim and Yau in the context of ASEP. They have been developed in several directions but essentially for lattice gas dynamics. Our main sources of inspiration are given by Refs. 5, 16, 23, and 24. The key ingredients of generalized duality techniques are the following.

- If $(\mu_\rho)_\rho$ is a shift invariant family of stationary measures indexed by the conserved quantity ρ (e.g., the density), then $L^2(\mu_\rho)$ can be decomposed in an orthogonal sum $\bigoplus_{n \geq 0} \mathcal{H}_n$.
- The symmetric part S of the generator sends \mathcal{H}_n into \mathcal{H}_n (S conserves the degree) and is the generator of a reversible Markov process with a finite number of particles n .
- In general, the asymmetric part A of the generator does not conserve the degree but the action of A on each subspace \mathcal{H}_n is a bounded operator from $(\mathcal{H}_n, \|\cdot\|_1)$ into $(\bigoplus_{j=n-n_0}^{n+n_0} \mathcal{H}_j, \|\cdot\|_{-1})$, where n_0 is a fixed positive integer. Here $\|\cdot\|_{\pm 1}$ are Sobolev norms corresponding to the operator S (see Sec. IV).

The paper is organized as follows. In Sec. II, we define AEM and introduce the diffusion coefficient $D(t)$. Section III is devoted to the generalized duality properties of the process. Section IV contains the technical lemmas necessary for the proof of Theorem 1. The proofs of the main theorem for the 1, 2, and $d \geq 3$ cases are given in Secs. V–VII. The paper is ended by remarks in Sec. VIII.

II. THE AEM

The system is composed of atoms indexed by $x \in \mathbb{Z}^d$. The canonical basis of \mathbb{R}^d is denoted by (e_1, \dots, e_d) . Each atom has a momentum $p_x \in \mathbb{R}$. Momenta are exchanged during the stochastic evolution in such a way that kinetic energy is conserved. The generator $L = S + A$ of AEM is defined by

$$(Sf)(p) = \sum_{i=1}^d \sum_{x \in \mathbb{Z}^d} (p_{x+e_i} \partial_{p_x} - p_x \partial_{p_{x+e_i}})^2$$

and

$$(Af)(p) = \sum_{i=1}^d a_i \sum_{x \in \mathbb{Z}^d} p_x p_{x+e_i} (p_{x+e_i} \partial_{p_x} - p_x \partial_{p_{x+e_i}}).$$

Here $p = (p_x)_{x \in \mathbb{Z}^d}$ is an element of the state space $\Omega = \mathbb{R}^{\mathbb{Z}^d}$ and f is a smooth local function of p . Parameters $(a_i)_{i=1, \dots, d}$ regulate the strength of the asymmetry in each direction. Let

$$\mu_T(dp) = \otimes_{x \in \mathbb{Z}^d} (2\pi T)^{-1/2} \exp(-p_x^2/2T) dp_x$$

be the Gaussian product measure with temperature T . μ_T is an invariant probability measure for L . Moreover S is symmetric and A is antisymmetric in $L^2(\Omega, \mu_T)$. We fix now $T > 0$ and denote μ_T by $\langle \cdot \rangle$. Energy of site x is denoted by $E_x = p_x^2$. The formal total energy $\sum_{x \in \mathbb{Z}^d} E_x$ is a conserved quantity of the dynamics, and one has

$$L(E_x) = 2\Delta(E_x) + 2 \sum_{i=1}^d a_i \nabla_{e_i} (E_{x-e_i} E_x),$$

where Δ is the usual d -dimensional discrete Laplacian,

$$\Delta E_x = \sum_{i=1}^d \{E_{x+e_i} + E_{x-e_i} - 2E_x\},$$

and ∇_{e_i} is the discrete gradient in the direction e_i ,

$$\nabla_{e_i} E_x E_{x-e_i} = E_{x+e_i} E_x - E_x E_{x-e_i}.$$

The microscopic instantaneous current in the direction e_i is given by

$$j_{x, x+e_i}(p) = 2\nabla_{e_i}(E_x) + 2a_i E_x E_{x+e_i},$$

and one has the following microscopic continuity equation:

$$E_x(t) - E_x(0) = \sum_{i=1}^d \int_0^t (\nabla_{e_i} j_{x-e_i, x})(p(s)) ds + M_x(t),$$

where $M_x(t)$ is a martingale.

We are interested in the energy-energy correlation function $S(x, t)$ defined by

$$S(x, t) = \langle E_x(t)E_0(0) \rangle - T^2.$$

At equilibrium, the mean value of the current in the direction e_i is $j_i(T) = \mu_T(j_{x,x+e_i}) = 2a_i T^2$. By the conservation law (cf., Ref. 25, pp. 263–264), one has the two following informations for the average location and velocity of the structure function:

$$\sum_x S(x, t) = 2T^2$$

and

$$\frac{1}{2T^2} \sum_x x S(x, t) = tv,$$

where $v = \sum_{i=1}^d j'_i(T) e_i = 4T \sum_{i=1}^d a_i e_i$.

The third natural quantity to study is the bulk diffusion coefficient which is defined by

$$D_{i,j}(t) = \frac{1}{4T^2 t} \left\{ \sum_{x \in \mathbb{Z}^d} x_i x_j S(x, t) - 2T^2 (v_i t)(v_j t) \right\}. \tag{3}$$

Based on mode-coupling theory,⁴ it is expected that

$$D(t) \sim \begin{cases} t^{1/3}, & d = 1 \\ (\log t)^{2/3}, & d = 2 \\ 1, & d \geq 3, \end{cases}$$

for large t . Let $w_i(p)$ be the normalized current in the direction e_i ,

$$w_i(p) = j_{0,e_i}(p) - j_i(T) - j'_i(T)(p_0^2 - T),$$

or more explicitly

$$w_i(p) = 2(1 + a_i T) \nabla_{e_i}(E_0) + 2a_i(E_0 - T)(E_{e_i} - T).$$

For local functions f and g in $L^2(\Omega)$, we define the semi-inner product

$$\langle\langle f, g \rangle\rangle = \sum_{z \in \mathbb{Z}^d} (\langle \tau_z f, g \rangle - \langle f \rangle \langle g \rangle).$$

Here τ_z is the usual shift on Ω . The seminorm corresponding to $\langle\langle \cdot, \cdot \rangle\rangle$ is denoted by $\|\cdot\|$. Note that (discrete) gradient terms $g = \tau_x h - h$ and constants vanish in this norm.

A formal integration by parts gives the following formula for the diffusion coefficient (see Ref. 15):

$$D_{i,j}(t) = \frac{\delta_{i,j}}{2} + \frac{1}{4T^2} \left\langle \left\langle t^{-1/2} \int_0^t w_i(p(s)) ds, t^{-1/2} \int_0^t w_j(p(s)) ds \right\rangle \right\rangle. \tag{4}$$

The Laplace transform of the diffusion coefficient is then given by

$$\int_0^\infty e^{-\lambda t} t D_{i,j}(t) dt = \frac{1}{2\lambda^2} + \frac{1}{4T^2 \lambda^2} \langle\langle w_i, (\lambda - L)^{-1} w_j \rangle\rangle. \tag{5}$$

We have to mention that we do not have a rigorous proof of equality (4). In fact, even the existence of the equilibrium infinite volume dynamics is a nontrivial problem.⁹ Several methods exist in the literature (Refs. 10 and 18 and references therein) but they are not directly applicable and we plan to extend these methods for AEM in a future work. In the rest of the paper, we assume the following.

Assumption 1: The operator L defined on the set of local integrable smooth functions of Ω is closable and its closure also denoted by L is the generator of a strong Markov process. Moreover the set of local smooth integrable functions on Ω is a core for L .

This assumption is needed to define the dynamics from the generator and make sense of $(\lambda - L)^{-1}$. Moreover the fact that the set of local smooth functions is a core permits to restrict the set of functions appearing in the variational formula (15) to this one.

In the following right hand side of (4) will be used as definition of $D(t)$. For ASEP, the validity of (4) can be established by coupling techniques one can note translate in our context (see Ref. 15 for more informations on this subject). It follows that in terms of Laplace transform, behavior of $D(t)$ for large t is, in a Tauberian sense, equivalent to behavior for small λ of

$$\langle\langle w_i, (\lambda - L)^{-1} w_j \rangle\rangle.$$

For simplicity, we will restrict ourselves to the case $i=j$. We will prove the following theorem.

Theorem 1: There exists a constant $C > 0$ such that

$$\langle\langle w_i, (\lambda - L)^{-1} w_i \rangle\rangle \geq \begin{cases} C\lambda^{-1/4}, & d = 1 \\ C \log(|\log \lambda|), & d = 2 \\ C, & d \geq 3. \end{cases}$$

Moreover, if $d \geq 3$, we have also an upper bound

$$\langle\langle w_i, (\lambda - L)^{-1} w_i \rangle\rangle \leq C^{-1}.$$

In a Tauberian sense, this theorem means that $D(t) \geq Ct^{1/4}$ for $d=1$, $D_{i,i}(t) \geq \log(\log t)$ for $d=2$, and $C^{-1} \geq D_{i,i}(t) \geq C > 0$ for $d \geq 3$.

Observe that the assumptions are not so relevant when we are only interested in lower and upper bounds. Without any assumption one can obtain similar bounds if we define the diffusion coefficient by a finite volume limit procedure (see Ref. 3, Sec. 3, for a discussion). It means we can define $D(t)$ by the following limit (when it exists):

$$D_{i,j}(t) = \lim_{N \rightarrow \infty} D_{i,j}^N(t), \quad D_{i,j}^{(N)}(t) = \frac{1}{4T^2 t} \left\{ \sum_{x \in \mathbb{T}_N^d} x_i x_j S(x, t) - 2T^2 (v_i t)(v_j t) \right\},$$

where \mathbb{T}_N^d is the d -dimensional discrete torus of length N and the dynamics is now defined on $\mathbb{R}^{\mathbb{T}_N^d}$. Then one can prove (without any assumption) the following lower bounds:

$$\lambda^2 \int_0^\infty e^{-\lambda t} D_{i,j}^{(N)}(t) dt \geq \begin{cases} C\lambda^{-1/4}, & d = 1 \\ C \log(|\log \lambda|), & d = 2 \\ C, & d \geq 3, \end{cases}$$

where $C > 0$ is independent of λ and N . Moreover we have the corresponding upper bound if $d \geq 3$. The advantage to deal directly with the infinite volume definition of the diffusion coefficient is that it simplifies notations and avoids working with discrete Fourier transform but directly with continuous Fourier transform.

III. DUALITY

For simplicity, we fix the temperature T equal to one and we denote by μ or $\langle \cdot \rangle$ the standard Gaussian product measure μ_1 . The Hilbert space associated with μ is denoted by $L^2(\Omega)$ and the corresponding inner product is denoted by $\langle \cdot, \cdot \rangle$. Let $h_n, n \geq 0$ be the sequence of Hilbert polynomials (see the Appendix) which is an orthogonal basis in $L^2(\mathbb{R}, (2\pi)^{-1/2} e^{-x^2/2} dx)$,

$$\int_{\mathbb{R}} h_n(x) h_m(x) \frac{e^{-x^2/2}}{\sqrt{2\pi}} = \delta_{m,n} n!, \quad n, m \geq 0.$$

The dual space of AEM is $\mathbb{N}^{\mathbb{Z}^d}$ and elements ξ of $\mathbb{N}^{\mathbb{Z}^d}$ are seen as configurations of the “generalized dual process.” If $\xi=(\xi_x; x \in \mathbb{Z}^d)$ is a configuration of the dual space, we will say that ξ is local if $\xi_x \neq 0$ only for a finite number of sites $x \in \mathbb{Z}^d$. In such a case the number of particles $|\xi|$ is defined by

$$|\xi| = \sum_{x \in \mathbb{Z}^d} \xi_x.$$

For any local configuration $\xi: \mathbb{Z}^d \rightarrow \mathbb{N}$, we define the multivariate Hermite polynomial function H_ξ by

$$H_\xi(p) = \prod_{x \in \mathbb{Z}^d} \frac{h_{\xi_x}(p_x)}{n(\xi_x)},$$

where $n: \mathbb{N} \rightarrow \mathbb{N} \setminus \{0\}$ is a suitable normalization function we will precise later. Remark that

$$\langle H_\xi, H_\eta \rangle = \frac{\xi!}{n^2(\xi)} \delta_{\xi, \eta}.$$

Here, the notations $\xi!$ and $n(\xi)$ are for $\prod_x (\xi_x!)$ and $\prod_x n(\xi_x)$.

Any local function $f \in \mathbb{L}^2(\Omega)$ can be decomposed uniquely as a finite linear combination of local functions

$$f = \sum_{\xi} F(\xi) H_\xi = \sum_{n \geq 0} \sum_{|\xi|=n} F(\xi) H_\xi.$$

The coefficients of this linear combination are given by a real valued function F defined on the dual space $\mathbb{N}^{\mathbb{Z}^d}$. Such a function is said to be of degree $n \geq 0$ if $F(\xi) = 0$ as soon as $|\xi| \neq n$. For example, the normalized current $w_i(p)$ has the following decomposition:

$$w_i = 2a_i \sum_{\xi} \delta_{\xi_i}(\xi) H_\xi + \text{gradient term}, \tag{6}$$

where ξ_i is the configuration with two particles on site 0 and two particles on site e_i ,

$$(\xi_i)_x = \begin{cases} 2 & \text{if } x = 0 \\ 2 & \text{if } x = e_i \\ 0 & \text{otherwise.} \end{cases}$$

We now examine how the generator acts on the dual space. For this purpose we need to introduce notations. We introduce a cemetery configuration “o” which does not belong to the dual space. Any function $F: \mathbb{N}^{\mathbb{Z}^d} \rightarrow \mathbb{R}$ is extended to a function F on $\mathbb{N}^{\mathbb{Z}^d} \cup \{o\}$ by $F(o) = 0$. If ξ is a configuration belonging to the dual space $\mathbb{N}^{\mathbb{Z}^d}$ and x, y are two sites of \mathbb{Z}^d then $\xi^{x,+2,y,-2}$ is the configuration obtained from ξ by moving two particles from site y to site x . If $\xi_y \leq 1$ then $\xi^{x,+2,y,-2} = o$. The configuration $\xi^{x,+2}$ ($\xi^{x,-2}$) is the configuration obtained from ξ by adding (by removing) two particles on site x . In the second case, if $\xi_x \leq 1$ then $\xi^{x,-2} = o$. If $f \in \mathbb{L}^2(\Omega)$ is a smooth local function such that

$$f = \sum_{\xi \in \mathbb{N}^{\mathbb{Z}^d}} F(\xi) H_\xi = \sum_{n \geq 0} \sum_{|\xi|=n} F(\xi) H_\xi,$$

then

$$Lf = \sum_{i=1}^d \{S_i f + a_i A_i f\} = \sum_{i=1}^d \left\{ \sum_{\xi} (S_i F)(\xi) H_{\xi} + a_i \sum_{\xi} (A_i F)(\xi) H_{\xi} \right\}.$$

We take

$$n(k) = k! = k(k-2)(k-4), \dots, \quad n(0) = 1.$$

A long but elementary computation (see Appendix) shows that

$$(S_i F)(\xi) = \sum_x (\xi_x + 1) \xi_{x-e_i} [F(\xi^{x+2, x-e_i-2}) - F(\xi)] + \sum_x (\xi_x + 1) \xi_{x+e_i} [F(\xi^{x+2, x+e_i-2}) - F(\xi)] \tag{7}$$

and

$$A_i = A_i^0 + A_i^+ + A_i^-,$$

with

$$(A_i^0 F)(\xi) = \sum_x [(\xi_{x+e_i} + 1) \xi_x F(\xi^{x, -2, x+e_i+2}) - (\xi_{x-e_i} - 1) \xi_x F(\xi^{x, -2, x-e_i+2})] \tag{8}$$

and

$$(A_i^+ F)(\xi) = \sum_x \xi_x (\xi_{x+e_i} - \xi_{x-e_i}) F(\xi^{x, -2}) \tag{9}$$

and

$$(A_i^- F)(\xi) = \sum_x (\xi_x + 1) (\xi_{x+e_i} - \xi_{x-e_i}) F(\xi^{x, 2}). \tag{10}$$

Remark that $\mathcal{S} = \sum_{i=1}^d S_i$ is the generator of a Markov process reversible with respect to the measure

$$m(\xi) = \frac{\xi!}{n^2(\xi)}.$$

This Markov process is not irreducible, but if we restrict this process to the invariant subspace,

$$\mathcal{E}_{\varepsilon, n} = \{\xi; |\xi| = n, \xi_x = \varepsilon_x \pmod{2}\}, \quad \varepsilon \in \{0, 1\}^{\mathbb{Z}^d},$$

then the restriction is irreducible. Let $\mathcal{H}_{\varepsilon, n}$ be the subspace of functions F vanishing outside $\mathcal{E}_{\varepsilon, n}$.

Remark that the process corresponding to the generator \mathcal{S} is the same as the dual process derived in Ref. 11. Nevertheless it is important to observe that our basis $\{H_{\xi}\}$ is different from the basis of Ref. 11. The key advantage of our choice is that the basis $\{H_{\xi}\}$ is orthogonal and the computations simplify considerably.

The operators A_i^0 , A_i^+ , and A_i^- are not Markov generators. The operator A_i^0 conserves the degree and is antisymmetric in $L^2(m)$. The operator A_i^+ increases the degree by 2 and A_i^- decreases the degree by 2. In $L^2(m)$, one has $(A_i^-)^* = -A_i^+$.

We will use the following abusive but very convenient notations in the sequel. If $f, g \in L^2(\Omega)$ are smooth local functions with coefficients in the basis H_{ξ} given by local functions F and G , then we have

$$\langle f, g \rangle = \langle F, G \rangle,$$

with $\langle F, G \rangle$ defined by

$$\langle F, G \rangle = \sum_{\xi \in \mathbb{N}^{\mathbb{Z}^d}} F(\xi)G(\xi)m(\xi).$$

We write similarly

$$\langle\langle f, g \rangle\rangle = \langle\langle F, G \rangle\rangle,$$

with

$$\langle\langle F, G \rangle\rangle = \sum_{x \in \mathbb{Z}^d} \sum_{\xi \in \mathbb{N}^{\mathbb{Z}^d} \setminus \{0\}} F(\xi)G(\tau_x \xi)m(\xi), \tag{11}$$

where $\tau_x \xi$ is the shifted configuration by x , meaning $(\tau_x \xi)_z = \xi_{x+z}$. Remark that in (11), the sum is carried over configurations ξ with at least one particle.

IV. FREE PARTICLE APPROXIMATION

We introduce the generator Δ (discrete Laplacian) of independent random walks on \mathbb{Z}^d . It is given by

$$(\Delta G)(\eta) = \sum_{\substack{x, y \in \mathbb{Z}^d \\ |x-y|=1}} \eta_x [G(\eta^{x,-1,y,+1}) - G(\eta)].$$

Here $\eta \in \mathbb{N}^{\mathbb{Z}^d}$ and $\eta^{x,-1,y,+1}$ is the configuration obtained from η by moving a particle from site x to site y . G is a real valued local function defined on $\mathbb{N}^{\mathbb{Z}^d}$. A configuration η can be seen as an element of $(\mathbb{Z}^d)/\Sigma_n$, where Σ_n is the symmetric group of order n . The identification is given by the map

$$(x_1, \dots, x_n) \in (\mathbb{Z}^d)^n/\Sigma_n \rightarrow \eta,$$

with

$$\eta(x) = \sum_{i=1}^n \delta_{x_i}(x).$$

A function G in the domain of Δ is then identified with an element of $\oplus_{n=1}^{\infty} \mathbb{L}^2((\mathbb{Z}^d)^n/\Sigma_n)$ (with respect to the counting measure). We denote the standard inner product on $\oplus_{n=1}^{\infty} \mathbb{L}^2((\mathbb{Z}^d)^n/\Sigma_n)$ by $\langle \cdot, \cdot \rangle_0$,

$$\forall F, G \in \mathbb{L}^2((\mathbb{Z}^d)^n/\Sigma_n), \quad \langle F, G \rangle_0 = \sum_{\mathbf{x}=(x_1, \dots, x_n) \in (\mathbb{Z}^d)^n/\Sigma_n} F(\mathbf{x})G(\mathbf{x}) = \frac{1}{n!} \sum_{\mathbf{x}=(x_1, \dots, x_n) \in (\mathbb{Z}^d)^n} F(\mathbf{x})G(\mathbf{x}). \tag{12}$$

We define also the scalar product with translations associated with $\langle \cdot, \cdot \rangle_0$,

$$\langle\langle F, G \rangle\rangle_0 = \sum_{x \in \mathbb{Z}^d} (\langle F, \tau_x G \rangle_0 - \langle F \rangle_0 \langle G \rangle_0).$$

We recall that Δ is a positive self-adjoint operator with respect to $\langle \cdot, \cdot \rangle_0$, and that for any local function G , we have

$$\langle -\Delta G, G \rangle_0 = \frac{1}{2} \sum_{\eta, |x-y|=1} \frac{\eta_x}{\eta!} [G(\eta^{x,-1,y,+1}) - G(\eta)]^2.$$

Lemma 1: Let $F(\xi)$ be a local function belonging to $\mathcal{H}_{0,2n}$ and define $G(\eta) = F(2\eta)$, then

$$C^{-1}2^{-2n}\langle -\Delta G, G \rangle_0 \leq \langle -SF, F \rangle \leq Cn\langle -\Delta G, G \rangle_0,$$

where C is a positive constant independent of n and F .

Proof: One has

$$\langle -SF, F \rangle = \sum_{\xi \in \mathcal{E}_{0,n}} \sum_{|x-y|=1} \xi_x(1 + \xi_y)[F(\xi^{x,-2,y,+2}) - F(\xi)]^2 m(\xi).$$

For any integer k , we have

$$m(2k) = \frac{(2k)!}{(2k!!)^2} = \frac{\prod_{j=1}^k (2j-1)}{\prod_{j=1}^k 2j} = \prod_{j=1}^k \left(1 - \frac{1}{2j}\right)$$

and

$$m(2k+1) = \frac{(2k+1)!}{((2k+1)!!)^2} = \frac{\prod_{j=1}^k (2j)}{\prod_{j=1}^k (2j+1)} = \prod_{j=1}^k \left(1 - \frac{1}{2j+1}\right)$$

so that $2^{-k} \leq m(2k) \leq 1$ and $(2/3)^k \leq m(2k+1) \leq 1$. It follows that

$$2^{-2n} \leq \inf_{\xi \in \mathcal{E}_{0,2n}} m(\xi) \leq \sup_{\xi \in \mathcal{E}_{0,2n}} m(\xi) \leq 1.$$

We use now the fact that $1 \leq (1 + \xi_y) \leq (1 + 2n)$ and we conclude. □

It follows that the same lemma is also true for the inner product with translations since

$$\langle \langle F, G \rangle \rangle = \lim_{k \rightarrow \infty} \frac{1}{(2k+1)^d} \left\langle \left(\sum_{|x| \leq k} \tau_x(F - \langle F \rangle) \right), \left(\sum_{|x| \leq k} \tau_x(G - \langle G \rangle) \right) \right\rangle \tag{13}$$

and

$$\langle \langle F, G \rangle \rangle_0 = \lim_{k \rightarrow \infty} \frac{1}{(2k+1)^d} \left\langle \left(\sum_{|x| \leq k} \tau_x(F - \langle F \rangle_0) \right), \left(\sum_{|x| \leq k} \tau_x(G - \langle G \rangle_0) \right) \right\rangle. \tag{14}$$

Lemma 2: Let $F(\xi)$ a local function belonging to $\mathcal{H}_{0,2n}$ and define $G(\eta) = F(2\eta)$. There exists a constant C independent of n and F such that

$$C^{-1}2^{-2n}\langle -\Delta G, G \rangle_0 \leq \langle -SF, F \rangle \leq Cn\langle -\Delta G, G \rangle_0.$$

The $H_{-1,\lambda}$ norms are defined by

$$\|F\|_{-1,\lambda}^2 = \langle \langle F, (\lambda - S)^{-1}F \rangle \rangle = \sup_G \{2\langle \langle F, G \rangle \rangle - \langle \langle G, (\lambda - S)G \rangle \rangle\}$$

and

$$\|F\|_{-1,\lambda,0}^2 = \langle \langle F, (\lambda - \Delta)^{-1}F \rangle \rangle_0 = \sup_G \{2\langle \langle F, G \rangle \rangle_0 - \langle \langle G, (\lambda - \Delta)G \rangle \rangle_0\}.$$

In these formulas, function F is a local function from \mathbb{N}^d into \mathbb{R} and the supremum is carried over local functions.

It follows easily from lemma 2 and Eq. (12) that there exists a positive constant $C(n)$ such that for every local square integrable function F belonging to $\mathcal{H}_{0,2n}$,

$$\frac{1}{C(n)} \|mF\|_{-1,\lambda,0}^2 \leq \|F\|_{-1,\lambda}^2 \leq C(n) \|mF\|_{-1,\lambda,0}^2.$$

In this inequality, the function mF is defined by $(mF)(\xi) = m(\xi)F(\xi)$.

Every function belonging to $\mathcal{H}_{0,2n}$ can be seen as a symmetric function \tilde{F} from $(\mathbb{Z}^d)^n$ into \mathbb{R} ,

$$\tilde{F}(x_1, \dots, x_n) = F(2\delta_{x_1} + \dots + 2\delta_{x_n}),$$

where δ_x denotes the configuration with only one particle on site x . The sum of two configurations and multiplication by an integer are defined in the standard way. In the sequel, we will identify F with \tilde{F} .

To obtain a lower bound, we use the following variational formula for the Laplace transform (see Ref. 5):

$$\begin{aligned} \langle\langle w_i, (\lambda - L)^{-1} w_i \rangle\rangle &= \sup_f \{2\langle\langle f, w_i \rangle\rangle - \langle\langle f, (\lambda - S)f \rangle\rangle - \langle\langle Af, (\lambda - S)^{-1} Af \rangle\rangle\} \\ &= \sup_F \{2\langle\langle F, \delta_{\xi_i} \rangle\rangle - \|F\|_{1,\lambda}^2 - \|AF\|_{-1,\lambda}^2\}, \end{aligned} \tag{15}$$

and we restrict the supremum over smooth local functions $f = \sum_{\xi} F(\xi) H_{\xi}$ such that F belongs to $\mathcal{H}_{0,4}$.

Lemma 3: *Let $d \geq 1$ and $F, G: (\mathbb{Z}^d)^n \rightarrow \mathbb{R}$ be symmetric local functions. There exists a constant $C := C(d, n)$ such that*

$$|\langle G, \mathcal{A}^0 F \rangle| \leq C \langle G, (\lambda - S)G \rangle^{1/2} \langle F, (\lambda - S)F \rangle^{1/2}.$$

The same inequality is valid for the inner product $\langle\langle \cdot, \cdot \rangle\rangle$,

$$|\langle\langle G, \mathcal{A}^0 F \rangle\rangle| \leq C \langle\langle G, (\lambda - S)G \rangle\rangle^{1/2} \langle\langle F, (\lambda - S)F \rangle\rangle^{1/2}.$$

Proof: By definition of m , for any non-negative integers k, ℓ , we have

$$m(k+2)(k+2)m(\ell-2)(\ell-2) = (k+1)m(k)\ell m(\ell), \tag{16}$$

and by simple changes of variables, one has

$$\begin{aligned} \langle \mathcal{A}_i^0 F, G \rangle &= \frac{1}{2} \sum_{\xi, x} \xi_x (\xi_{x+e_i} + 1) [F(\xi^{x-2, x+e_i+2}) - F(\xi)] [G(\xi) - G(\xi^{x-2, x+e_i+2})] m(\xi) \\ &\quad - \frac{1}{2} \sum_{\xi, x} \xi_x (\xi_{x-e_i} + 1) [F(\xi^{x-2, x-e_i+2}) - F(\xi)] [G(\xi) - G(\xi^{x-2, x-e_i+2})] m(\xi). \end{aligned}$$

The estimate is then a simple consequence of Schwarz's inequality. The second part of the lemma follows from the definition of $\langle\langle \cdot, \cdot \rangle\rangle$. □

Corollary 1: *Let $d \geq 1$ and $F: (\mathbb{Z}^d)^n \rightarrow \mathbb{R}$ be a symmetric local function. There exists a constant $C = C(d, n) > 0$ such that*

$$\|\mathcal{A}^0 F\|_{-1,\lambda}^2 \leq C \|F\|_{1,\lambda}^2.$$

Recall we want to obtain a lower bound for the formula (15). Corollary 1 implies we can forget the term $\|\mathcal{A}^0 F\|_{-1,\lambda}^2$ in the variational formula (15). This is because by Schwarz inequality there exists a positive constant C , such that for any test function F , we have

$$2\langle\langle F, \delta_{\xi_i} \rangle\rangle - \|F\|_{1,\lambda}^2 - \|AF\|_{-1,\lambda}^2 \leq 2\langle\langle F, \delta_{\xi_i} \rangle\rangle - (C+1)\|F\|_{1,\lambda}^2 - \left\| \sum_{i=1}^d a_i (\mathcal{A}_i^+ F + \mathcal{A}_i^- F) \right\|_{-1,\lambda}^2.$$

V. THE ONE DIMENSIONAL CASE

We recall that every function F belonging to $\mathcal{H}_{0,2n}$ is identified with a symmetric function \tilde{F} from \mathbb{Z}^n into \mathbb{R} ,

$$\tilde{F}(x_1, \dots, x_n) = F(2\delta_{x_1} + \dots + 2\delta_{x_n}).$$

In the sequel, we forget the \sim and consider a function $F \in \mathcal{H}_{0,n}$ as a symmetric function on \mathbb{Z}^n .

To obtain a lower bound, we restrict the supremum over functions $f = \sum_{\xi} F(\xi) H_{\xi}$ such that F belongs to $\mathcal{H}_{0,4}$. We identify hence F with a symmetric function on \mathbb{Z}^2 .

By Corollary 1, $\|\mathcal{A}^0 F\|_{-1,\lambda}^2$ is of no importance and can be forgotten.

Since $F \in \mathcal{H}_{0,4}$, we have that $\mathcal{A}^+ F$ belongs to $\mathcal{H}_{0,6}$. $\mathcal{A}^+ F$ is identified with a symmetric function on \mathbb{Z}^3 .

For all symmetric function $F(x, y)$ ($G(x, y, z)$) from \mathbb{Z}^2 (\mathbb{Z}^3) to \mathbb{R} , we define

$$F^*(\alpha) = \sum_{z \in \mathbb{Z}} F(z, z + \alpha),$$

respectively,

$$G^*(\alpha, \beta) = \sum_{z \in \mathbb{Z}} G(\alpha + z, \beta + z, z).$$

One can check that

$$\langle\langle F, F \rangle\rangle_0 = \sum_{\alpha \in \mathbb{Z}} (F^*(\alpha))^2, \quad \langle\langle G, G \rangle\rangle_0 = \sum_{\alpha, \beta \in \mathbb{Z}} (G^*(\alpha, \beta))^2$$

and

$$\langle\langle F, (-\Delta F) \rangle\rangle_0 = \sum_{u \in \mathbb{Z}} \sum_e [F^*(u+e) - F^*(u)]^2, \quad \langle\langle G, (-\Delta G) \rangle\rangle_0 = \sum_{u \in \mathbb{Z}^2} \sum_e [G^*(u+e) - G^*(u)]^2, \quad (17)$$

where the sum over e is carried over all $e \in \mathbb{Z}^2$ such that $e = \pm(0, 1), \pm(1, 0), \pm(1, 1)$.

The advantage of this notation is to reduce the degree of functions. We take now a function $F \in \mathcal{H}_{0,4}$ (i.e., a symmetric function on \mathbb{Z}^2) and we compute $G^* = (m\mathcal{A}^+ F)^*$, where \mathcal{A}^+ is defined by (9). One has

$$(m\mathcal{A}^+ F)(x, x, x+1) = \frac{3}{2}(F(x, x+1) - F(x, x)),$$

$$(m\mathcal{A}^+ F)(x, x, x-1) = \frac{3}{2}(F(x, x) - F(x-1, x)),$$

$$(m\mathcal{A}^+ F)(x-1, x, x+1) = \frac{1}{2}(F(x, x+1) - F(x-1, x)),$$

$$(m\mathcal{A}^+ F)(x, x+1, y) = \frac{1}{2}(F(x+1, y) - F(x, y)), \quad y \neq x-1, x, x+1, x+2,$$

$$(m\mathcal{A}^+ F)(x, y, z) = 0 \quad \text{otherwise.}$$

A long but elementary computation shows that

$$\begin{aligned}
 G^*(x,y) = & \frac{1}{4}[\delta_1(y-x) - \delta_1(x-y)](F^*(y) - F^*(x)) - \frac{1}{4}\delta_1(|x|)\delta_1(y-x)(F^*(x+1) - F^*(x)) \\
 & - \frac{1}{4}\delta_1(|y|)\delta_1(x-y)(F^*(y+1) - F^*(y)) + \frac{3}{2}(\delta_1(x)\delta_1(y) - \delta_{-1}(x)\delta_{-1}(y))(F^*(0) - F^*(1)) \\
 & + \frac{1}{4}\delta_{-1}(x)\delta_{\geq 2}(y)(F^*(y-1) - F^*(y)) + \frac{1}{4}\delta_{-1}(y)\delta_{\geq 2}(x)(F^*(x-1) - F^*(x)) \\
 & + \frac{1}{4}\delta_1(x)\delta_{\geq 3}(y)(F^*(y) - F^*(y+1)) + \frac{1}{4}\delta_1(y)\delta_{\geq 3}(x)(F^*(x) - F^*(x+1)). \tag{18}
 \end{aligned}$$

We choose now the following test function F :

$$F^*(x) = \lambda^{-1/4} e^{-\lambda^{3/4}|x|} \tag{19}$$

and evaluate for this test function the terms appearing in the variational formula (15). Remark that variational formula (15) is defined by taking the supremum over local smooth functions. F is not local but a standard approximation procedure can be applied. In the sequel, C denotes a positive constant independent of λ which can change from line to line.

By using the decomposition (6) of w_i , we have that for small $\lambda > 0$,

$$\langle\langle \delta_{\xi_1}, F \rangle\rangle = 2a_1 \sum_x F(x, x+1) = 2a_1 F^*(1) \sim \lambda^{-1/4}.$$

Moreover, since F belongs to $\mathcal{H}_{0,4}$, there exists a positive constant C such that

$$C^{-1} \langle F, F \rangle_0 \leq \langle F, F \rangle \leq C \langle F, F \rangle_0,$$

and by (13) and (14), the same is true for the inner products $\langle\langle \cdot, \cdot \rangle\rangle_0 / \langle\langle \cdot, \cdot \rangle\rangle$. The norm of F with respect to $\langle\langle \cdot, \cdot \rangle\rangle_0$ is easy to evaluate and is of the order $\lambda^{-1/4}$. The third term to estimate is $\langle\langle F, -SF \rangle\rangle$. By Lemma 2, this term is of the same order as $\langle\langle F, -SF \rangle\rangle_0$. Thanks to formula (17) and explicit form of F^* , we obtain that $\langle\langle F, (-S)F \rangle\rangle$ is of the order $\lambda^{1/4}$. Hence we proved

$$2\langle\langle \delta_{\xi_1}, F \rangle\rangle \sim \lambda^{-1/4}, \quad \lambda \langle\langle F, F \rangle\rangle \sim \lambda^{-1/4}, \quad \langle\langle F, -SF \rangle\rangle \sim \lambda^{1/4}. \tag{20}$$

We now evaluate the last term

$$\langle\langle [m(\mathcal{A}^+F)], (\lambda - \Delta)^{-1} [m(\mathcal{A}^+F)] \rangle\rangle_0.$$

Lemma 4: Let $F^*(x) = \lambda^{-1/4} e^{-\lambda^{3/4}|x|}$. We have

$$\langle\langle [m(\mathcal{A}^+F)], (\lambda - \Delta)^{-1} [m(\mathcal{A}^+F)] \rangle\rangle_0 \sim \lambda^{-1/4}. \tag{21}$$

Proof: For any function $f: \mathbb{Z}^n \rightarrow \mathbb{R}$, we introduce the Fourier transform \hat{f} of f defined by

$$\hat{f}(\xi) = \sum_{x \in \mathbb{Z}^n} f(x) e^{2i\pi x \cdot \xi}, \quad \xi \in [0, 1]^n.$$

By (17), we have

$$\langle\langle [m(\mathcal{A}^+F)], (\lambda - \Delta)^{-1} [m(\mathcal{A}^+F)] \rangle\rangle_0 = \int_{[0,1]^2} ds dt \frac{|\hat{G}^*(s,t)|^2}{\lambda + 4\theta(s) + 4\theta(t) + 4\theta(s+t)},$$

where $\theta(u) = \sin^2(\pi u)$. We have

$$\langle\langle [m(\mathcal{A}^+F)], (\lambda - \Delta)^{-1}[m(\mathcal{A}^+F)] \rangle\rangle_0 \leq \int_{[0,1]^2} dsdt \frac{|\hat{G}^*(s,t)|^2}{\lambda + 4\theta(s) + 4\theta(t)} = \langle G^*, (\lambda - \Delta)^{-1}G^* \rangle_0.$$

We can express Fourier transform of G^* in terms of \hat{F}^* . We write $G^* = G_1^* + G_2^* + \dots + G_8^*$ with G_1^*, \dots, G_8^* the eight terms appearing in (18). We claim that

$$\langle\langle G_j, (\lambda - \Delta)^{-1}G_j \rangle\rangle_0 \sim \lambda^{-1/4}, \quad j = 1, 5, 6, 7, 8,$$

$$\langle\langle G_j, (\lambda - \Delta)^{-1}G_j \rangle\rangle_0 = O(\lambda \log \lambda), \quad j = 2, 3, 4.$$

We begin by the proof of the first claim. We have

$$G_1^*(x, y) = \frac{1}{4}[\delta_1(y - x) - \delta_1(x - y)](F^*(y) - F^*(x))$$

and

$$\hat{G}_1^*(s, t) = -\frac{i}{2}\{\sin(2\pi s) + \sin(2\pi t)\}\hat{F}^*(s + t).$$

The Fourier transform of \hat{F}^* is easy to compute and is given by

$$\hat{F}^*(s) = \frac{\lambda^{-1/4}(1 - 2e^{-2\lambda^{3/4}})}{(1 - e^{-\lambda^{3/4}})^2 + 4e^{-\lambda^{3/4}} \sin^2(\pi s)},$$

and we have

$$|\hat{F}^*(s)| \leq \frac{C\sqrt{\lambda}}{\lambda^{3/2} + \sin^2(\pi s)}$$

as soon as λ is sufficiently small. Moreover, a simple computation shows that

$$\begin{aligned} \int_{[0,1]^2} dsdt \frac{|\hat{G}_1^*(s,t)|^2}{\lambda + 4\theta(s) + 4\theta(t)} &= \int_{[0,1]^2} dsdt \frac{(\sin(2\pi s) + \sin(2\pi t))^2 |\hat{F}^*(s+t)|^2}{\lambda + 4\theta(s) + 4\theta(t)} \\ &= \int_0^1 ds |\hat{F}^*(s)|^2 \sin^2(\pi s) U_\lambda(s), \end{aligned} \tag{22}$$

where

$$U_\lambda(s) = \int_0^1 \frac{\cos^2(\pi u)}{\lambda + 2 \sin^2\left(\frac{\pi(u+s)}{2}\right) + 2 \sin^2\left(\frac{\pi(u-s)}{2}\right)} du.$$

We are interested in the behavior of (22) as λ goes to zero. The critical points of the integrand are 0 and 1, and by symmetry arguments, we can restrict ourselves to 0. By Eq. (4.12) of Ref. 5, one has

$$U_\lambda(s) \leq \frac{C}{\sqrt{\lambda + s^2}}.$$

It follows that

$$4 \int_0^{1/2} ds |\hat{F}^*(s)|^2 \sin^2(\pi s) U_\lambda(s) \leq C\lambda \int_0^{1/2} ds \frac{\sin^2(\pi s)}{(\lambda^{3/2} + \sin^2(\pi s))^2 \sqrt{\lambda + s^2}}.$$

Standard analysis shows that this last term is of order $\lambda^{-1/4}$ which proves the first claim.

Let us examine the term G_8 (terms G_5, G_6, G_7 are evaluated in the same way). We have

$$\hat{G}_8^*(x, y) = \frac{\lambda^{-1/4}(1 - e^{-\lambda^{3/4}})}{4} \delta_1(y) \delta_{x \geq 3} e^{-\lambda^{3/4}x}.$$

By a direct computation, we obtain

$$|\hat{G}_8^*(s, t)|^2 \leq C \frac{\lambda}{\lambda^{3/2} + \sin^2(\pi s)}.$$

It follows that

$$\begin{aligned} \langle\langle G_8, (\lambda - \Delta)^{-1} G_8 \rangle\rangle_0 &= \int_{[0,1]^2} ds dt \frac{|\hat{G}_8^*(s, t)|^2}{\lambda + 4\theta(s) + 4\theta(t) + \theta(s+t)} \\ &\leq C\lambda \int_0^1 ds \frac{1}{\lambda^{3/2} + \sin^2(\pi s)} \left(\int_0^1 \frac{dt}{\lambda + \theta(s) + \theta(t)} \right) \leq C\lambda \int_0^1 \frac{ds}{(\lambda^{3/2} + s^2)(\sqrt{\lambda + s^2})} \\ &= O(\lambda^{-1/4}) \end{aligned}$$

because we have

$$\int_0^1 \frac{dt}{\lambda + \theta(s) + \theta(t)} \leq \frac{C}{\sqrt{\lambda + \theta(s)}}.$$

For the terms G_2, G_3, G_4 , we use the fact that $|\hat{G}_j^*(s, t)| \leq C\lambda^{1/2}$ if $j=2, 3, 4$, and we observe that

$$\int_{[0,1]^2} \frac{ds dt}{\lambda + \theta(s) + \theta(t)} = O(\log \lambda).$$

The lemma is proven. □

We can now conclude the proof of Theorem 1 for the one dimensional case. We use the variational formula (15) and the test function

$$f(p) = a \sum_{x,y} F(x, y) H_{2\delta_x + 2\delta_y}(p).$$

Here a is a positive constant we will fix later and the function F is defined in (19). By (20), Lemma 4, and Eq. (15), we obtain

$$\langle\langle w_1, (\lambda - L)^{-1} w_1 \rangle\rangle \geq C_1 a \lambda^{-1/4} - C_2 a^2 \lambda^{1/4} - C_3 a^2 \lambda^{-1/4},$$

where C_1, C_2, C_3 are positive constants independent of a and λ . If a is chosen sufficiently small, the lower bound is of order $\lambda^{-1/4}$ and Theorem 1 is proven.

VI. THE TWO DIMENSIONAL CASE

For simplicity, we assume $a_2=0$ and $a_1=1$. In this case, one can, in fact, obtain better lower bounds [see (24)]. The general case $a_2^2+a_1^2>0$ can be treated in a similar way but Fourier's analysis is slightly more difficult and lower bounds are of order $\log|\log \lambda|$ instead of $(|\log \lambda|)^{1/2}$ (see Ref. 23 Proof of Proposition 1.3, for a detailed analysis).

Lemma 5: Fix $R>0$ and assume $d \geq 2$. There exists a positive constant $C=C(d)$ independent

of λ and n such that for any local symmetric function $F: (\mathbb{Z}^d)^n \rightarrow \mathbb{R}$,

$$\sum_{i,j,k,l=1}^n \langle \mathbf{1}_{|x_i-x_j|+|x_k-x_l| \leq R} F^2 \rangle_0 \leq Cn^3 \langle F, (\lambda - \Delta)F \rangle_0.$$

Proof: This lemma is proven in Lemma 4.2 of Ref. 16. □

Lemma 6: Let $d \geq 2$ and $n \geq 1$. There exists a constant $C(n)$ such that for any local symmetric functions $F: (\mathbb{Z}^d)^{n+1} \rightarrow \mathbb{R}$ and $G: (\mathbb{Z}^d)^n \rightarrow \mathbb{R}$,

$$|\langle G, \mathcal{A}^\pm F \rangle - \langle G, \mathcal{A}^\pm F \rangle_0| \leq C(n) \langle F, (\lambda - \Delta)F \rangle_0^{1/2} \langle G, (\lambda - \Delta)G \rangle_0^{1/2},$$

and it follows that

$$|\langle \langle G, \mathcal{A}^\pm F \rangle \rangle - \langle \langle G, \mathcal{A}^\pm F \rangle \rangle_0| \leq C(n) \langle \langle F, (\lambda - \Delta)F \rangle \rangle^{1/2} \langle \langle G, (\lambda - \Delta)G \rangle \rangle_0^{1/2}.$$

Proof: One has

$$(\mathcal{A}_i^+ F)(\mathbf{x}) = \sum_{k,\ell=1}^n \mathbf{1}_{\{x_k=x_\ell+e_i\}} (F(\mathbf{x}^k) - F(\mathbf{x}^\ell)),$$

with \mathbf{x}^k the vector \mathbf{x} where the coordinate x^k has been removed. Observe it is the operator defined in formula (3.4) of Ref. 16. We have

$$\langle G, \mathcal{A}_i^+ F \rangle - \langle G, \mathcal{A}_i^+ F \rangle_0 = \sum_{k,\ell=1}^n \mathbf{1}_{\{x_k=x_\ell+e_i\}} G(\mathbf{x})(m(\mathbf{x}) - 1)(F(\mathbf{x}^k) - F(\mathbf{x}^\ell)). \tag{23}$$

If $\mathbf{x} \in (\mathbb{Z}^d)^n$ is such that $x_i \neq x_j$ for every $i \neq j$, then $m(\mathbf{x}) - 1 = 0$. Hence we can introduce the sum of the following indicator function:

$$\sum_{i \neq j} \mathbf{1}_{x_i=x_j},$$

in the sum (23). We recall that $|m(\mathbf{x}) - 1| \leq C(n)$ for a constant $C(n)$. It follows that

$$|\langle G, \mathcal{A}_i^+ F \rangle - \langle G, \mathcal{A}_i^+ F \rangle_0| \leq C(n) \sum_{i,j,k,\ell} \sum_{\mathbf{x}} |G(\mathbf{x}) \mathbf{1}_{|x_k-x_\ell| \leq 1} \mathbf{1}_{|x_i-x_j| \leq 1}| |F(\mathbf{x}^k) - F(\mathbf{x}^\ell)|.$$

One concludes by Schwarz inequality. □

Corollary 2: Let $d \geq 2$ and $F: (\mathbb{Z}^d)^n \rightarrow \mathbb{R}$. There exists a constant $C(n)$ such that

$$\|\mathcal{A}^\pm F\|_{-1,\lambda}^2 \leq C(n) (\|\mathcal{A}^\pm F\|_{-1,\lambda,0}^2 + \|F\|_{1,\lambda,0}^2).$$

Proof: We have only to prove the lemma with the inner product $\langle \langle \cdot, \cdot \rangle \rangle$ replaced with the standard inner product without translations $\langle \cdot, \cdot \rangle$ thanks to (13) and (14). Recall that

$$\langle \mathcal{A}^+ F, (\lambda - S)^{-1} \mathcal{A}^+ F \rangle = \sup_G \{2 \langle \mathcal{A}^+ F, G \rangle - \lambda \langle G, G \rangle - \langle G, (-S)G \rangle\}.$$

Then we use Lemmas 6 and 1 and Eq. (12) to conclude. □

We can now complete the proof. Recall that we want obtain a lower bound for the right hand side of (15) and remark that in the dual basis, one has

$$\langle \langle w_1, f \rangle \rangle = 2a_1 \sum_{z \in \mathbb{Z}^d} F(2\delta_z + 2\delta_{z+e_1}),$$

where F is the function corresponding to f in the dual basis. We restrict the supremum over functions belonging to $\mathcal{H}_{0,4}$. Hence, by Corollaries 1 and 2 and Lemma 2, one has

$$\langle\langle w_1, (\lambda - L)^{-1} w_1 \rangle\rangle \geq \sup_{F \in \mathcal{H}_{0,4}} \left\{ 4a_1 \sum_z F(2z, 2z + 2e_1) - C \|F\|_{1,\lambda,0}^2 - C \|A^+ F\|_{-1,\lambda,0}^2 \right\}.$$

The maximizer for this new variational problem can be computed using Fourier transform (cf. Ref. 16, Lemma 3.3, and Ref. 23, Proof of Proposition 1.3). It turns out that

$$\langle\langle w_1, (\lambda - L)^{-1} w_1 \rangle\rangle \geq \begin{cases} C |\log \lambda|^{1/2} & d = 2 \\ C & d \geq 3. \end{cases} \tag{24}$$

For $d=2$ and $a_2 \neq 0$, one can only show (see Ref. 23)

$$\langle\langle w_1, (\lambda - L)^{-1} w_1 \rangle\rangle \geq C \log(|\log \lambda|).$$

VII. THE d -DIMENSIONAL CASE FOR $d \geq 3$

In this section, we show that the diffusion coefficient is finite and strictly positive if the dimension d is greater than 3. For the ASEP, this has been proved in Ref. 17. The lower bound follows from the previous section. The upper bound is obtained by ignoring the asymmetric part of the generator,

$$\begin{aligned} \langle\langle w_1, (\lambda - L)^{-1} w_1 \rangle\rangle &= \sup_f \{ 2\langle\langle f, w_1 \rangle\rangle - \langle\langle f, (\lambda - S)f \rangle\rangle - \langle\langle Af, (\lambda - S)^{-1} Af \rangle\rangle \} \\ &\leq \sup_f \{ 2\langle\langle f, w_1 \rangle\rangle - \langle\langle f, (\lambda - S)f \rangle\rangle \}. \end{aligned} \tag{25}$$

We write this last variational formula in the dual basis H_ξ and we recall that

$$\langle\langle w_1, f \rangle\rangle = 2a_1 \sum_{z \in \mathbb{Z}^d} F(2\delta_z + 2\delta_{z+e_1}),$$

with $f(p) = \sum_\xi F(\xi) H_\xi(p)$. Let us decompose F in the orthogonal sum composed of subspaces $\mathcal{H}_{n,\varepsilon}$ (ε are elements of $\{0, 1\}^{\mathbb{Z}^d}$),

$$F = \sum_{n,\varepsilon} F_{n,\varepsilon}, \quad F_{n,\varepsilon} \in \mathcal{H}_{n,\varepsilon}.$$

Since S sends an element of $\mathcal{H}_{n,\varepsilon}$ on an element of $\mathcal{H}_{n,\varepsilon}$, we have

$$\|F\|_{1,\lambda}^2 = \sum_{n,\varepsilon} \|F_{n,\varepsilon}\|_{1,\lambda}^2.$$

It follows that in the third line of (25), one can restrict the supremum over functions F belonging to $\mathcal{H}_{2,0}$. If F belongs to $\mathcal{H}_{2,0}$ then F is identified with a symmetric function on $(\mathbb{Z}^d)^2$, and we have

$$\|F\|_{1,\lambda}^2 \geq C \|F\|_{1,\lambda,0}^2$$

for a positive constant C . The supremum is then easily computed using Fourier transform. In the supremum appears the Green function of the discrete Laplacian which is finite only for dimension $d \geq 3$. We have

$$\langle\langle w_1, (\lambda - L)^{-1} w_1 \rangle\rangle \leq C, \quad d \geq 3.$$

VIII. CONCLUSIONS AND FINAL REMARKS

For $d=2$, the diffusion coefficient is expected to be of order $(\log t)^{2/3}$. It has been proven by Yau in Ref. 26 in a very technical paper for ASEP. It is not clear that the method of Ref. 26 can be applied for AEM. Indeed, constants $C(n)$ appearing in Lemma 1 are exponential in n and the method of Ref. 26 seems to be restricted to polynomial dependence in n . Remark that ASEP is the

only model belonging to KPZ class for which such behavior is proven for $d=2$. For the case $d \geq 3$, it would be interesting using generalized duality techniques to establish a fluctuation-dissipation equation¹⁷ for AEM, meaning a decomposition of the current in the form

$$w_i = \nabla \varphi_i + L h_i \quad (26)$$

for functions φ_i and h_i in a suitable Hilbert space.

In this paper we obtained lower (and upper for $d \geq 3$) bounds for the diffusion coefficient for AEM. The strategy was based on generalized duality techniques similar to Refs. 5 and 16. A recent paper of Balázs and Seppäläinen (cf., Ref. 2) improves considerably the lower bounds obtained in Ref. 16 in the one dimensional case for the diffusion coefficient. For nearest neighbor ASEP (but not for general ASEP), the authors of Ref. 2 are able to prove upper and lower bounds with the right order. Nevertheless, their method has restrictions: a key ingredient is attractivity of the process (AEM is not) and the method has only been developed for the one dimensional case. Of course, generalized duality techniques have also restrictions (see the Introduction).

A model introduced by Seppäläinen in Ref. 22 belongs to the KPZ universality class and should have anomalous behavior in low dimension as ASEP and AEM. Even if one consider this attractive process, methods of Ref. 2 are difficult to apply (there is no concept of second class particle for this process). This process is also difficult to study with duality techniques because the symmetric part of the generator does not conserve the degree. But a suitable modification of the process can be studied with duality techniques. The slight modification is the discrete counterpart of the AEOM. The symmetric part is given by the KMP process¹³ and the asymmetric part by the asymmetric part of the process defined in Ref. 22. The state space is $\mathbb{R}^{\mathbb{Z}^d}$. For a real valued local function $f(\xi)$ defined on the state space of the process, the action of the generator L on f is given by

$$(Lf)(\xi) = \sum_{i=1}^d \sum_{x \in \mathbb{Z}^d} \int_0^1 dp [f(E_{x,x+e_i,p}\xi) - f(\xi)] + a_i \xi_x [f(T_{x,x+e_i,p}\xi) - f(T_{x,x-e_i,p}\xi)], \quad (27)$$

where the exchange operator E and the transfer operator T are defined by

$$E_{x,y,p}\xi = \xi + [p(\xi_x + \xi_y) - \xi_x] \delta_x + [(1-p)(\xi_x + \xi_y) - \xi_y] \delta_y,$$

$$T_{x,y,p}\xi = \xi + p \xi_x (\delta_y - \delta_x).$$

The existence of this Markov process is also a difficult problem like for AEM. We conjecture that Assumption 1 is also valid here. Then the process is well defined, conserves the energy $\sum_x \xi_x$, and centered exponential product measures of the form

$$\mu_\lambda(d\xi) = \prod_x \lambda e^{-\lambda \xi_x} d\xi_x$$

are invariant for the dynamics. The symmetric part of the generator is the KMP process. At the difference of the symmetric part of the process defined in Ref. 22, it conserves the degree and generalized duality technique presented here (the basis is composed of multivariate Laguerre polynomials) can be applied to the process defined by (27), and one obtains similar lower bounds for the corresponding diffusion coefficient.

ACKNOWLEDGMENTS

The author thanks C. Landim, F. Redig, and S. Sethuraman for valuable discussions and J. Fritz for useful comments on the existence problems of the dynamics. He thanks also the anonymous referee for pertinent comments. The author acknowledges the support of the French Ministry of Education through Grant No. ANR BLAN07-2184264.

APPENDIX

We derive here the expression of the generator L in the dual basis H_ξ . Recall that the Hermite polynomials $(h_n)_n$ satisfy the following equations:

$$\frac{d^2 h_n}{du^2} - u \frac{dh_n}{du} + nh_n = 0,$$

$$\frac{dh_n}{du} = nh_{n-1},$$

$$h_{n+1}(u) = uh_n(u) - nh_{n-1}(u),$$

$$h_0 = 1,$$

(A1)

where we adopt the convention that $h_n=0$ if $n<0$. We denote by \tilde{H}_ξ the multivariate Hermite polynomial without the normalization factor $n(\xi)$,

$$\tilde{H}_\xi = n(\xi)H_\xi.$$

Fix $x \neq y$ in \mathbb{Z}^d . By (A1), we have

$$(p_y \partial_{p_x} - p_x \partial_{p_y}) \tilde{H}_\xi = \xi_x \tilde{H}_{\xi^{x,-1,y,+1}} - \xi_y \tilde{H}_{\xi^{y,-1,x,+1}}.$$

It follows that

$$(p_y \partial_{p_x} - p_x \partial_{p_y})^2 \tilde{H}_\xi = \xi_x(\xi_x - 1) \tilde{H}_{\xi^{x,-2,y,+2}} - \xi_x(\xi_y + 1) \tilde{H}_\xi + \xi_y(\xi_y - 1) \tilde{H}_{\xi^{x,+2,y,-2}} - \xi_y(\xi_x + 1) \tilde{H}_\xi,$$

and we get finally

$$\begin{aligned} (p_y \partial_{p_x} - p_x \partial_{p_y})^2 H_\xi &= (\xi_x - 1)(\xi_y + 2) H_{\xi^{x,-2,y,+2}} - \xi_x(\xi_y + 1) H_\xi \\ &\quad + (\xi_x + 2)(\xi_y - 1) H_{\xi^{x,+2,y,-2}} - \xi_y(\xi_x + 1) H_\xi \end{aligned}$$

because we have $n(k-2)/n(k)=1/k$. To obtain the expression for $(S_i F)(\xi)$, we write $f = \sum_\xi F(\xi) H_\xi$, and we have

$$\begin{aligned} S_i f &= \sum_{\xi \in \mathbb{N}^d, x \in \mathbb{Z}^d} F(\xi) \{ (\xi_x - 1)(\xi_{x+e_i} + 2) H_{\xi^{x,-2,x+e_i,+2}} - \xi_x(\xi_{x+e_i} + 1) H_\xi \} \\ &\quad + \sum_{\xi \in \mathbb{N}^d, x \in \mathbb{Z}^d} F(\xi) \{ (\xi_x + 2)(\xi_{x+e_i} - 1) H_{\xi^{x,+2,x+e_i,-2}} - \xi_{x+e_i}(\xi_x + 1) H_\xi \}. \end{aligned}$$

Recall here that our conventions are such that $H_\xi=0$ if there exists $z \in \mathbb{Z}^d$ such that $\xi_z < 0$. Moreover, any function $F: \mathbb{N}^d \rightarrow \mathbb{R}$ is extended to a function F on $\mathbb{N}^d \cup \{o\}$ by $F(o)=0$ with the convention that $\xi=o$ if there exists $z \in \mathbb{Z}^d$ such that $\xi_z < 0$. By a suitable obvious change in variables, we get

$$\begin{aligned} S_i f &= \sum_{\xi \in \mathbb{N}^d, x \in \mathbb{Z}^d} \left\{ \sum_x (\xi_x + 1) \xi_{x-e_i} [F(\xi^{x,+2,x-e_i,-2}) - F(\xi)] \right\} H_\xi \\ &\quad + \sum_{\xi \in \mathbb{N}^d, x \in \mathbb{Z}^d} \left\{ \sum_x (\xi_x + 1) \xi_{x-e_i} [F(\xi^{x,-2,x+e_i,+2}) - F(\xi)] \right\} H_\xi. \end{aligned}$$

The computations are similar for the asymmetric part. We have

$$p_x p_y (p_y \partial_{p_x} - p_x \partial_{p_y}) \tilde{H}_\xi = p_x p_y (\xi_x \tilde{H}_{\xi_x-1, y, +1} - \xi_y \tilde{H}_{\xi_y-1, x, +1}).$$

By symmetry we can restrict ourselves to compute

$$p_x p_y (\xi_x \tilde{H}_{\xi_x-1, y, +1}),$$

which is equal to

$$\xi_x \tilde{H}_{\xi_y+2} + \xi_x (\xi_y + 1) (\xi_x - 1) \tilde{H}_{\xi_x-2} + \xi_x (\xi_x - 1) \tilde{H}_{\xi_x-2, y, +2} + \xi_x (\xi_y + 1) \tilde{H}_\xi,$$

and we obtain that

$$p_x p_y (p_y \partial_{p_x} - p_x \partial_{p_y}) \tilde{H}_\xi$$

is equal to

$$\begin{aligned} & \xi_x \tilde{H}_{\xi_y+2} + \xi_x (\xi_y + 1) (\xi_x - 1) \tilde{H}_{\xi_x-2} - \xi_y \tilde{H}_{\xi_x+2} - \xi_y (\xi_x + 1) (\xi_y - 1) \tilde{H}_{\xi_y-2} \\ & + (\xi_x - \xi_y) \tilde{H}_\xi + \xi_x (\xi_x - 1) \tilde{H}_{\xi_x-2, y, +2} - \xi_y (\xi_y - 1) \tilde{H}_{\xi_y-2, x, +2}. \end{aligned}$$

Since $\tilde{H}_\xi = n(\xi) H_\xi$, we obtain

$$\begin{aligned} p_x p_y (p_y \partial_{p_x} - p_x \partial_{p_y}) \tilde{H}_\xi &= \xi_x (\xi_y + 2) H_{\xi_y+2} - \xi_y (\xi_x + 2) H_{\xi_x+2} + (\xi_y + 1) (\xi_x - 1) H_{\xi_x-2} \\ & - (\xi_y - 1) (\xi_x + 1) H_{\xi_y-2} + (\xi_y - \xi_x) H_\xi + (\xi_x - 1) (\xi_y + 2) H_{\xi_x-2, y, +2} \\ & - (\xi_y - 1) (\xi_x + 2) H_{\xi_x+2, y, -2}. \end{aligned}$$

If f is a local smooth function such that $f(p) = \sum_{\xi} F(\xi) H_\xi$, then

$$\begin{aligned} \mathcal{A}_i f &= \sum_{\xi, x} F(\xi) \{ \xi_x (\xi_{x+e_i} + 2) H_{\xi_x+e_i+2} - \xi_{x+e_i} (\xi_x + 2) H_{\xi_x+2} \} \\ & + \sum_{\xi, x} F(\xi) \{ (\xi_{x+e_i} + 1) (\xi_x - 1) H_{\xi_x-2} - (\xi_{x+e_i} - 1) (\xi_x + 1) H_{\xi_x+e_i-2} \} \\ & + \sum_{\xi, x} F(\xi) \{ (\xi_x - 1) (\xi_{x+e_i} + 2) H_{\xi_x-2, x+e_i+2} - (\xi_{x+e_i} - 1) (\xi_x + 2) H_{\xi_x+2, x+e_i-2} \}. \end{aligned}$$

By suitable changes in variables, we get the announced expression for $\mathcal{A}_i F$.

¹Balázs, M. and Komjáthy, J., e-print arXiv:0804.1397v1.

²Balázs, M. and Seppäläinen, T., e-print arXiv:math/0608400v2.

³Basile, G., Bernardin, C., and Olla, S., "Thermal conductivity for a momentum conserving model," Commun. Math. Phys. (to be published); e-print arXiv:cond-mat/0601544.

⁴van Beijeren, H., Kutner, R., and Spohn, H., "Excess noise for driven diffusive systems," Phys. Rev. Lett. **54**, 2026 (1985).

⁵Bernardin, C., "Fluctuations in the occupation time of a site in the asymmetric simple exclusion process," Ann. Probab. **32**, 855 (2004).

⁶Ferrari, P. L., e-print arXiv:math-ph/0701021v1.

⁷Ferrari, P. L. and Prähofer, M., "One-dimensional stochastic growth and Gaussian ensembles of random matrices," Markov Processes Relat. Fields **12**, 203 (2006).

⁸Ferrari, P. L. and Spohn, H., "Scaling limit for the space-time covariance of the stationary totally asymmetric simple exclusion process," Commun. Math. Phys. **265**, 1 (2006).

⁹Fritz, J. private communication.

¹⁰Fritz, J., "Infinite lattice systems of interacting diffusion processes, existence and regularity properties," Z. Wahrscheinlichkeitstheor. Verwandte Geb. **59**, 291 (1982).

¹¹Giardinà, C., Kurchan, J., and Redig, F., "Duality and exact correlations for a model of heat conduction," J. Math. Phys. **48**, 033301 (2007).

¹²Kardar, M., Parisi, G., and Zhang, Y. C., "Dynamical scaling of growing interfaces," Phys. Rev. Lett. **56**, 889 (1986).

- ¹³Kipnis, C., Marchioro, C., and Presutti, E., "Heat flow in an exactly solvable model," *J. Stat. Phys.* **27**, 65 (1982).
- ¹⁴Krug, J., Meakin, P., and Halpin-Healy, T., "Amplitude universality for driven interfaces and directed polymers in random media," *Phys. Rev. A* **45**, 638 (1992).
- ¹⁵Landim, C., Olla, S., and Yau, H. T., "Properties of the diffusion coefficients for the Asymmetric Simple Exclusion Processes," *Ann. Probab.* **24**, 1779 (1997).
- ¹⁶Landim, C., Quastel, C. J., Salmhofer, M., and Yau, H. T., "Superdiffusivity of asymmetric exclusion process in dimensions one and two," *Commun. Math. Phys.* **244**, 455 (2004).
- ¹⁷Landim, C. and Yau, H. T., "Fluctuation-dissipation equation of asymmetric simple exclusion processes," *Probab. Theory Relat. Fields* **108**, 321 (1997).
- ¹⁸Olla, S. and Tremoulet, C., "Equilibrium fluctuations for interacting Ornstein-Uhlenbeck particles," *Commun. Math. Phys.* **233**, 463 (2003).
- ¹⁹Prähofer, M. and Spohn, H., in *In and Out of Equilibrium (Mambucaba, 2000)*, Progress in Probability Vol. 51 (Birkhauser, Boston, MA, 2002), pp. 185–204.
- ²⁰Prähofer, M. and Spohn, H., "Exact scaling functions for one-dimensional stationary KPZ growth," *J. Stat. Phys.* **115**, 255 (2004).
- ²¹Quastel, J. and Valko, B., " $t^{1/3}$ Superdiffusivity of finite-range asymmetric exclusion processes on Z ," *Commun. Math. Phys.* **273**, 379 (2007).
- ²²Seppäläinen, T., "A microscopic model for the Burgers equation and longest increasing subsequences," *Electron. J. Probab.* **1**, 51 (1996).
- ²³Sethuraman, S., "Superdiffusivity of occupation-time variance in 2-dimensional asymmetric exclusion processes with density $\rho=1/2$," *J. Stat. Phys.* **123**, 787 (2006).
- ²⁴Sethuraman, S., "Diffusive variance for a tagged particle in $d \leq 2$ asymmetric simple exclusion," *ALEA Lat. Am. J. Probab. Math. Stat.* **1**, 305 (2006).
- ²⁵Spohn, H., *Large Scale Dynamics of Interacting Particles* (Springer-Verlag, New-York, 1991).
- ²⁶H. T. Yau, " $(\log t)^{2/3}$ law of the two dimensional asymmetric simple exclusion process," *Ann. Math.* **159**, 377 (2002).