

SYMMETRIC SPACES AND THE KASHIWARA-VERGNE METHOD

Reykjavik, August 15, 2007
François Rouvière
(Université de Nice, France)

til Sigurðar Helgasonar með þökkum, vinsemd og virðingu

Introduction

Let G/K denote a Riemannian symmetric space of the non-compact type.

"The action of the G -invariant differential operators on G/K on the radial functions on G/K is isomorphic with the action of certain differential operators with constant coefficients. The isomorphism in question is used in Harish-Chandra's work on the Fourier analysis on G and is related to the Radon transform on G/K .

For the case when G is complex a more direct isomorphism of this type is given, again as a consequence of results of Harish-Chandra."

This is a quote from S. Helgason [8] *Fundamental solutions of invariant differential operators on symmetric spaces* (1964), one of the first mathematical papers I studied. Those two results were fascinating to me, they still are and they motivated my everlasting interest in invariant differential operators as well as Radon transforms. Yet I was dreaming simpler proofs could be given, without relying on Harish-Chandra's deep study of semisimple Lie groups...

Then, in the fall of 1977, came a preprint by Kashiwara and Vergne [11] *The Campbell-Hausdorff formula and invariant hyperfunctions*, showing that similar results could be obtained - for solvable Lie groups at least - only by means of "elementary" (but very clever) computations with the exponential mapping and the Campbell-Hausdorff formula.

After briefly recalling this method for Lie groups I will give an overview of its extension to general symmetric spaces, including several recent results.

1. The Kashiwara-Vergne method for Lie groups

Let G be a Lie group with Lie algebra \mathfrak{g} . The Campbell-Hausdorff formula

$$\log(\exp X \exp Y) = X + Y + \frac{1}{2}[X, Y] + \frac{1}{12}[X, [X, Y]] - \frac{1}{12}[Y, [X, Y]] + \dots \quad (1)$$

expresses (locally) the group law of G in the exponential chart $\exp : \mathfrak{g} \rightarrow G$ as a convergent series of iterated Lie brackets. Kashiwara and Vergne [11] stated the following

Conjecture 1 *The formula can be written as*

$$Z := \log(\exp Y \exp X) = X + Y + (e^{-\text{ad} X} - 1)F(X, Y) + (1 - e^{\text{ad} Y})G(X, Y) \quad (2)$$

where $F, G : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ are given by convergent power series near $(0, 0)$ and satisfy the trace condition

$$\text{tr}_{\mathfrak{g}}(\text{ad} X \cdot \partial_X F + \text{ad} Y \cdot \partial_Y G) = \frac{1}{2} \text{tr}_{\mathfrak{g}} \left(\frac{\text{ad} X}{e^{\text{ad} X} - 1} + \frac{\text{ad} Y}{e^{\text{ad} Y} - 1} - \frac{\text{ad} Z}{e^{\text{ad} Z} - 1} - 1 \right) . \quad (3)$$

Formula (2) is rather easily obtained: roughly speaking one can group together all brackets $[X, \dots]$ resp. $[Y, \dots]$ in the right-hand side of (1), though not in a unique way. But (3) is far from obvious!

Let $j(X) = \det_{\mathfrak{g}} \left(\frac{1 - e^{-\text{ad} X}}{\text{ad} X} \right)$, the jacobian of \exp . Working on the domain of the chart \exp , we associate to a function f on \mathfrak{g} the function \tilde{f} on G such that

$$f(X) = j(X)^{1/2} \tilde{f}(\exp X) .$$

This correspondence extends to distributions u (on \mathfrak{g}) and \tilde{u} (on G) with

$$\langle \tilde{u}, \tilde{f} \rangle = \langle u, f \rangle$$

for any test function f on \mathfrak{g} .

A distribution u on \mathfrak{g} is said to be *invariant* if $\langle u(X), f(\text{Ad} g(X)) \rangle = \langle u(X), f(X) \rangle$ for all $g \in G$, $f \in \mathcal{D}(\mathfrak{g})$.

Theorem 2 (*Kashiwara-Vergne*) *If the conjecture is true, then*

$$\tilde{u} *_G \tilde{v} = \widetilde{u *_G v}$$

for any invariant distributions u, v such that the convolutions make sense.

Taking v supported at the origin it follows that bi-invariant differential operators on G correspond to constant coefficients differential operators on the Lie algebra by means of \sim .

Theorem 3 (*Kashiwara-Vergne*) *The conjecture is true for solvable Lie algebras.*

The conjecture is also true for $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{R})$ (R. 1981 [14], a lengthy and boring computation). What about general Lie algebras?

What can you do when you fail to prove a difficult conjecture? Either you give up or you extend the conjecture. I did both: I extended it to symmetric spaces, then I gave up. Decisive progress was made in the last decade fortunately:

- true for any quadratic Lie algebra (Vergne 1999 [20], Alekseev and Meinrenken 2002 [1])

- true for all Lie algebras (Andler, Sahi and Torossian 2001 [4], Torossian 2002 [18], Alekseev and Meinrenken 2006 [2]). See the June 2007 Bourbaki seminar [19] by Torossian for a nice survey. Those works rely on M. Kontsevich's fundamental paper [12] on quantization.

Let us also mention two interesting recent works by Alekseev and Petracchi (2006) [3] and by Burgunder (2006) [5], discussing the uniqueness of functions F, G in the conjecture. The former paper proves the uniqueness, up to order one in Y , of F and G satisfying (2) and (3) and the natural symmetry $G(X, Y) = F(-Y, -X)$. The latter gives all solutions F, G of (2) alone.

2. e-functions of symmetric spaces

2.a. Definition

Let G/H denote a general symmetric space and $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{s}$ the decomposition of the Lie algebra given by the symmetry. Here \mathfrak{h} is the Lie algebra of H and \mathfrak{s} identifies to the tangent space to G/H at the origin. Let $\exp : \mathfrak{g} \rightarrow G$ and $\text{Exp} : \mathfrak{s} \rightarrow G/H$ be the respective exponential mappings. *We shall always work on neighborhoods of the origin where they are diffeomorphisms.* To simplify the exposition this will not be repeated in the sequel.

Let

$$J(X) = \det_{\mathfrak{s}} \left(\frac{\text{sh ad } X}{\text{ad } X} \right), \quad X \in \mathfrak{s},$$

be the Jacobian of Exp . As before we relate a function f , resp. a distribution u , on \mathfrak{s} to \tilde{f} , resp. \tilde{u} , on G/H by

$$f(X) = J(X)^{1/2} \tilde{f}(\text{Exp } X), \quad \langle \tilde{u}, \tilde{f} \rangle = \langle u, f \rangle.$$

The distribution u on \mathfrak{s} is said to be *H-invariant* if $\langle u(X), f(\text{Ad } h(X)) \rangle = \langle u(X), f(X) \rangle$ for all $h \in H$, $f \in \mathcal{D}(\mathfrak{s})$. The nice convolution formula of the group case no longer holds for general symmetric spaces. One must consider "twisted convolutions" instead and give the following

Definition 4 *An analytic function $e : \mathfrak{s} \times \mathfrak{s} \rightarrow \mathbb{R}$ is called an **e-function** of G/H if $e(h \cdot X, h \cdot Y) = e(X, Y)$ for all $h \in H$, $X, Y \in \mathfrak{s}$ (adjoint action) and, for all H -invariant distributions u, v and all test functions f on \mathfrak{s} ,*

$$\langle \tilde{u} *_{G/H} \tilde{v}, \tilde{f} \rangle = \langle u(X) \otimes v(Y), e(X, Y) f(X + Y) \rangle. \quad (4)$$

*The symmetric space is said to be **special** if it admits an e-function which is identically 1.*

The convolution on the left is defined by

$$\langle \tilde{u} *_{G/H} \tilde{v}, \tilde{f} \rangle = \langle \tilde{u}(xH), \langle \tilde{v}(yH), \tilde{f}(xyH) \rangle \rangle$$

where x, y are variables in G (see [15] §1 for more details).

Example. Let G/K be a *rank one* Riemannian symmetric space of the non-compact type, $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ as usual, and let p, q be the respective multiplicities of the roots $\alpha, 2\alpha$. The dimension of G/K is $n = p + q + 1$. Then an e-function of G/K is given by (Flensted-Jensen & R., unpublished)

$$e(X, Y) = \left(4 \frac{x}{\text{sh } x} \cdot \frac{y}{\text{sh } y} \cdot \frac{z}{\text{sh } z} \cdot \frac{\text{ch}(x+y) - \text{ch } z}{(x+y)^2 - z^2} \cdot \frac{\text{ch } z - \text{ch}(x-y)}{z^2 - (x-y)^2} \right)^{(n-3)/2} \times \\ \times {}_2F_1 \left(1 - \frac{q}{2}, \frac{q}{2}; \frac{n-1}{2}; \frac{(\text{ch}(x+y) - \text{ch } z)(\text{ch } z - \text{ch}(x-y))}{4 \text{ch } x \text{ch } y \text{ch } z} \right) \quad (5)$$

where $X, Y \in \mathfrak{p}$, $x = \|X\| = (-B(X, \theta X)/2(p+4q))^{1/2}$ (normalized Killing form), $y = \|Y\|$, $z = \|X+Y\|$. The result comes out of manipulations of various integral

formulas. Note that the hypergeometric factor ${}_2F_1$ is 1 if $q = 0$ i.e. $G/K = H^n(\mathbb{R})$, and this space is special if $n = 3$.

2.b. Application to invariant differential operators

Let $p \in S(\mathfrak{s})^{\mathfrak{h}}$ be an \mathfrak{h} -invariant element in the symmetric algebra of \mathfrak{s} , identified to an H -invariant differential operator $p(\partial)$ on \mathfrak{s} with constant coefficients. The distribution $v = {}^t p \delta$ (where t means transpose and δ is the Dirac measure at the origin of \mathfrak{s}) is then H -invariant, supported at the origin. Applying (4) we obtain, for any H -invariant distribution u on \mathfrak{s} ,

$$\langle \tilde{u} * ({}^t p \delta)^\sim, \tilde{f} \rangle = \langle u(X) \otimes {}^t p \delta(Y), e(X, Y) f(X + Y) \rangle .$$

The left-hand side is $\langle {}^t \tilde{p} \tilde{u}, \tilde{f} \rangle$, where \tilde{p} is the G -invariant differential operator on G/H defined by

$$\tilde{p} \varphi(\text{Exp } X) = p(\partial_Y) \left(J(Y)^{1/2} \varphi(\exp X \cdot \text{Exp } Y) \right) \Big|_{Y=0} , \quad \varphi \in C^\infty(G/H) .$$

In the chart Exp , the operator \tilde{p} acting on H -invariant distributions is then expressed by

$${}^t \tilde{p} \tilde{u} = ({}^t p_e(X, \partial_X) u)^\sim \quad (6)$$

where $p_e(X, \partial_X)$ is the H -invariant differential operator on \mathfrak{s} defined by

$$p_e(X, \partial_X) f(X) = p(\partial_Y) (e(X, Y) f(X + Y)) \Big|_{Y=0} ,$$

with symbol

$$p_e(X, \xi) = \sum_{\alpha} \frac{1}{\alpha!} \partial_Y^\alpha e(X, 0) \cdot p^{(\alpha)}(\xi) , \quad X \in \mathfrak{s} , \quad \xi \in \mathfrak{s}^*$$

(finite sum).

Example 1. Let G/H be (pseudo-)Riemannian, with Laplace operator $L_{G/H}$, and let $p = L_{\mathfrak{s}}$ be the Laplacian of \mathfrak{s} . For any H -invariant function (or distribution) u on \mathfrak{s} formula (6) reduces to

$$L_{G/H} \tilde{u} = \left(L_{\mathfrak{s}} u - J^{-1/2} L_{\mathfrak{s}} J^{1/2} \cdot u \right)^\sim$$

(Helgason, 1972; see [9] p. 273). In this simple example only second order derivatives of e are needed, and they can be expressed by means of $J^{1/2}$.

Example 2. If G/H is special ($e = 1$) then $p_e(X, \partial_X) = p(\partial_X)$.

The map $p \mapsto \tilde{p}$ is thus a linear isomorphism of $S(\mathfrak{s})^{\mathfrak{h}}$ onto $\mathbb{D}(G/H)$, the algebra of G -invariant differential operators on the symmetric space, but not in general an isomorphism of algebras. Instead we have the following corollary of (6) :

Theorem 5 (*R. 1991 [16]*) *Let $S(\mathfrak{s})^{\mathfrak{h}}$ be equipped with the product \times defined by*

$$(p \times q)(\xi) = e(\partial_\xi, \partial_\eta) p(\xi) q(\eta) \Big|_{\xi=\eta} = \sum_{\alpha, \beta} \frac{1}{\alpha! \beta!} \partial_X^\alpha \partial_Y^\beta e(0, 0) p^{(\alpha)}(\xi) q^{(\beta)}(\xi) .$$

Then $p \mapsto \tilde{p}$ is an isomorphism of algebras of $(S(\mathfrak{s})^{\mathfrak{h}}, \times)$ onto $(\mathbb{D}(G/H), \circ)$, i.e. $\widetilde{p \times q} = \tilde{p} \circ \tilde{q}$.

Of course \times is the usual product in the symmetric algebra if G/H is special ($e = 1$).

2.c. A general construction of e-functions

The analogue of $\log(\exp X \exp Y)$ above is now $Z(X, Y) \in \mathfrak{s}$ defined by

$$\text{Exp } Z(X, Y) = \exp X \cdot \text{Exp } Y, \quad X, Y \in \mathfrak{s}.$$

Thus Z expresses the natural action of G on G/H read in exponential coordinates. It is readily checked by means of the symmetry that $2Z(X, Y) = \log(\exp X \exp 2Y \exp X)$; thus Z is given by a variant of the classical Campbell-Hausdorff formula. Working on this expression the following can be shown.

Theorem 6 (R. 1986 [15]) *There exist two analytic maps $a, b : \mathfrak{s} \times \mathfrak{s} \rightarrow H$ such that*

$$(X, Y) \longmapsto \Phi(X, Y) = (a(X, Y) \cdot X, b(X, Y) \cdot Y)$$

(adjoint action of H on \mathfrak{s}) is a diffeomorphism of $\mathfrak{s} \times \mathfrak{s}$ onto itself such that

$$Z \circ \Phi(X, Y) = X + Y.$$

Thus Φ transforms Z into its analogue for the flat symmetric space \mathfrak{s} . It is obtained as the value for $t = 1$ of a one-parameter family of diffeomorphisms Φ_t solving differential equations related to the evolution for $0 \leq t \leq 1$ of $Z_t(X, Y) = t^{-1}Z(tX, tY)$, $Z_0(X, Y) = X + Y$. In others words one works with *contractions of the symmetric space G/H* into its tangent space \mathfrak{s} .

For the sake of simplicity let us now *assume G/H admits a G -invariant measure*, a non-essential assumption so as to avoid some modular factors in the formulas.

Changing variables by means of the diffeomorphism Φ leads to

Theorem 7 (R. 1990 [16]) *The formula*

$$e(X, Y) = \left(\frac{J(X)J(Y)}{J(X+Y)} \right)^{1/2} \det_{\mathfrak{s} \times \mathfrak{s}} D\Phi(X, Y) \quad (7)$$

gives an e-function of G/H , strictly positive and such that

$$e(X, Y) = e(-X, -Y) = e(Y, X)$$

Let $b = B_{\mathfrak{g}} - 2B_{\mathfrak{h}}$ where $B_{\mathfrak{g}}, B_{\mathfrak{h}}$ are the Killing forms of \mathfrak{g} and \mathfrak{h} . Then

$$e(X, Y) = 1 - \frac{1}{240}b([X, Y], [X, Y]) + \frac{1}{1512}b([X, Y], u[X, Y]) + \dots$$

with $u = (\text{ad } X)^2 + \text{ad } X \text{ ad } Y + (\text{ad } Y)^2$.

It is difficult to compute the Jacobian $\det D\Phi$ explicitly. Nevertheless Theorem 7 implies the following:

- (R. 1990 [16]) G/H is special if G is solvable.
- $G_{\mathbb{C}}/G_{\mathbb{R}}$ and $G \times G/\text{diagonal}$ are special for any G (this result relies on the Kashiwara-Vergne conjecture for G).

- (R. 2007) For $p \in S(\mathfrak{s})^{\mathfrak{h}}$ let $p_e(X, \partial_X)f(X) = p(\partial_Y)(e(X, Y)f(X + Y))|_{Y=0}$ as above. Then, for any $f \in C^\infty(\mathfrak{s})$,

$$\tilde{p}f = (p_e f + r f)^\sim, \quad (8)$$

where r is a differential operator on \mathfrak{s} belonging to the left ideal generated by the adjoint vector fields

$$\zeta_V f(X) = \left. \frac{d}{dt} f((\exp -tV) \cdot X) \right|_{t=0}, \quad V \in \mathfrak{h}, \quad X \in \mathfrak{s}.$$

In particular $rf = 0$ whenever f is an H -invariant function. This extends to symmetric spaces Proposition 4.2 of [11] in the case of Lie groups.

Remark. In view of the above factors $J^{1/2}$ it would be more natural to work with half-densities rather than functions on the symmetric space. The theory actually extends to line bundles over G/H ([17]).

2.d. A new construction of e -functions

Adapting Kontsevich's quantization to symmetric spaces ($\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{s}$ as above) Cattaneo and Torossian ([6], §3) define a function $E(X, Y)$ of $X, Y \in \mathfrak{s}$ and a $*$ -product

$$(p * q)(\xi) = E(\partial_\xi, \partial_\eta)(p(\xi)q(\eta))|_{\xi=\eta},$$

with $p, q \in S(\mathfrak{s})$, $\xi \in \mathfrak{s}^*$. When restricted to H -invariant elements p, q , this product is associative and commutative.

They show that

$$\log E(X, Y) = \sum_{\Gamma} w_{\Gamma} (\text{tr}_{\mathfrak{s}}(x_1 x_2 \cdots x_n) + (-1)^n \text{tr}_{\mathfrak{h}}(x_n \cdots x_2 x_1))$$

where $x_i = \text{ad } X_i$ and each X_i consists of iterated brackets of X and Y . The sum runs over a family of graphs Γ adapted from Kontsevich's diagrams, and the w_{Γ} are certain coefficients. The main properties of E can thus be read off "easily" from the graphs.

In particular $E(X, Y) = 1$ if \mathfrak{g} solvable, or for a quadratic Lie algebra viewed as a symmetric pair.

Cattaneo and Torossian also obtain an explicit expression, by means of $*$ -products, of invariant differential operators on G/H written in exponential coordinates ([6], §4).

Besides the product $*$ above, restricted to H -invariant $p, q \in S(\mathfrak{s})$, coincides with \times in Theorem 5

$$\widetilde{p * q} = \widetilde{p \times q} = \widetilde{p} \circ \widetilde{q},$$

as differential operators acting on H -invariant functions.

Thus the function $E(X, Y)$ is very similar to $e(X, Y)$ in Theorem 7 but the proofs of its main properties are much easier (inspection of graphs instead of difficult identities in non-commutative algebra). It is reasonable to conjecture that $E(X, Y)$ and $e(X, Y)$ are (essentially) the same function; see [6] §4.2.2 for a more precise statement. But this remains to be proved.

3. Back to the introduction

For special symmetric spaces the problems mentioned in the introduction are solved by the map $\tilde{\cdot}$. For more general spaces one should expect it can be done by means of an e -function. We now explain this in the case of a Riemannian symmetric space G/K , using the classical semi-simple notations as in Helgason's books [9][10].

4.a. We recall that Harish-Chandra's spherical transform $F_{G/K}$ is related to the Abel transform $\mathcal{A} : \mathcal{D}(G/K)^K \rightarrow \mathcal{D}(\mathfrak{a})^W$ by

$$\int_{G/K} f(x)\varphi_\lambda(x) dx = F_{G/K}f(\lambda) = F_{\mathfrak{a}}\mathcal{A}f(\lambda), \lambda \in \mathfrak{a}^*, \quad (9)$$

where $F_{\mathfrak{a}}f(\lambda) = \int_{\mathfrak{a}} f(H)e^{i\langle \lambda, H \rangle} dH$ is the classical Fourier transform on the vector space \mathfrak{a} .

By the Weyl group invariance $\varphi_{w\lambda} = \varphi_\lambda$ and a suitable version of Chevalley's restriction theorem ([9] p. 468-469) this W -invariant equality in $\lambda \in \mathfrak{a}^*$ extends to a K -invariant equality in $\xi \in \mathfrak{p}^*$:

$$\int_{G/K} f(x)\varphi_\xi(x) dx = F_{G/K}f(\xi) = F_{\mathfrak{p}}\mathcal{T}f(\xi), \xi \in \mathfrak{p}^*, \quad (10)$$

where $F_{\mathfrak{p}}f(\xi) = \int_{\mathfrak{p}} f(X)e^{i\langle \xi, X \rangle} dX$ is the Fourier transform on \mathfrak{p} . This defines an **operator** $\mathcal{T} : \mathcal{D}(G/K)^K \rightarrow \mathcal{D}(\mathfrak{p})^K$ which transforms convolutions of K -invariant functions on G/K into the abelian convolution on \mathfrak{p} and G -invariant differential operators on G/K into K -invariant differential operators on \mathfrak{p} with constant coefficients.

Let $\mathfrak{p} = \mathfrak{a} \oplus \mathfrak{q}$ (orthogonal sum) and, for $\psi \in \mathcal{D}(\mathfrak{p})^K$, $H \in \mathfrak{a}$, $\mathcal{A}_0\psi(H) = \int_{\mathfrak{q}} \psi(H+Y) dY$ (flat Abel transform; see [10] chap. IV §5). Then $F_{\mathfrak{p}}\psi(\lambda) = F_{\mathfrak{a}}\mathcal{A}_0\psi(\lambda)$, $\lambda \in \mathfrak{a}^*$; this equality is the flat analogue of (9). Comparing (9) and (10) we have $F_{\mathfrak{a}}\mathcal{A}f(\lambda) = F_{\mathfrak{p}}\mathcal{T}f(\lambda) = F_{\mathfrak{a}}\mathcal{A}_0\mathcal{T}f(\lambda)$ therefore

$$\mathcal{T} = \mathcal{A}_0^{-1} \circ \mathcal{A}.$$

4.b. This nice operator \mathcal{T} can be related to e -functions in the following way.

Conjecture 8 *Let e be the function introduced in Theorem 7. For $Y \in \mathfrak{p}'$ (a regular element in \mathfrak{p}) the following limit*

$$e_\infty(X, Y) = \lim_{t \rightarrow +\infty} e(X, tY)$$

exists, with uniform convergence when X runs in an arbitrary compact subset of \mathfrak{p} .

I am grateful to M. Flensted-Jensen for suggesting to consider this limit. Clearly, for $k \in K$ and $t > 0$,

$$e_\infty(X, tY) = e_\infty(X, Y) = e_\infty(k \cdot X, k \cdot Y).$$

Examples. As an easy consequence of (5), Conjecture 8 is true for rank one spaces and

$$e_\infty(X, Y) = \left(2 \frac{x}{\operatorname{sh} x} \frac{\operatorname{ch} x - \operatorname{ch} s}{x^2 - s^2} \right)^{(n-3)/2} {}_2F_1 \left(1 - \frac{q}{2}, \frac{q}{2}; \frac{n-1}{2}; \frac{\operatorname{ch} x - \operatorname{ch} s}{2 \operatorname{ch} x} \right) \quad (11)$$

with $X, Y \in \mathfrak{p}$, $\|X\| = x$, $\|Y\| = 1$ and $s = X \cdot Y$ (the K -invariant scalar product on \mathfrak{p} corresponding to the norm $\|\cdot\|$).

Besides Conjecture 8 is obviously true when G/K is special ($e = 1$), i.e. when G admits a complex structure.

Theorem 9 *Assume Conjecture 8. Then the spherical functions (extended to $\xi \in \mathfrak{p}^{*l}$, identified with a regular element of \mathfrak{p} by duality) are*

$$\varphi_\xi(\text{Exp } X) = J(X)^{-1/2} \int_K e^{i\langle \xi, k \cdot X \rangle} e_\infty(k \cdot X, \xi) dk, \quad X \in \mathfrak{p}, \quad (12)$$

and the above operator \mathcal{T} is given by the oscillatory integral

$$\mathcal{T}\tilde{u}(Y) = \int_{\mathfrak{p} \times \mathfrak{p}^{*l}} e^{i\langle \xi, X-Y \rangle} e_\infty(X, \xi) u(X) dX d\xi, \quad u \in \mathcal{D}(\mathfrak{p})^K, \quad Y \in \mathfrak{p}.$$

Thus, up to the map $u \mapsto \tilde{u}$, \mathcal{T} is a **pseudo-differential operator** of order 0 on \mathfrak{p} defined by the e -function. If G admits a complex structure it boils down to $\mathcal{T}\tilde{u} = u$.

Remarks. An expression of spherical functions similar to (12) has been obtained by Duistermaat (1983) [7] by means of a diffeomorphism of the boundary K/M transforming the Iwasawa projection into the orthogonal projection $\mathfrak{p} \rightarrow \mathfrak{a}$. Theorem 6 draws inspiration from his construction, which in turn appears as some boundary limit of ours in the spirit of Conjecture 8. A proof of this conjecture might follow from unifying Duistermaat's diffeomorphism and our Φ in Theorem 6.

For rank one spaces (11) and (12) give back a formula for spherical functions already proved by Koornwinder (1975) [13].

References

- [1] Alekseev, A. and Meinrenken, E., *Poisson geometry and the Kashiwara-Vergne conjecture*, C.R. Acad. Sci. Paris, Série I, 335 (2002), 723-728.
- [2] Alekseev, A. and Meinrenken, E., *On the Kashiwara-Vergne conjecture*, Invent. Math. 164 (2006), 615-634.
- [3] Alekseev, A. and Petracchi, E., *Low order terms of the Campbell-Hausdorff series and the Kashiwara-Vergne conjecture*, J. Lie Theory 16 (2006), 531-538; arXiv:math.QA/0508077.
- [4] Andler, M., Sahi, S. and Torossian, C., *Convolution of invariant distributions: proof of the Kashiwara-Vergne conjecture*, Lett. Math. Phys. 69 (2004), 177-203; arXiv:math.QA/0104100.
- [5] Burgunder, E., *Eulerian idempotent and Kashiwara-Vergne conjecture*, arXiv:math.QA/0612548 (2006).
- [6] Cattaneo, A. and Torossian, C., *Quantification pour les paires symétriques et diagrammes de Kontsevich*, arXiv:math.RT/0609693 (2006).
- [7] Duistermaat, J.J., *On the similarity between the Iwasawa projection and the diagonal part*, Mémoire n°15 Soc. Math. France (1984), 129-138.
- [8] Helgason, S., *Fundamental solutions of invariant differential operators on symmetric spaces*, Amer. J. Maths. 86 (1964), 565-601.
- [9] Helgason, S., *Groups and geometric analysis*, Academic Press 1984.
- [10] Helgason, S., *Geometric analysis on symmetric spaces*, Math. Surveys and Monographs 39, Amer. Math. Soc. 1994.
- [11] Kashiwara, M. and Vergne, M., *The Campbell-Hausdorff formula and invariant hyperfunctions*, Invent. Math. 47 (1978), 249-272.
- [12] Kontsevich, M., *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. 66 (2003), 157-216; arXiv:math.QA/9709040.
- [13] Koornwinder, T., *A new proof of a Paley-Wiener type theorem for the Jacobi transform*, Arkiv för Mat. 13 (1975), 145-159.
- [14] Rouvière, F., *Démonstration de la conjecture de Kashiwara-Vergne pour l'algèbre $sl(2)$* , C.R. Acad. Sci. Paris, Série I 292 (1981), 657-660.
- [15] Rouvière, F., *Espaces symétriques et méthode de Kashiwara-Vergne*, Ann. Sci. École Norm. Sup. 19 (1986), 553-581.
- [16] Rouvière, F., *Invariant analysis and contractions of symmetric spaces*, I, Compositio Math. 73 (1990), 241-270; II, Compositio Math. 80 (1991), 111-136.
- [17] Rouvière, F., *Fibrés en droites sur un espace symétrique et analyse invariante*, J. Funct. Anal. 124 (1994), 263-291.
- [18] Torossian, C., *Sur la conjecture combinatoire de Kashiwara-Vergne*, J. Lie Theory 12 (2002), 597-616.
- [19] Torossian, C., *La conjecture de Kashiwara-Vergne [d'après Alekseev et Meinrenken]*, Séminaire Bourbaki n° 980 (Juin 2007).
- [20] Vergne, M., *Le centre de l'algèbre enveloppante et la formule de Campbell-Hausdorff*, C.R. Acad. Sci. Paris, Série I 329 (1999), 767-772.

Laboratoire J.-A. Dieudonné
Parc Valrose
06108 Nice cedex 2, France
frou@math.unice.fr