

# On the monodromy conjecture for non-degenerate hypersurfaces <sup>\*</sup>

Alexander ESTEROV <sup>†</sup>, Ann LEMAHIEU <sup>‡</sup> and Kiyoshi TAKEUCHI

March 12, 2019

## Abstract

The monodromy conjecture is an umbrella term for several expected relationships between poles of zeta functions, monodromy eigenvalues and roots of Bernstein-Sato polynomials in the context of arithmetic geometry and singularity theory. Even the weakest of these relations — the Denef–Loeser conjecture on topological zeta functions — is not proved even for surface singularities so far.

We prove it for a wide class of multidimensional singularities that are non-degenerate with respect to their Newton polyhedra, including all such singularities of functions of four variables.

A crucial difference from the known case of three variables is the existence of degenerate singularities arbitrarily close to a non-degenerate one. Thus, even aiming at the study of non-degenerate singularities, we have to go beyond the setting of Newton polyhedra and toric resolutions.

Besides this, in the course of the proof we detect other phenomena that look unexpected from the lower-dimensional perspective, and develop new tools to deal with them (some may be of independent interest in convex geometry and analysis; see e.g. a special class of supermodular functions that we call hypermodular). Based on these observations, we conjecture how the proof of the Denef–Loeser conjecture for non-degenerate singularities of arbitrarily many variables might look like.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>The monodromy conjecture for the topological zeta function</b>	<b>5</b>
2.1	The conjecture . . . . .	5
2.2	The topological zeta function and Newton polyhedra . . . . .	6
2.3	The monodromy zeta function and Newton polyhedra . . . . .	8

---

<sup>\*</sup>**2010 Mathematics Subject Classification:** 14M25, 32S40

<sup>†</sup>HSE University

<sup>‡</sup>This research is partially supported by the MCI-Spain grant MTM2010-21740-C02, the ANR ‘SUSI’ project and the ANR ‘Defigeo’ project.

<b>3</b>	<b>Candidate poles of the topological zeta function and <math>B</math>-facets</b>	<b>9</b>
3.1	$B_1$ -faces . . . . .	9
3.2	Critical edges . . . . .	11
3.3	Some non-contributing configurations of $B_1$ -facets . . . . .	12
3.4	$B_2$ -facets . . . . .	15
<b>4</b>	<b>Fake poles of the topological zeta function in arbitrary dimension</b>	<b>17</b>
4.1	Contributions . . . . .	18
4.2	The main theorem: the plan of the proof . . . . .	18
4.3	Intersections of $B$ -facets . . . . .	19
4.4	Bases and apices . . . . .	21
4.5	Sprouts and cancellation of contributions . . . . .	22
4.6	Critical cones . . . . .	23
4.7	A tubular neighborhood of the critical subfan . . . . .	25
<b>5</b>	<b>Fake poles of the topological zeta function in dimension 4</b>	<b>27</b>
5.1	The main theorem . . . . .	27
5.2	Very critical cones . . . . .	29
5.3	A tubular neighborhood of the critical subfan revisited . . . . .	31
5.4	Generalizing the notion of $B$ -facets . . . . .	34
<b>6</b>	<b>Eigenvalues of monodromy and corners</b>	<b>36</b>
6.1	Motivation and results . . . . .	36
6.2	Hypermultiplicities . . . . .	39
6.3	Reduction to the case $k = n - 1$ and $r = n - 1$ . . . . .	39
6.4	The proof of the case $r = n - 1$ . . . . .	40
<b>7</b>	<b>On non-convenient Newton polyhedra</b>	<b>46</b>
7.1	Nearby singularities at coordinate lines . . . . .	46
7.2	Nearby singularities outside coordinate lines . . . . .	47
<b>8</b>	<b>The monodromy conjecture for <math>n = 4</math></b>	<b>49</b>
8.1	Two- and three-dimensional case . . . . .	50
8.2	Dimension 4 : Looking for the eigenvalue $t_0$ outside the origin : part 1. . . . .	51
8.3	Dimension 4 : Looking for the eigenvalue $t_0$ outside the origin : part 2. . . . .	53
8.4	Monodromy conjecture for non-degenerate singularities of four variables . . . . .	56
8.5	Exotic families . . . . .	58

# 1 Introduction

Over the fields  $\mathbb{R}$  and  $\mathbb{C}$  it is well-known that the poles of the local zeta function associated to a polynomial  $f$  are contained in the set of roots of the Bernstein-Sato polynomial and integer shifts of them. By a celebrated theorem of Kashiwara and Malgrange, this implies that for any such pole  $s_0 \in \mathbb{Q}$  the complex number  $\exp(2\pi i s_0) \in \mathbb{C}$  is an eigenvalue of the monodromies of the complex hypersurface defined by  $f$ . Igusa predicted a similar beautiful relationship between the poles of  $p$ -adic integrals and the complex monodromies. This is now called the monodromy conjecture (see the papers of Denef [5], Nicaise [21] and

Denef and Loeser [9] for excellent reviews on this subject). Later in [7], Denef and Loeser introduced the local topological zeta function  $Z_{\text{top},f}(s)$  associated to  $f$  and proposed a weaker version of the monodromy conjecture. However, even this weaker version, proposed a quarter of a century ago, is proved so far without any restrictions in dimension two only ([18]).

For other important contributions to this Denef-Loeser conjecture, see for example [19] by Loeser, where he studies the monodromy conjecture for non-degenerate singularities satisfying some non-resonance conditions. The result holds for isolated singularities. In [16], the second author and Van Proeyen restrict to non-degenerate surface singularities and then prove the monodromy conjecture without further conditions. In the works [2] and [3], Artal Bartolo–Cassou-Noguès–Luengo–Melle Hernández prove the monodromy conjecture for some particular classes of singularities for which they establish an explicit formula for the topological zeta function. Other important contributions to the Denef-Loeser conjecture and related results include [25], [17], [20], [13], [4].

The aim of the present paper is to explore to what extent the results of [16] hold true in higher dimensions, and what we are missing to step one dimension higher for non-degenerate singularities.

A crucial difference that we observe in dimension four is the existence of degenerate singularities arbitrarily close to a non-isolated non-degenerate singularity. So, even aiming at the study of non-degenerate singularity, we have to go beyond the setting of Newton polyhedra and toric resolutions at some point.

This is in contrast to all preceding results on non-isolated singularities: in the three-dimensional setting of [16], all singularities close to a non-degenerate one are non-degenerate, and, in the setting of [3], all singularities close to a quasi-ordinary one are quasi-ordinary.

The paper consists of three parts. In Sections 3–5, we study configurations of facets of the Newton polyhedron that do not assure the existence of the corresponding pole of the topological zeta function. In Sections 6–7, we study configurations of faces that, on the contrary, always non-trivially contribute to the multiplicity of the expected monodromy eigenvalue. Finally, in the last section, we use these results to prove the monodromy conjecture for singularities of non-degenerate functions of four variables.

**Theorem 1.1.** *The Denef–Loeser monodromy conjecture (and moreover its polyhedral version from [11]) holds true for all non-degenerate hypersurface singularities of four variables.*

In Section 2 we recall the exact statement of the monodromy conjecture and the notion of non-degenerate singularity.

In Section 3, as a generalization of the notion of  $B_1$ -facets in [16], we introduce  $B_1$ -facets of the Newton polyhedron  $\Gamma_+(f)$  (Definition 3.10) and discover so called  $B_2$ -facets (Definition 3.9) that behave similarly, although do not exist in the lower-dimensional setting.

In Section 4, we show that many configurations of  $B_1$  and  $B_2$ -facets never assure the existence of the corresponding pole of the topological zeta function (Theorem 4.3). In the course of the proof we introduce an important notion of a critical face of the Newton polyhedron (Definition 4.25). Its role in the proof indicates that it might be possible to find a similarly important notion of a critical stratum of the exceptional divisor in the context of arbitrary singularities and their non-toric resolutions.

In Section 5, we apply the tools from the preceding two sections to completely classify configurations of facets of four-dimensional Newton polyhedra that never assure the existence of the corresponding pole of the topological zeta function (Theorem 5.2). Besides the previously found configurations, we discover so called  $B^2$ -borders (Definition 5.1). It would be important for the further research to detect a proper multidimensional generalization of this object.

In Section 6, following the strategy of [16], we prove that the candidate poles of  $Z_{\text{top},f}(s)$  contributed by certain non- $B_1$ -facets of  $\Gamma_+(f)$  always yield monodromy eigenvalues.

In particular we obtain Theorem 6.4. Its proof relies upon the new notion of a hypermodular function (Definition 6.8), which is inspired by supermodular functions in convex geometry and analysis, and may be of independent interest.

As a corollary, we can confirm the monodromy conjecture of Denef-Loeser for many non-degenerate hypersurfaces in higher dimensions, see e.g. Theorem 6.6.

In Section 7 we prove that singularities adjacent to a Newton non-degenerate singularity along a coordinate line are themselves Newton non-degenerate (see Proposition 7.2).

However, we notice that starting from dimension 4, not all singularities adjacent to Newton non-degenerate singularities are non-degenerate themselves (see Example 7.4). In particular, even in dimension 4 it is not possible to prove the Denef-Loeser conjecture for Newton non-degenerate singularities within the framework of non-degenerate singularities.

To this end, the first author introduced in [11] the notion of tropical nearby monodromy eigenvalues and the corresponding monodromy conjecture, which implies the Denef-Loeser conjecture and (in contrast to the latter) turns into a purely combinatorial statement on the Newton polyhedron for non-degenerate singularities. This tool is essential in the study of monodromy eigenvalues of singularities that are adjacent to a singularity with a given resolution.

In particular, it allows us to prove in Section 8 the monodromy conjecture for non-degenerate functions of four variables: if the sought monodromy eigenvalue is not a tropical monodromy eigenvalue outside the origin, this imposes lots of restrictions on the combinatorial structure of the Newton polyhedron, and a detailed study of this structure shows that the sought monodromy eigenvalue is a root of the monodromy zeta function at the origin.

Note that the order of this reasoning is opposite to the one usually seen in the literature: first try to find the sought monodromy eigenvalue at the origin, then in the case of a trouble switch to nearby singularities.

Altogether, the mentioned observations allow us to conjecture how the proof of the Denef-Loeser conjecture for non-degenerate singularities of arbitrarily many variables might look like, see Plan 5.20.

**Acknowledgements:** The authors are very grateful to the University of Nice for its hospitality.

## 2 The monodromy conjecture for the topological zeta function

In this section, we recall the monodromy conjecture for the topological zeta function and related results.

### 2.1 The conjecture

Let  $f : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$  be a germ of a non-trivial analytic function. We assume that  $f$  is defined on an open neighborhood  $X$  of the origin  $0 \in \mathbb{C}^n$ . Let  $\pi : Y \rightarrow X$  be an embedded resolution of the complex hypersurface  $f^{-1}(0) \subset X$  and  $E_j$  ( $j \in J$ ) the irreducible components of the normal crossing divisor  $\pi^{-1}(f^{-1}(0)) \subset Y$ . For  $j \in J$  we denote by  $N_j$  (resp.  $\nu_j - 1$ ) the multiplicity of the divisor associated to  $f \circ \pi$  (resp.  $\pi^*(dx_1 \wedge \cdots \wedge dx_n)$ ) along  $E_j \subset Y$ . For a non-empty subset  $I \subset J$  we set

$$E_I = \bigcap_{i \in I} E_i, \quad E_I^\circ = E_I \setminus \left( \bigcup_{j \notin I} E_j \right).$$

In [7] Denef and Loeser defined the local topological zeta function  $Z_{\text{top},f}(s) \in \mathbb{C}(s)$  associated to  $f$  (at the origin) by

$$Z_{\text{top},f}(s) = \sum_{I \neq \emptyset} \chi(E_I^\circ \cap \pi^{-1}(0)) \prod_{i \in I} \frac{1}{N_i s + \nu_i},$$

where  $\chi(\cdot)$  denotes the topological Euler characteristic. More precisely, they introduced  $Z_{\text{top},f}(s)$  by  $p$ -adic integrals and showed that it does not depend on the choice of the embedded resolution  $\pi : Y \rightarrow X$  by algebraic methods. Later in [8] and [9], they redefined  $Z_{\text{top},f}(s)$  by using the motivic zeta function of  $f$  and reproved this independence of  $\pi$  more elegantly. For a point  $x \in f^{-1}(0) \cap X$  let  $F_x \subset X \setminus f^{-1}(0)$  be the Milnor fiber of  $f$  at  $x$  and  $\Phi_{j,x} : H^j(F_x; \mathbb{C}) \xrightarrow{\sim} H^j(F_x; \mathbb{C})$  ( $j \in \mathbb{Z}$ ) the Milnor monodromies associated to it. Then the monodromy conjecture of Denef-Loeser for the local topological zeta function  $Z_{\text{top},f}(s)$  is stated as follows.

**Monodromy Conjecture:** (Denef-Loeser [7, Conjecture 3.3.2]) Assume that  $s_0 \in \mathbb{C}$  is a pole of  $Z_{\text{top},f}(s)$ . Then  $\exp(2\pi i s_0) \in \mathbb{C}$  is an eigenvalue of the monodromy  $\Phi_{j,x} : H^j(F_x; \mathbb{C}) \xrightarrow{\sim} H^j(F_x; \mathbb{C})$  for some point  $x \in f^{-1}(0) \cap X$  in a neighborhood of the origin  $0 \in \mathbb{C}^n$  and some  $j \in \mathbb{Z}$ .

In [7] the authors also formulated an even stronger conjecture concerning the Bernstein-Sato polynomial  $b_f(s)$  of  $f$ . Namely they conjectured that the poles of  $Z_{\text{top},f}(s)$  are roots of  $b_f(s)$ .

From now on, we assume that  $f$  is a non-trivial polynomial on  $\mathbb{C}^n$  such that  $f(0) = 0$  and recall the results of Denef-Loeser [7, Section 5] and Varchenko [24]. For  $f(x) = \sum_{v \in \mathbb{Z}_+^n} c_v x^v \in \mathbb{C}[x_1, \dots, x_n]$  ( $\mathbb{Z}_+ = \{m \in \mathbb{Z} \mid m \geq 0\}$ ), its support  $\text{supp} f \subset \mathbb{Z}_+^n$  is the subset

$$\text{supp} f = \{v \in \mathbb{Z}_+^n \mid c_v \neq 0\} \subset \mathbb{Z}_+^n.$$

We denote by  $\Gamma_+(f) \subset \mathbb{R}_+^n$  the convex hull of  $\cup_{v \in \text{supp} f} (v + \mathbb{R}_+^n)$  in  $\mathbb{R}_+^n$ . It is called the Newton polyhedron of  $f$  at the origin  $0 \in \mathbb{C}^n$ . The polynomial  $f$  such that  $f(0) = 0$  is called convenient if  $\Gamma_+(f)$  intersects the positive part of any coordinate axis of  $\mathbb{R}^n$ .

**Definition 2.1.** (Kouchnirenko [15]) The polynomial  $f$  is non-degenerate (at the origin  $0 \in \mathbb{C}^n$ ) if for any compact face  $\tau \prec \Gamma_+(f)$  the complex hypersurface

$$\{x \in (\mathbb{C}^*)^n \mid f_\tau(x) = 0\} \subset (\mathbb{C}^*)^n$$

in  $(\mathbb{C}^*)^n$  is smooth and reduced, where we set

$$f_\tau(x) = \sum_{v \in \tau \cap \mathbb{Z}_+^n} c_v x^v \in \mathbb{C}[x_1, \dots, x_n].$$

It is well-known that generic polynomials having a fixed Newton polyhedron are non-degenerate (see for example [23, Chapter V, paragraph 2]).

## 2.2 The topological zeta function and Newton polyhedra

For  $u = (u_1, \dots, u_n) \in \mathbb{R}_+^n$  we set

$$N(u) = \min_{v \in \Gamma_+(f)} \langle u, v \rangle, \quad \nu(u) = |u| = \sum_{i=1}^n u_i$$

and

$$F(u) = \{v \in \Gamma_+(f) \mid \langle u, v \rangle = N(u)\} \prec \Gamma_+(f).$$

We call  $F(u)$  the supporting face of the vector  $u \in \mathbb{R}_+^n$  on  $\Gamma_+(f)$ . To a face  $\tau \prec \Gamma_+(f)$  one can associate a dual cone

$$\tau^\circ = \overline{\{u \in \mathbb{R}_+^n \mid F(u) = \tau\}} \subset \mathbb{R}_+^n.$$

Note that  $\tau^\circ$  is an  $(n - \dim \tau)$ -dimensional rational polyhedral convex cone in  $\mathbb{R}_+^n$ . The subdivision of  $\mathbb{R}_+^n$  by the cones  $\tau^\circ$  ( $\tau \prec \Gamma_+(f)$ ) satisfies the axiom of fans and is called the dual fan of  $\Gamma_+(f)$ . Let  $\Delta = \mathbb{R}_+ a(1) + \dots + \mathbb{R}_+ a(l)$  ( $a(i) \in \mathbb{Z}_+^n$ ) be a rational simplicial cone in  $\mathbb{R}_+^n$ , where the lattice vectors  $a(i) \in \mathbb{Z}_+^n$  are linearly independent over  $\mathbb{R}$  and primitive. Let  $\text{aff}(\Delta) \simeq \mathbb{R}^l$  be the affine span of  $\Delta$  in  $\mathbb{R}^n$  and  $s(\Delta) \subset \Delta$  the  $l$ -dimensional lattice simplex whose vertices are  $a(1), \dots, a(l)$  and the origin  $0 \in \Delta \subset \mathbb{R}_+^n$ . We denote by  $\text{mult}(\Delta) \in \mathbb{Z}_{>0}$  the  $l$ -dimensional normalized volume  $\text{Vol}_{\mathbb{Z}}(s(\Delta))$  of  $s(\Delta)$ , i.e.  $l!$  times the usual volume of  $s(\Delta)$  with respect to the affine lattice  $\text{aff}(\Delta) \cap \mathbb{Z}^n \simeq \mathbb{Z}^l$  in  $\text{aff}(\Delta)$ . By using this integer  $\text{mult}(\Delta)$  we set

$$J_\Delta(s) = \frac{\text{mult}(\Delta)}{\prod_{i=1}^l \{N(a(i))s + \nu(a(i))\}} \in \mathbb{C}(s).$$

For a face  $\tau \prec \Gamma_+(f)$  we choose a decomposition  $\tau^\circ = \cup_{1 \leq i \leq r} \Delta_i$  of its dual cone  $\tau^\circ$  into rational simplicial cones  $\Delta_i$  of dimension  $l = \dim \tau^\circ$  such that  $\dim(\Delta_i \cap \Delta_j) < l$  ( $i \neq j$ ) and set

$$J_\tau(s) = \sum_{i=1}^r J_{\Delta_i}(s) \in \mathbb{C}(s).$$

By the following result of Denef-Loeser [7], this rational function  $J_\tau(s)$  does not depend on the choice of the decomposition of  $\tau^\circ$ . Let us set

$$\mathbf{1} = \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{R}_+^n.$$

**Lemma 2.2.** (see the proof of [7, Lemme 5.1.1]) *We have an equality*

$$J_\tau(s) = \int_{\tau^\circ} \exp\left(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle\right) du,$$

where  $P$  is a point in  $\tau$  and  $du$  is the  $l$ -dimensional volume form on the affine span  $\text{aff}(\tau^\circ) \simeq \mathbb{R}^l$  for which the volume of the parallelepiped spanned by a basis of the affine lattice  $\text{aff}(\tau^\circ) \cap \mathbb{Z}^n \simeq \mathbb{Z}^l$  is equal to 1.

It is also well-known that one can decompose  $\tau^\circ$  into rational simplicial cones without adding new edges. Then we have the following formula for  $Z_{\text{top},f}(s)$ .

**Theorem 2.3.** (Denef-Loeser [7, Théorème 5.3 (ii)]) *Assume that  $f(x) \in \mathbb{C}[x_1, \dots, x_n]$  is non-degenerate. Then*

$$Z_{\text{top},f}(s) = \sum_{\gamma} J_\gamma(s) + \frac{s}{s+1} \sum_{\tau} (-1)^{\dim \tau} \text{Vol}_{\mathbb{Z}}(\tau) \cdot J_\tau(s),$$

where in the sum  $\sum_{\gamma}$  (resp.  $\sum_{\tau}$ ) the face  $\gamma \prec \Gamma_+(f)$  (resp.  $\tau \prec \Gamma_+(f)$ ) ranges through the vertices of  $\Gamma_+(f)$  (resp. the compact ones such that  $\dim \tau \geq 1$ ) and  $\text{Vol}_{\mathbb{Z}}(\tau) \in \mathbb{Z}_{>0}$  is the  $(\dim \tau)$ -dimensional normalized volume of  $\tau$  with respect to the affine lattice  $\text{aff}(\tau) \cap \mathbb{Z}^n \simeq \mathbb{Z}^{\dim \tau}$  in  $\text{aff}(\tau) \simeq \mathbb{R}^{\dim \tau}$ .

Recall that a face  $\tau$  of  $\Gamma_+(f)$  is called a facet if  $\dim \tau = n - 1$ . For a facet  $\tau \prec \Gamma_+(f)$  let  $a(\tau) = (a(\tau)_1, \dots, a(\tau)_n) \in \tau^\circ \cap \mathbb{Z}_+^n$  be its primitive conormal vector and set

$$N(\tau) = \min_{v \in \Gamma_+(f)} \langle a(\tau), v \rangle,$$

$$\nu(\tau) = |a(\tau)| = \sum_{i=1}^n a(\tau)_i = \langle a(\tau), \mathbf{1} \rangle.$$

We call  $N(\tau)$  the lattice distance of  $\tau$  from the origin  $0 \in \mathbb{R}^n$ . It follows from Theorem 2.3 that any pole  $s_0 \neq -1$  of  $Z_{\text{top},f}(s)$  is contained in the finite set

$$\left\{ -\frac{\nu(\tau)}{N(\tau)} \mid \tau \prec \Gamma_+(f) \text{ is a facet not lying in a coordinate hyperplane} \right\} \subset \mathbb{Q}.$$

Its elements are called candidate poles of  $Z_{\text{top},f}(s)$ .

## 2.3 The monodromy zeta function and Newton polyhedra

Finally we recall the result of Varchenko [24]. For a polynomial  $f(x) \in \mathbb{C}[x_1, \dots, x_n]$  such that  $f(0) = 0$ , its monodromy zeta function  $\zeta_{f,0}(t) \in \mathbb{C}(t)$  at the origin  $0 \in \mathbb{C}^n$  is defined by

$$\zeta_{f,0}(t) = \prod_{j \in \mathbb{Z}} \left\{ \det(\text{id} - t\Phi_{j,0}) \right\}^{(-1)^j} \in \mathbb{C}(t).$$

Similarly one can define also  $\zeta_{f,x}(t) \in \mathbb{C}(t)$  for any point  $x \in f^{-1}(0)$ . Then by considering the decomposition of the nearby cycle perverse sheaf  $\psi_f(\mathbb{C}_{\mathbb{C}^n})[n-1]$  with respect to the monodromy eigenvalues of  $f$  and the concentrations of its components at generic points  $x \in f^{-1}(0)$  (see e.g. [10] and [14]), in order to prove the monodromy conjecture, it suffices to show that for any pole  $s_0 \in \mathbb{C}$  of  $Z_{\text{top},f}(s)$  the complex number  $\exp(2\pi i s_0) \in \mathbb{C}$  is a root or a pole of  $\zeta_{f,x}(t)$  for some point  $x \in f^{-1}(0)$  in a neighborhood of the origin  $0 \in \mathbb{C}^n$  (see Denef [6, Lemma 4.6]).

For a subset  $S \subset \{1, 2, \dots, n\}$  we define a coordinate subspace  $\mathbb{R}^S \simeq \mathbb{R}^{|S|}$  of  $\mathbb{R}^n$  by

$$\mathbb{R}^S = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i = 0 \text{ for any } i \notin S\}$$

and set

$$\mathbb{R}_+^S = \mathbb{R}^S \cap \mathbb{R}_+^n \simeq \mathbb{R}_+^{|S|}.$$

For a compact face  $\tau \prec \Gamma_+(f)$  we take the minimal coordinate subspace  $\mathbb{R}^S$  of  $\mathbb{R}^n$  containing  $\tau$  and set  $s_\tau = |S|$ . If  $\tau$  satisfies the condition  $\dim \tau = s_\tau - 1$  we set

$$\zeta_\tau(t) = \left(1 - t^{N(\tau)}\right)^{\text{Vol}_{\mathbb{Z}}(\tau)} \in \mathbb{C}[t],$$

where  $N(\tau) \in \mathbb{Z}_{>0}$  is the lattice distance of the affine hyperplane  $\text{aff}(\tau) \simeq \mathbb{R}^{\dim \tau}$  in  $\mathbb{R}^S$  from the origin  $0 \in \mathbb{R}^S$ .

**Theorem 2.4.** (Varchenko [24]) *Assume that  $f(x) \in \mathbb{C}[x_1, \dots, x_n]$  is non-degenerate. Then one has*

$$\zeta_{f,0}(t) = \prod_{\tau} \left\{ \zeta_\tau(t) \right\}^{(-1)^{\dim \tau}},$$

where in the product  $\prod_{\tau}$  the face  $\tau \prec \Gamma_+(f)$  ranges through the compact ones satisfying the condition  $\dim \tau = s_\tau - 1$ .

**Definition 2.5.** We say that a face  $\tau$  of  $\Gamma_+(f)$  is a  $V$ -face (or a Varchenko face) if it is compact and satisfies the condition  $\dim \tau = s_\tau - 1$ .

**Definition 2.6.** 1. We say that a candidate pole  $s_0 \in \mathbb{C}$  of  $Z_{\text{top},f}(s)$  is contributed by a facet  $\tau \prec \Gamma_+(f)$  or that  $\tau$  contributes  $s_0$  if we have  $s_0 = -\nu(\tau)/N(\tau)$ .

2. Let  $\sigma$  be a  $V$ -face in  $\Gamma_+(f)$ . We say that  $\sigma$  contributes to (the multiplicity of)  $t_0 \in \mathbb{C}$  if  $t_0$  is a root of the polynomial  $\zeta_\sigma(t)$ .

Also for a general face  $\tau \prec \Gamma_+(f)$  we define its lattice distance  $N(\tau) \in \mathbb{Z}_{>0}$  to be that of the affine hyperplane  $\text{aff}(\tau) \simeq \mathbb{R}^{\dim \tau}$  in  $\text{aff}(\tau \cup \{0\}) \simeq \mathbb{R}^{\dim \tau + 1}$  from the origin  $0 \in \text{aff}(\tau \cup \{0\}) \subset \mathbb{R}^n$ . To end this section, we note the following simple fact.

**Lemma 2.7.** *For two faces  $\tau, \gamma \prec \Gamma_+(f)$  such that  $\gamma \prec \tau$ , we have  $N(\gamma)|N(\tau)$ .*

*Proof.* We may assume that  $\gamma$  is a facet of  $\tau$ . Let  $\Sigma_0$  be the dual fan of  $\Gamma_+(f)$  and  $\Sigma$  a smooth subdivision of  $\Sigma_0$ . Then there exists an  $(n - \dim\gamma)$ -dimensional smooth cone  $\Delta \in \Sigma$  contained in  $\gamma^\circ$  such that  $\dim(\Delta \cap \tau^\circ) = \dim\tau^\circ = n - \dim\tau = (n - \dim\gamma) - 1$ . Set  $l = n - \dim\gamma$  and let  $a(1), \dots, a(l) \in \mathbb{Z}_+^n$  be the primitive vectors on the edges of  $\Delta$ . Then it is easy to see that for any  $v \in \gamma$  we have

$$N(\tau) = \gcd(\langle a(1), v \rangle, \dots, \langle a(l), v \rangle).$$

We have also a similar description of  $N(\tau)$  in terms of the primitive vectors on the edges of  $\Delta \cap \tau^\circ \prec \Delta$ . If we use a point  $v \in \gamma \prec \tau$  to express  $N(\gamma)$  and  $N(\tau)$  simultaneously, we find  $N(\gamma)|N(\tau)$ . This completes the proof.  $\square$

**Remark 2.8.** For two  $V$ -faces  $\tau, \gamma \prec \Gamma_+(f)$  such that  $\gamma \prec \tau$  we can prove Lemma 2.7 more easily as follows. Let  $\mathbb{R}^{S_\tau}$  (resp.  $\mathbb{R}^{S_\gamma}$ ) be the minimal coordinate subspace of  $\mathbb{R}^n$  containing  $\tau$  (resp.  $\gamma$ ) and

$$\text{aff}(\tau) : \langle a(\tau), v \rangle = N(\tau)$$

the equation of the affine hyperplane  $\text{aff}(\tau) \subset \mathbb{R}^{S_\tau}$ . Then by restricting it to  $\mathbb{R}^{S_\gamma} \subset \mathbb{R}^{S_\tau}$  to find that of  $\text{aff}(\gamma)$  we obtain the equality

$$\gcd\{a(\tau)_i \mid (i \in S_\tau)\} \cdot N(\gamma) = N(\tau).$$

### 3 Candidate poles of the topological zeta function and $B$ -facets

In this section, we develop new tools (Lemmas 3.3 and 3.6) to detect configurations of facets contributing fake poles of  $Z_{\text{top},f}(s)$  — so that once a candidate pole is contributed only by facets from this configuration, then it is definitely fake.

As a first example, we define  $B_1$ -pyramid facets (Definition 3.1) of the Newton polyhedron  $\Gamma_+(f)$ . Our definition is a straightforward generalization of that of Lemahieu-Van Proeyen [16]. However, starting from dimension  $n = 4$ , there exist many other combinatorial types of facets and configurations that may contribute fake poles. In particular, we introduce so called  $B_2$ -facets (Definition 3.9) and detect some non-contributing configurations of  $B_1$  and  $B_2$ -facets, see Propositions 3.7, 3.8 and 3.11. The proofs of these facts are intended to motivate general constructions in the next section, where we prove a more general Theorem 4.3.

From now on, we introduce the following convention on figures in this paper: whenever we depict some configuration of cones in  $\mathbb{R}^n$ , we draw its projectivization, resulting in an  $(n - 1)$ -dimensional figure.

#### 3.1 $B_1$ -faces

For a subset  $S \subset \{1, 2, \dots, n\}$  let  $\pi_S : \mathbb{R}_+^n \longrightarrow \mathbb{R}_+^{S^c} \simeq \mathbb{R}_+^{n-|S|}$  be the natural projection. We say that a polyhedron  $\tau$  in  $\mathbb{R}_+^n$  is non-compact for  $S \subset \{1, 2, \dots, n\}$  if the Minkowski sum  $\tau + \mathbb{R}_+^S$  is contained in  $\tau$ .

**Definition 3.1.** (cf. Lemahieu-Van Proeyen [16]) Let  $\tau$  be a polyhedron in  $\mathbb{R}_+^n$ .

1. We say that  $\tau$  is a  $B_1$ -pyramid of compact type for the variable  $x_i$  if  $\tau$  is a compact pyramid over the base  $\gamma = \tau \cap \{x_i = 0\}$  and its unique vertex  $P \prec \tau$  such that  $P \notin \gamma$  has height one from the hyperplane  $\{x_i = 0\} \subset \mathbb{R}_+^n$ .
2. We say that  $\tau$  is a  $B_1$ -pyramid of non-compact type if there exists a non-empty subset  $S \subset \{1, 2, \dots, n\}$  such that  $\tau$  is non-compact for  $S$  and  $\pi_S(\tau) \subset \mathbb{R}_+^{S^c} \simeq \mathbb{R}_+^{n-|S|}$  is a  $B_1$ -pyramid of compact type for some variable  $x_i$  ( $i \notin S$ ).
3. We say that  $\tau$  is a  $B_1$ -pyramid if it is a  $B_1$ -pyramid of compact or non-compact type.
4. We say that a face  $\tau$  of the Newton polyhedron  $\Gamma_+(f)$  is a  $B_1$ -face if it is a  $B_1$ -pyramid. In particular,  $B_1$ -faces of dimension  $n - 1$  and  $1$  will be called  $B_1$ -facets and  $B_1$ -segments respectively.

We shall see later in this section that  $B_1$ -facets alone tend not to contribute poles to the topological zeta function.

**Remark 3.2.** The fact that  $B_1$ -facets might not give rise to eigenvalues of monodromy was already discovered by Loeser (see [19], Remark 6.3). The condition he requires on the facets expels among others all  $B_1$ -facets. Let us recall this condition.

For two distinct facets  $\tau$  and  $\tau'$  of  $\Gamma_+(f)$  let  $\beta(\tau, \tau') \in \mathbb{Z}$  be the greatest common divisor of the  $2 \times 2$  minors of the matrix  $(a(\tau), a(\tau')) \in M(n, 2; \mathbb{Z})$ . Recall that  $a(\tau) \in \tau^\circ \cap \mathbb{Z}^n$  is the primitive conormal vector of  $\tau$ , and  $\beta(\tau, \tau') \in \mathbb{Z}$  is equal to the lattice area of the triangle spanned by  $a(\tau)$  and  $a(\tau')$ .

If  $N(\tau) \neq 0$  (e.g. if  $\tau$  is compact) we set

$$\lambda(\tau, \tau') = \nu(\tau') - \frac{\nu(\tau)}{N(\tau)}N(\tau'), \quad \mu(\tau, \tau') = \frac{\lambda(\tau, \tau')}{\beta(\tau, \tau')} \in \mathbb{Q}.$$

In [19] the author considered only compact facets  $\tau$  of  $\Gamma_+(f)$  which satisfy the following technical condition:

“For any facet  $\tau' \prec \Gamma_+(f)$  such that  $\tau' \neq \tau$  and  $\tau' \cap \tau \neq \emptyset$  we have  $\mu(\tau, \tau') \notin \mathbb{Z}$ .”

He showed that if  $f$  is non-degenerate, the candidate pole of  $Z_{\text{top},f}(s)$  associated to such a compact facet  $\tau$  is a root of the local Bernstein-Sato polynomial of  $f$ .

Now let  $\tau \prec \Gamma_+(f)$  be a facet containing a  $B_1$ -pyramid of compact type for the variable  $x_i$  and set  $\gamma = \tau \cap \{x_i = 0\}$ . Let  $\tau_0 \prec \Gamma_+(f)$  be the unique (non-compact) facet such that  $\gamma \prec \tau_0$ ,  $\tau_0 \neq \tau$  and  $\tau_0 \subset \{x_i = 0\}$ . Then we can easily show that  $\beta(\tau, \tau_0) = \mu(\tau, \tau_0) = 1$ . Namely such a facet  $\tau$  does not satisfy the above-mentioned condition of [19].

The atypical behavior of candidate poles of  $Z_{\text{top},f}(s)$  associated to  $B_1$ -facets essentially arises from the following simple computation (cf. Lemma 2.2).

For a subcone  $C$  of the dual cone  $\tau^\circ$  to a  $k$ -dimensional face  $\tau$  of the Newton polyhedron  $\Gamma_+(f)$ , define the contribution of  $C$  to the topological  $\zeta$ -function  $Z_{\text{top},f}$  as

$$\int_C \exp(-N(u)s - \langle u, \mathbf{1} \rangle) du^{(n)}$$

for  $k = 0$  and otherwise

$$(-1)^k \text{Vol}_{\mathbb{Z}}(\tau) \frac{s}{s+1} \int_C \exp(-N(u)s - \langle u, \mathbf{1} \rangle) du^{(n-k)},$$

where  $N(\cdot)$  is the support function of the Newton polyhedron, and  $du^{(m)}$  is the  $m$ -dimensional lattice volume form. This definition is chosen so that the topological  $\zeta$ -function of  $f$  equals the sum of contributions of the dual cones to all bounded faces of  $\Gamma_+(f)$ .

**Lemma 3.3.** *Assume that a  $B_1$ -face  $\tau$  of  $\Gamma_+(f)$  is the convex hull of its base  $\gamma$  in the coordinate hyperplane  $\{x_n = 0\}$  and its apex  $P = (*, \dots, *, 1)$ . Furthermore, assume that  $C \subset \gamma^\circ$  is the convex hull of a rational polyhedral subcone  $C' \subset \tau^\circ$  and the  $n$ -th coordinate axis  $O_n = \mathbb{R}_+(0, \dots, 0, 1) \subset \mathbb{R}_+^n$ . Then the sum of the contributions from the cones  $C \subset \gamma^\circ$  and  $C' \subset \tau^\circ$  to  $Z_{\text{top},f}(s)$  is equal to*

$$\int_C \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du_1 \cdots du_n$$

if  $\tau$  is a  $B_1$ -segment, and is 0 otherwise.

*Proof.* In the second case, the contributions from  $C$  and  $C'$  are equal up to sign and cancel each other. Indeed, if we decompose  $C'$  into simplicial cones  $\Delta'_i \subset C'$  and take the convex hulls  $\Delta_i \subset C$  of them and the coordinate axis  $O_n = \mathbb{R}_+ \cdot (0, \dots, 0, 1)$ , then by the condition  $P = (*, \dots, *, 1)$  we have  $\text{mult}(\Delta_i) = \text{mult}(\Delta'_i)$ . In the first case, we may assume that  $C'$  is simplicial. Let  $a(i) \in C' \cap \mathbb{Z}_+^n$  ( $1 \leq i \leq n-1$ ) be the primitive vectors on the edges of the  $(n-1)$ -dimensional cone  $C'$ . Then the sum of the contributions is equal to

$$\frac{1}{\prod_{i=1}^{n-1} \{N(a(i))s + \nu(a(i))\}} - \frac{s}{s+1} \frac{1}{\prod_{i=1}^{n-1} \{N(a(i))s + \nu(a(i))\}} \\ = \frac{1}{\{\langle e_n, P \rangle s + \langle e_n, \mathbf{1} \rangle\} \cdot \prod_{i=1}^{n-1} \{\langle a(i), P \rangle s + \langle a(i), \mathbf{1} \rangle\}},$$

where  $e_n = (0, \dots, 0, 1)$ . The right hand side is equal to the sought integral by Lemma 2.2.  $\square$

## 3.2 Critical edges

We now introduce our main tool to prove that a given number is not a pole of the topological zeta function.

**Lemma 3.4.** *Assume that  $s_0 \neq -1$ . Then for the points  $P$  in the summit of a  $B_1$ -facet of  $\Gamma_+(f)$  the equation  $\langle u, P \rangle s_0 + \langle u, \mathbf{1} \rangle = 0$  is non-trivial. Namely it defines a hyperplane  $L_P$  in  $\mathbb{R}^n$ .*

*Proof.* If it is trivial, then we have  $P = -\mathbf{1}/s_0$ . Since  $s_0 \neq -1$ , none of the coordinates of  $P$  is equal to 1, so it is not in the summit.  $\square$

**Definition 3.5.** We say that a closed set  $C \subset \mathbb{R}^n$  is a polyhedral cone if it is a union of finitely many closed convex polyhedral cones. A ray  $R$  on the boundary  $\partial C$  of an  $n$ -dimensional polyhedral cone  $C$  in  $\mathbb{R}^n$  is called an edge of  $C$ , if, in an arbitrarily small neighborhood of a point of  $\text{rel.int}R = R \setminus \{0\}$ , the cone  $C$  is not affinely isomorphic to a product  $\mathbb{R}^2 \times C'$  for some subset  $C' \subset \mathbb{R}^{n-2}$ . Moreover for  $s_0 \in \mathbb{R}$  and a point  $P \in \mathbb{R}^n$  we say that a ray  $R$  in  $\mathbb{R}^n$  is critical with respect to the pair  $(s_0, P)$  if for its generator  $u \in R$  we have  $\langle u, P \rangle s_0 + \langle u, \mathbf{1} \rangle = 0$ , i.e.  $u \perp (s_0 P + \mathbf{1})$ .

**Lemma 3.6.** *Let  $C \subset \mathbb{R}^n$  be an  $n$ -dimensional polyhedral cone in  $\mathbb{R}^n$ . Assume that for  $s_0 \in \mathbb{R}$  and a point  $P \in \mathbb{R}^n$  no edge of  $C$  is critical with respect to  $(s_0, P)$ . Then the integral*

$$\int_C \exp\left(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle\right) du_1 \cdots du_n \quad (3.1)$$

*is a rational function of  $s$  holomorphic at  $s = s_0 \in \mathbb{C}$ .*

*Proof.* Subdivide  $C$  into simplicial cones. This subdivision is combinatorially stable under small perturbation of each ray  $R$  within its ambient face of  $C$ . Since no edge of this ambient face is not critical with respect to  $(s_0, P)$ , the same is true also for almost all rays in the face. We can thus perturb the rays in the subdivision in their ambient faces so that all of them become non-critical with respect to  $(s_0, P)$ . Then by integrating  $\exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle)$  over each of the simplicial cones in the resulting subdivision, we obtain a rational function holomorphic at  $s = s_0 \in \mathbb{C}$  (see Lemma 2.2).  $\square$

### 3.3 Some non-contributing configurations of $B_1$ -facets

We first show that a candidate pole contributed by a unique facet is always fake, once this facet is  $B_1$ . Then we discuss what happens in other cases (when the same candidate pole is contributed by several  $B_1$ -facets or by a non- $B_1$  facet).

The following result is not used in the sequel and, on the contrary, is a special case of the subsequent Theorem 4.3. Nevertheless, we prefer to give it an independent proof, keeping things as simple and explicit as possible. This proof is a good illustration of a more general construction (of so called sprouts) used later on to prove Theorem 5.2 leading to the monodromy conjecture in dimension 4.

**Proposition 3.7.** *Assume that  $f$  is non-degenerate and let  $\tau \prec \Gamma_+(f)$  be a  $B_1$ -facet. Assume also that the candidate pole*

$$s_0 = -\frac{\nu(\tau)}{N(\tau)} \neq -1$$

*of  $Z_{\text{top},f}(s)$  is contributed only by  $\tau$ . Then  $s_0$  is fake, i.e. not an actual pole of  $Z_{\text{top},f}(s)$ .*

*Proof.* Since the proof for  $B_1$ -pyramids of non-compact type is similar, we prove the assertion only for  $B_1$ -pyramids of compact type. Without loss of generality we may assume that  $\tau$  is a compact pyramid over the base  $\gamma = \tau \cap \{x_n = 0\}$  and its unique vertex  $P \prec \tau$  such that  $P \notin \gamma$  has height one from the hyperplane  $\{x_n = 0\} \subset \mathbb{R}_+^n$ . Let  $A_1, A_2, \dots, A_m$  ( $m \geq n - 1$ ) be the vertices of the  $(n - 2)$ -dimensional polytope  $\gamma$ . For

$1 \leq i \leq m$  we denote the dual cone  $A_i^\circ$  of  $A_i \prec \Gamma_+(f)$  by  $C_{A_i}$ . Similarly we set  $C_P = P^\circ$ . Let  $a(\tau) \in \mathbb{Z}_+^n$  be the primitive vector on the ray  $\tau^\circ$ . Then we have

$$a(\tau) \in \text{Int}(C_P \cup C_{A_1} \cup \cdots \cup C_{A_m}).$$

Note that  $C_P \cup C_{A_1} \cup \cdots \cup C_{A_m}$  is an  $n$ -dimensional polyhedral cone in the sense of Definition 3.5. In order to construct a nice  $n$ -dimensional polyhedral subcone  $\square$  of  $C_P \cup C_{A_1} \cup \cdots \cup C_{A_m}$  such that  $a(\tau) \in \text{Int}\square$ , we shall introduce a new dummy vector  $b \in \text{Int}C_P \cap \mathbb{Z}_+^n$  satisfying the condition

$$-\frac{\nu(b)}{N(b)} \neq s_0 \quad (3.2)$$

in the following way. First, by our assumption  $s_0 \neq -1$  and Lemma 3.4, for the summit  $P = (*, *, \dots, *, 1) \in \mathbb{Z}_+^n$  of the  $B_1$ -pyramid  $\tau$  the equation  $\langle u, P \rangle s_0 + \langle u, \mathbf{1} \rangle = 0$  is non-trivial. It thus defines a hyperplane  $L_P$  in  $\mathbb{R}^n$ . Then by taking a primitive vector  $b \in \text{Int}C_P \cap \mathbb{Z}_+^n$  such that  $b \notin L_P$  we get the desired condition  $N(b)s_0 + \nu(b) \neq 0$ . Let  $\tau_0 \prec \Gamma_+(f)$  be the unique facet such that  $\gamma \prec \tau_0$ ,  $\tau_0 \neq \tau$  and  $\tau_0 \subset \{x_n = 0\}$ . Then the primitive vector  $a(\tau_0) \in \mathbb{Z}_+^n$  on the ray  $\tau_0^\circ$  is given by  $a(\tau_0) = (0, 0, \dots, 0, 1)$ . For  $1 \leq i \leq m$  let  $\sigma_i \prec \tau$  be the edge of  $\tau$  connecting the two points  $P$  and  $A_i$  and  $F_i \prec C_P$  the corresponding facet of the cone  $C_P$  containing  $\tau^\circ = \mathbb{R}_+a(\tau) \prec C_P$ . All the facets of  $C_P$  containing  $\tau^\circ$  are obtained in this way. Since the point  $A_i \prec \sigma_i$  is a vertex of  $\tau_0$ , its dual cone  $C_{A_i} = A_i^\circ$  contains not only  $F_i$  but also the ray  $\tau_0^\circ = \mathbb{R}_+a(\tau_0)$ . For  $1 \leq i \leq m$  set

$$F_i^\sharp = \mathbb{R}_+a(\tau_0) + F_i, \quad F_i^\flat = \mathbb{R}_+b + F_i.$$

In Figure 1 below we presented the transversal hyperplane sections of the cones  $F_i$ ,  $F_i^\sharp$  and  $F_i^\flat$ .

Then by our construction,  $\square = \cup_{i=1}^m (F_i^\sharp \cup F_i^\flat)$  is an  $n$ -dimensional polyhedral cone in  $\mathbb{R}^n$  and satisfies the desired condition  $a(\tau) \in \text{Int}\square$ . By Lemma 3.3 (or the argument in Case 1 of the proof of [16, Proposition 14]) we obtain an equality

$$Z_{\text{top},f}(s) \equiv \int_{\square} \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du_1 \cdots du_n$$

modulo holomorphic functions at  $s = s_0 \in \mathbb{C}$ . By our assumption and the condition (3.2) no edge of the polyhedral cone  $\square$  is critical with respect to  $(s_0, P)$  in the sense of Definition 3.5. Then by Lemma 3.6 the rational function

$$\int_{\square} \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du_1 \cdots du_n$$

of  $s$  is holomorphic at  $s = s_0 \in \mathbb{C}$ . This implies that also  $Z_{\text{top},f}(s)$  is holomorphic there.  $\square$

We now discuss what happens when the same candidate pole is contributed by several  $B_1$ -facets.

If two  $B_1$ -facets for different variables are adjacent (i. e. have a common  $(n-2)$ -dimensional face) and contribute to the same candidate pole, this may happen to be an actual pole of the topological  $\zeta$ -function. This is always so for  $n=3$  (see [16]), but may fail starting from  $n=4$  (see  $B^2$ -borders in Theorem 5.2).




Figure 1: The proof of Proposition 3.7

On the contrary, two adjacent  $B_1$ -faces for the same variable cannot alone yield an actual pole (similarly to [16, Proposition 14] for  $n = 3$ ).

This result is not used in the sequel and, on the contrary, is a special case of the subsequent Theorem 4.3. Nevertheless, we prefer to give it an independent proof, keeping things as simple and explicit as possible. This proof is a good illustration of a more general phenomenon (so called critical cones) that we later encounter in the proof of Theorem 5.2 leading to the monodromy conjecture in dimension 4. The following result is a higher-dimensional analogue of the one in the proof of [16, Proposition 14].

**Proposition 3.8.** *Assume that  $f$  is non-degenerate and let  $\tau_1, \dots, \tau_k \prec \Gamma_+(f)$  be  $B_1$ -facets such that*

$$s_0 := -\frac{\nu(\tau_1)}{N(\tau_1)} = \dots = -\frac{\nu(\tau_k)}{N(\tau_k)} \neq -1$$

*and their common candidate pole  $s_0 \in \mathbb{Q}$  of  $Z_{\text{top},f}(s)$  is contributed only by them. Assume also that if  $\tau_i$  and  $\tau_j$  ( $i \neq j$ ) have a common facet they are  $B_1$ -pyramids for the same variable. Then  $s_0$  is fake i.e. not an actual pole of  $Z_{\text{top},f}(s)$ .*

*Proof.* If  $\tau_i$  and  $\tau_j$  ( $i \neq j$ ) do not have a common facet, then by the proof of Proposition 3.7 after a suitable subdivision of the dual fan of  $\Gamma_+(f)$  into rational simplicial cones we can calculate their contributions to  $Z_{\text{top},f}(s)$  separately. So we may assume that the  $B_1$ -facets  $\tau_1, \dots, \tau_k$  have the common summit  $P \in \tau_1 \cap \dots \cap \tau_k$ . For the sake of simplicity, here we shall treat only the case where  $k = 2$ ,  $\tau_1$  (resp.  $\tau_2$ ) is a compact  $B_1$ -pyramid over the base  $\gamma_1 = \tau_1 \cap \{v_n = 0\}$  (resp.  $\gamma_2 = \tau_2 \cap \{v_n = 0\}$ ) and  $\tau_1 \cap \tau_2$  is the (unique) common facet of  $\tau_1$  and  $\tau_2$ . The proofs for the other cases are similar. Let  $\tau_0 \prec \Gamma_+(f)$  be the unique

facet of  $\Gamma_+(f)$  such that  $\gamma_1, \gamma_2 \prec \tau_0$ ,  $\tau_0 \neq \tau_i$  ( $i = 1, 2$ ) and  $\tau_0 \subset \{v_n = 0\}$ . We denote by  $P$  the common summit of  $\tau_1$  and  $\tau_2$ . Since  $\tau_1 \cap \tau_2$  is a common facet of  $\tau_1$  and  $\tau_2$ , there exists a 2-dimensional face of the dual cone  $C_P$  of  $P \prec \Gamma_+(f)$  containing both  $\tau_1^\circ = \mathbb{R}_+ a(\tau_1)$  and  $\tau_2^\circ = \mathbb{R}_+ a(\tau_2)$ . As in the proof of Proposition 3.7, let  $F_1, F_2, \dots, F_m$  be the facets of  $C_P$  containing the ray  $\tau_1^\circ$  or  $\tau_2^\circ$  and subdivide  $F_1 \cup \dots \cup F_m \subset \partial C_P$  into rational simplicial cones without adding new edges. Let  $\Delta_1, \dots, \Delta_r$  be the  $(n-1)$ -dimensional simplicial cones thus obtained in  $F_1 \cup \dots \cup F_m \subset \partial C_P$  and containing  $\tau_1^\circ$  or  $\tau_2^\circ$ . As in the proof of Proposition 3.7, we take a new primitive vector  $b \in \text{Int}C_P \cap \mathbb{Z}_+^n$  such that

$$-\frac{\nu(b)}{N(b)} \neq s_0. \quad (3.3)$$

For  $1 \leq i \leq r$  set

$$\Delta_i^\sharp = \mathbb{R}_+ a(\tau_0) + \Delta_i, \quad \Delta_i^\flat = \mathbb{R}_+ b + \Delta_i.$$

Then  $\square := \cup_{i=1}^r (\Delta_i^\sharp \cup \Delta_i^\flat)$  is an  $n$ -dimensional polyhedral cone in  $\mathbb{R}^n$  such that

$$a(\tau_1), a(\tau_2) \in \text{Int}\square.$$

By Lemma 3.3 (or the argument in Case 1 of the proof of [16, Proposition 14]) we obtain an equality

$$Z_{\text{top},f}(s) \equiv \int_{\square} \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du_1 \cdots du_n$$

modulo holomorphic functions at  $s = s_0 \in \mathbb{C}$ . By our assumption and the condition (3.3) no edge of the polyhedral cone  $\square$  is critical with respect to  $(s_0, P)$  in the sense of Definition 3.5. Then by Lemma 3.6 the rational function

$$\int_{\square} \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du_1 \cdots du_n$$

of  $s$  is holomorphic at  $s = s_0 \in \mathbb{C}$ . □

### 3.4 $B_2$ -facets

We now discuss what happens when a candidate pole is contributed by a non- $B_1$ -facet. Such pole  $s_0$  is always allowed not to be fake for  $n = 3$ , because  $\exp(2\pi i s_0)$  is always a nearby monodromy eigenvalue (see [16]). However, this is not the case starting from  $n = 4$  for some non- $B_1$  facets. In this subsection, we introduce one such example.

**Definition 3.9.** For all  $n \geq 4$  we define  $B_2$ -facets  $\tau \prec \Gamma_+(f)$  to be non- $B_1$  compact facets whose projection to a certain  $(n-2)$ -dimensional coordinate plane coincides with the standard  $(n-2)$ -dimensional simplex.

In particular, for  $n = 4$ , a facet  $\tau$  of  $\Gamma_+(f)$  is a  $B_2$ -facet if and only if, up to reordering the coordinates, it has the vertices  $A, B, P, Q, X, Y$  of the form

$$\left\{ \begin{array}{l} A = (1, 0, *, *) \\ B = (1, 0, *, *) \\ P = (0, 1, *, *) \\ Q = (0, 1, *, *) \\ X = (0, 0, *, *) \\ Y = (0, 0, *, *) \end{array} \right.$$

as in Figure 2 below (it can be degenerated so that  $X = Y$ ).




Figure 2: a  $B_2$ -facet in dimension 4

Note that this facet  $\tau$  splits into two  $B_1$ -pyramids for different variables whose intersection does not contain any 1-dimensional  $V$ -face. For example we have the decomposition  $\tau = AXPQ \cup QXABY$ .

**Definition 3.10.** A facet  $\tau$  of  $\Gamma_+(f)$  is called a  $B$ -facet if it is a  $B_1$ -facet or a  $B_2$ -facet.

The following result is not used in the sequel and, on the contrary, is a special case of the subsequent Theorem 4.3. Nevertheless, we prefer to give it an independent proof for  $n = 4$ , keeping things as simple and explicit as possible. This proof is a good illustration of a more general construction (of so called delimiter hyperplanes) that we later need in the proof of Theorem 5.2 leading to the monodromy conjecture in dimension 4.

**Proposition 3.11.** *In the case  $n \geq 4$  assume that  $f$  is non-degenerate and let  $\tau \prec \Gamma_+(f)$  be a  $B_2$ -facet. Assume also that the candidate pole*

$$s_0 = -\frac{\nu(\tau)}{N(\tau)} \neq -1$$

*of  $Z_{\text{top},f}(s)$  is contributed only by  $\tau$ . Then  $s_0$  is fake.*

*Proof.* We prove the assertion only for  $n = 4$ . The proof for the general case  $n \geq 4$  is similar. In the notation of Figure 2, we define facets  $\sigma_1, \sigma_2, \sigma_3$  of  $\tau$  by  $\sigma_1 = XAP$ ,  $\sigma_2 = PABQ$ ,  $\sigma_3 = YQB$  respectively. As in the proof of [16, Proposition 14] let  $\tau_i \prec \Gamma_+(f)$  ( $1 \leq i \leq 3$ ) be the unique facet such that  $\sigma_i \prec \tau_i$  and  $\tau_i \neq \tau$ . Moreover for  $i = 1, 2$  let  $\rho_i \prec \Gamma_+(f)$  be the unique facet such that  $\rho_i \subset \{x_i = 0\} \simeq \mathbb{R}^3$ ,  $\tau \cap \{x_i = 0\} \prec \rho_i$  and  $\rho_i \neq \tau$ . Since the three segments  $PQ$ ,  $AB$  and  $XY$  are parallel, their dual cones are on the same hyperplane  $H \simeq \mathbb{R}^3$  in  $\mathbb{R}^4$ . By the four facets  $\tau, \tau_1, \tau_2, \rho_1$  containing the vertex  $P$  we define a 4-dimensional simplicial cone  $\Delta_P \subset \mathbb{R}^4$  by

$$\Delta_P = \mathbb{R}_+a(\tau) + \mathbb{R}_+a(\tau_1) + \mathbb{R}_+a(\tau_2) + \mathbb{R}_+a(\rho_1).$$

Similarly we define 4-dimensional simplicial cones  $\Delta_A, \Delta_X, \Delta_Q, \Delta_B, \Delta_Y \subset \mathbb{R}^4$  for the vertices  $A, X, Q, B, Y$  and set

$$\square_P = \Delta_P \cup \Delta_A \cup \Delta_X, \quad \square_Q = \Delta_Q \cup \Delta_B \cup \Delta_Y.$$

Then the dual ray  $\tau^\circ$  of  $\tau$  is contained in  $\square_P \cap H = \square_Q \cap H = \square_P \cap \square_Q$ , but it is an edge of neither  $\square_P$  nor  $\square_Q$ . Moreover  $\square = \square_P \cup \square_Q$  is a 4-dimensional polyhedral cone such that  $a(\tau) \in \text{Int}\square$ . By Lemmas 2.2 and 3.3 we obtain the equality

$$Z_{\text{top},f}(s) \equiv \int_{\square_P} \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du_1 \cdots du_4 + \int_{\square_Q} \exp(-\langle u, Q \rangle s - \langle u, \mathbf{1} \rangle) du_1 \cdots du_4$$

modulo holomorphic functions at  $s = s_0 \in \mathbb{C}$ . Here we used the fact that the normalized area of the quadrilateral face  $\sigma_2 = PABQ$  of  $\tau$  is equal to the sum of the lengths of the segments  $PQ$  and  $AB$  (see Lemma 4.20 for the details). Then, by Lemma 3.6, we can show that the rational function

$$\int_{\square_P} \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du_1 \cdots du_4$$

of  $s$  is holomorphic at  $s = s_0 \in \mathbb{C}$ . The same is true also for the integral over  $\square_Q$ . Hence  $Z_{\text{top},f}(s)$  is holomorphic at  $s = s_0 \in \mathbb{C}$ .  $\square$

## 4 Fake poles of the topological zeta function in arbitrary dimension

In view of the observations from the preceding section, the following result does not look unexpected. From now on, by faces we mean faces of the Newton polyhedron  $\Gamma_+(f)$ , unless explicitly stated otherwise.

**Definition 4.1.** 1. A one-element set  $\{i\} \subset \{1, \dots, n\}$  is called a base direction for a compact  $B_1$ -facet  $\tau$ , if the  $i$ -th coordinate equals 1 for one vertex of  $\tau$ , and equals 0 for the other vertices.

2. An  $(n-2)$ -element set  $I \subset \{1, \dots, n\}$  is called a base direction for a  $B_2$ -facet, if its projection to the  $I$ -th coordinate plane is the standard  $(n-2)$ -dimensional simplex.

Note that a  $B_1$ -facet may have more than one base direction.

**Definition 4.2.** A collection of  $B$ -facets is said to be consistent, if their base directions can be chosen so that:

a pair of  $B$ -facets in the collection have a common facet  $\Rightarrow$   
the intersection of their base directions is non-empty.

**Theorem 4.3.** *Assume that  $f$  is non-degenerate and let  $\tau_1, \dots, \tau_k \prec \Gamma_+(f)$  be  $B$ -facets such that*

$$s_0 = -\frac{\nu(\tau_1)}{N(\tau_1)} = \dots = -\frac{\nu(\tau_k)}{N(\tau_k)} \neq -1$$

*and their common candidate pole  $s_0 \in \mathbb{Q}$  of  $Z_{\text{top},f}(s)$  is contributed only by them. If they are consistent, then  $s_0$  is fake i.e. not an actual pole of  $Z_{\text{top},f}(s)$ .*

This section is devoted to the proof of Theorem 4.3.

In the course of the proof we introduce some new tools that will be used in the next section, which is devoted to a sharper version of Theorem 4.3, completely classifying configurations of  $B$ -faces contributing fake poles for  $n = 4$ . This classification is a key point in the proof of the monodromy conjecture for  $n = 4$ .

## 4.1 Contributions

**Definition 4.4.** Let  $S \subset \mathbb{R}_+^n$  be a polyhedral set, that is, a disjoint union of the relative interiors of some (finitely many) closed convex polyhedral cones in  $\mathbb{R}_+^n$ . Then we define its contribution  $Z(S)(s) \in \mathbb{C}(s)$  (to the topological zeta function  $Z_{\text{top},f}$ ) by

$$\int_S \exp(-N(u)s - \langle u, \mathbf{1} \rangle) du_{\mathbb{R}_+^n} + \frac{s}{s+1} \sum_{\tau} (-1)^{\dim \tau} \text{Vol}_{\mathbb{Z}}(\tau) \int_{S \cap \tau^\circ} \exp(-N(u)s - \langle u, \mathbf{1} \rangle) du_{\tau^\circ},$$

where  $\tau$  ranges through all the positive-dimensional compact faces of  $\Gamma_+(f)$ ,  $N(\cdot)$  is the support function of  $\Gamma_+(f)$ , and  $du_C$  is the lattice volume form on a rational polyhedral cone  $C$  (so that all components of dimension smaller than  $\dim C$  in the intersection  $S \cap C$  do not affect the integral with respect to  $du_C$ ).

**Remark 4.5.** 1. As a function of  $S$ , the contribution to the topological zeta function is an additive measure.

2. By Lemma 2.2, the contribution  $Z(\mathbb{R}_+^n)$  of the open positive quadrant  $\mathbb{R}_+^n$  equals the topological zeta function of a generic  $f$  with the given Newton polyhedron  $\Gamma_+(f)$ .
3. We do not assume the argument  $S$  to be closed or open, because  $Z(S)$  changes significantly when passing to the closure or the interior of  $S$ , and indeed we shall need sets  $S$  that are neither closed nor open.

## 4.2 The main theorem: the plan of the proof

I. Very loosely, the proof of Theorem 4.3 will consist of constructing a particular subdivision of  $\mathbb{R}_+^n$  into pieces  $\sigma_i$  such that  $Z(\sigma_i)$  has no pole at  $s = s_0$ . The boldest hope would be to choose  $\sigma_i$  so that the key Lemma 3.6 is directly applicable to every  $\sigma_i$ , i.e.:

- Every  $\sigma_i$  is contained in the dual cone to some vertex  $P$  of the Newton polyhedron;
- No edge of  $\sigma_i$  is critical with respect to  $(s_0, P)$  in the sense of Lemma 3.6.

For instance, this is done in detail in the proof of Proposition 3.7 for the case of a unique facet contributing to the pole  $s_0$ .

II. Unfortunately, in general step (I) is not realistic as written, because some of the edges of the dual cone  $P^\circ$  will be critical, and they also have to be edges for some  $\sigma_i$ . These critical edges are exactly the ones dual to the contributing facets  $\tau_i$  of the Newton polyhedron.

Fortunately, all such facets  $\tau_i$  are  $B$ -facets in the assumptions of Theorem 4.3. This will help us to surround every such critical edge  $\tau_i^\circ$  by its personal conic neighborhood  $\sigma'_i$  (so called *sprout*) such that Lemma 3.6 is still applicable to it:

- The contribution  $Z(\sigma'_i)$  can be written as the integral from Lemma 3.6 for some appropriate vertex  $P_i$  (despite  $\sigma'_i$  is not contained in any individual cone of the form  $P^\circ$  anymore!)

- No edge of  $\sigma'_i$  is critical with respect to  $(s_0, P_i)$ .

III. Unfortunately, upon choosing neighborhoods in Step (II), we observe the next (and the last) obstacle: the complement to  $\bigcup_i \sigma'_i$  in any cone  $P^\circ$  may have new critical edges in the boundary of  $P^\circ$  (different from the edges of  $P^\circ$  itself). This happens because some cones (so called *critical cones*) in the dual fan  $\Sigma_0$  entirely (!) consist of critical rays,

so every 0-dimensional intersection of a face of  $\sigma'_i$  with such critical cone  $C$  will create a new critical edge of the complement of  $\sigma'_i$  in any cone containing  $C$ .

**Example 4.6.** Recall that here and in what follows we draw the projectivization of the fan  $\Sigma_0$  rather than the fan itself. On the left picture of Figure 3 below, the critical 1-dimensional cone  $\tau_1^\circ$ , whose dual facet  $\tau_1$  is a  $B_1$ -pyramid with the apex  $P$  with respect to the variable  $x_1$ , is surrounded with a conical neighborhood  $\sigma'_1$  (shown in bold). Since the dashed segment is a critical 2-dimensional cone, its intersection point with the boundary of  $\sigma'_1$  is a critical ray that is not an edge of  $\sigma'_1$ , but is a critical ray of the complement to  $\sigma'_1$  in the 3-dimensional cone  $P^\circ$ .




Figure 3: good neighborhoods of critical cones

Fortunately, choosing the neighborhoods  $\sigma'_i$  wisely, we can ensure that no new critical edge of the complement of an individual  $\sigma'_i$  is an edge of the complement to the whole  $\bigcup_i \sigma'_i$  (see the picture on the right of Figure 3). For instance, this is done in detail in the proof of Proposition 3.8 for the case of only two facets contributing the pole  $s_0$ .

In the general setting, this will be done using the geometry of so called *delimiter planes* and will allow to literally apply Step (I) of our plan to the complement of  $\bigcup_i \sigma'_i$ . Warning: the resulting cones  $\sigma'_i$  will not form a fan in the sense that  $\sigma'_i \cap \sigma'_j$  may be not a face of  $\sigma'_i$  and  $\sigma'_j$ .

We introduce all the aforementioned objects in the subsequent subsections.

### 4.3 Intersections of $B$ -facets

**Lemma 4.7.** *Assume that  $f$  is non-degenerate and does not have a Morse singularity at the origin  $0 \in \mathbb{C}^n$ . Then, if two  $B$ -facets of the Newton polyhedron  $\Gamma_+(f)$  have a common  $(n-2)$ -dimensional face, then it is a  $B_1$ -pyramid.*

The proof of this lemma requires the following observation.

**Lemma 4.8.** *If a homogeneous polynomial  $g$  of degree 2 on  $\mathbb{C}^4$  is non-degenerate with respect to its Newton polyhedron and its support  $\text{supp } g$  contains the points  $(1, 1, 0, 0)$  and  $(0, 0, 1, 1)$ , then  $g$  is non-degenerate as a quadratic form.*

*Proof.* The non-degeneracy of  $g$  as a quadratic form is equivalent to the smoothness of the hypersurface  $V = \{g = 0\}$  of  $\mathbb{P}^3$  defined by  $g$ . Indeed, the kernel of the symmetric matrix

associated to the quadratic form corresponds to the singular locus of  $V \subset \mathbb{P}^3$ . Recall that  $\mathbb{P}^3$  is naturally a toric variety on which the complex torus  $T = (\mathbb{C}^*)^3$  acts with 15 orbits. First of all, by the non-degeneracy of  $g$  w.r.t. its Newton polyhedron, the hypersurface  $V = \{g = 0\}$  of  $\mathbb{P}^3$  is smooth in  $T \subset \mathbb{P}^3$ . So we have only to analyze at other points  $x$  in  $\mathbb{P}^3 \setminus T$ . We shall do it step by step, considering  $x$  in each of the other 14  $T$ -orbits in  $\mathbb{P}^3$ . We denote the Newton polytope of  $g$  by  $Q \subset \{x_1 + x_2 + x_3 + x_4 = 2\}$ .

1) Assume that  $x = (1 : 0 : 0 : 0)$ . If the point  $(2, 0, 0, 0)$  is in  $\text{supp}g$ , then  $g(x) \neq 0$ . Otherwise, we have  $dg(x) \neq 0$ , because  $(1, 1, 0, 0) \in \text{supp}g$  and hence  $\partial g / \partial x_2(x) \neq 0$ . This implies that the hypersurface  $V = \{g = 0\}$  of  $\mathbb{P}^3$  is smooth at  $x \in V$ .

2) Assume that  $x$  is in one of the other three 0-dimensional  $T$ -orbits in  $\mathbb{P}^3$ . Then the reasoning is the same as in (1).

3) Assume that  $x = (s : t : 0 : 0)$  ( $s, t \neq 0$ ). Then, since the face  $F = Q \cap [(2, 0, 0, 0), (0, 2, 0, 0)]$  of  $Q$  is non-empty (containing at least the point  $(1, 1, 0, 0)$ ) and the restriction  $g_F$  of  $g$  to the face  $F \prec Q$  defines a smooth hypersurface in the 1-dimensional  $T$ -orbit in  $\mathbb{P}^3$  associated to  $[(2, 0, 0, 0), (0, 2, 0, 0)]$  (by the non-degeneracy of  $g$ ), the hypersurface  $V = \{g = 0\}$  of  $\mathbb{P}^3$  is smooth at  $x$ .

4) Assume that  $x = (0 : 0 : s : t)$  ( $s, t \neq 0$ ) or  $x$  is in one of the four 2-dimensional  $T$ -orbits in  $\mathbb{P}^3$ . Then the reasoning is the same as in (3).

5) Assume that  $x = (s : 0 : t : 0)$  ( $s, t \neq 0$ ). If  $\text{supp}g \cap [(2, 0, 0, 0), (0, 0, 2, 0)]$  is not empty, then the reasoning is the same as in (3). Otherwise, we have  $g(x) = 0$ . Assume that we have also  $dg(x) = 0$ . Then in particular  $\partial g / \partial x_2(x) = \partial g / \partial x_4(x) = 0$ . From these identities, we see that the restrictions  $g_A, g_B$  of  $g$  to the segments  $A = [(1, 1, 0, 0), (0, 1, 1, 0)]$ ,  $B = [(1, 0, 0, 1), (0, 0, 1, 1)]$  vanish at  $x$ , thus they are multiples of the same linear function. So the restriction  $g_P$  of  $g$  to the parallelogram  $P = \text{conv}(A \cup B)$  is a product of two linear functions. This implies that  $g_P$  defines a singular hypersurface in  $(\mathbb{C}^*)^4$ . This would contradict the non-degeneracy of  $g$  with respect to its Newton polyhedron.

6) Assume that  $x$  is in one of the other three 1-dimensional  $T$  orbits in  $\mathbb{P}^3$ . Then the reasoning is the same as in (5).  $\square$

**Lemma 4.9.** *Assume that  $n = 4$ , and two  $B_2$ -facets of  $\Gamma_+(f)$  have a common quadrilateral face. Then  $f$  has a Morse singularity at the origin  $0 \in \mathbb{C}^4$ .*

*Proof.* Projectivizing the ambient space of  $\Gamma_+(f)$ , we see the positive octant as a tetrahedron, and the two  $B_2$ -facets in it as two polytopes from Figure 4 with a common quadrilateral face. This is only possible if the common quadrilateral is a parallelogram, whose edges are parallel to two opposite edges of the tetrahedron, and whose vertices are contained in the four other edges, as shown on the picture.

Thus, reordering coordinates if necessary, the vertices are of the form  $(*, *, 0, 0)$ ,  $(0, *, *, 0)$ ,  $(0, 0, *, *)$ ,  $(*, 0, 0, *)$ , and all the stars are equal to 1 by the definition of  $B_2$ -facets. By Lemma 4.8, the quadratic part of  $f$  is non-degenerate, and hence  $f$  has a Morse singularity at the origin  $0 \in \mathbb{C}^4$ .  $\square$

*Proof of Lemma 4.7.* The only  $(n - 2)$ -dimensional non- $B_1$  face of a  $B$ -facet is the non- $V$  face of a  $B_2$ -facet combinatorially isomorphic to the product of a segment and an  $(n - 3)$ -simplex (let us call it the front face of a  $B_2$ -facet). So the only exception from the statement of the lemma could come from two  $B_2$ -facets with different base directions and the common front face. However, for  $n > 4$  this is impossible, because two products




Figure 4:  $B_2$ -facets with a common quadrangulal face

of a segment and an  $(n - 3)$ -simplex cannot be non-trivially combinatorially isomorphic, and for  $n = 4$  this is excluded by Lemma 4.9.  $\square$

#### 4.4 Bases and apices

**Definition 4.10.** The star of a compact face  $\sigma$  of the polyhedron  $\Gamma_+(f)$  is the set of all compact faces containing  $\sigma$ .

Similarly, the star of a cone  $C$  in the dual fan of  $\Gamma_+(f)$  is the set of all cones containing  $C$ .

For each  $B_1$ -face  $\tau$  we can choose its apex and preferred base

to be its vertex  $P$  and a number  $i \in \{1, \dots, n\}$  respectively, such that the  $i$ -th coordinate of  $P$  equals 1 and the  $i$ -th coordinates of the other vertices of  $\tau$  equals 0. Note that we may have several options for this choice.

We now fix once and for all the choice of apices  $P_\tau$  and preferred bases  $b_\tau$

- for all  $B_1$ -facets  $\tau$ , and
- for all  $B_1$ -facets  $\tau$  of  $B_2$ -facets (by a  $B_1$ -facet of a facet  $\sigma$  we mean a codimension 2 face of the Newton polyhedron that belongs to  $\sigma$  and is a  $B_1$ -pyramide).

Moreover, in the setting of Theorem 4.3, we can choose the apices and the preferred bases consistently, so that the preferred base of every aforementioned face  $\tau$  belongs to the base direction of every  $B$ -facet containing  $\tau$ . This in particular, ensures that

- every two  $B_1$ -facets intersecting by a codimension 2  $B_1$ -face have the same apex and preferred base, and
- if a codimension 2  $B_1$ -face  $\tau'$  is at the same time a facet of a  $B_1$ -facet  $\tau$  and of a  $B_2$ -facet (so that we have chosen the preferred base and apex for it), then  $\tau'$  and  $\tau$  have the same apex and preferred base.

**Definition 4.11.** Let  $L$  be the 2-dimensional coordinate plane along which the projection of a  $B_2$ -facet  $\tau$  equals the standard simplex, and let  $l \subset \mathbb{R}^n$  be the 1-dimensional vector space parallel to the intersection of  $L$  with the affine span of  $\tau$ .

The dual hyperplane to  $l$  will be denoted by  $D_\tau = D_{\tau^\circ}$  and called the delimiter of  $\tau$ .

In particular, if  $n = 4$ , then, in the notation of Figure 2, the delimiter is the 3-dimensional plane normal to the three parallel segments.

**Remark 4.12.** Most  $B_2$ -facets have a unique  $V$ -edge normal to the delimiter. However, we allow this edge to degenerate into a vertex (such  $B_2$ -facets are said to be degenerate).

However, various attributes of this  $V$ -edge make natural sense even for degenerate  $B_2$ -facets. For instance, “the length of the  $V$ -edge of  $\tau$ ” and “the dual cone of the  $V$ -edge of  $\tau$ ” refer to 0 and  $D_\tau \cap V^\circ$  respectively for a degenerate  $B_2$ -facet  $\tau$  with a vertex  $V$  instead of the  $V$ -edge. In what follows, this small abuse of terminology never causes confusion.

**Remark 4.13.** Every  $B_2$ -facet (including the degenerate ones) has three distinguished facets (i.e.  $(n - 2)$ -dimensional faces): namely, it has exactly one non-simplicial non- $V$ -facet and exactly two  $B_1$ -facets.

**Definition 4.14.** 1. We shall say that a vertex  $P$  of a  $B_2$ -facet  $\tau$  and a number  $i \in \{1, \dots, n\}$  are an apex and a preferred base of  $\tau$  on the side of a face  $F \prec \tau$ , if  $P$  and  $i$  are the apex and the preferred base of a  $B_1$ -facet  $\sigma$  of  $\tau$ , such that  $\sigma \cap F \neq \emptyset$ . If  $\tau$  has a unique preferred base on the side of  $F$  (that is, if  $\sigma$  is uniquely defined by the condition  $\sigma \cap F \neq \emptyset$ ), this preferred base will be denoted by  $b_\tau^F = b_{\tau^\circ}^F$ .

2. Similarly, if a cone  $C$  in the star of  $\tau^\circ$  does not intersect the delimiter  $D_\tau$ , then the apex and the preferred base of the  $B_2$ -facet  $\tau$  on the side of  $C$  are defined as the apex and the preferred base of the  $B_1$ -facet of  $\tau$ , whose dual cone is not separated from  $C$  by the delimiter.

3. For conformity, we shall say that a vertex  $P$  of a  $B_1$ -facet  $\tau$  and a number  $i$  are its apex and preferred base on the side of a face  $F \prec \tau$  (or a cone  $C$  in the star of  $\tau^\circ$ ), if  $P$  and  $i$  are the apex and the preferred base of  $\tau$  (independently of  $F$  and  $C$ ).

In particular, if  $n = 4$ , then, in the notation of Figure 2, the vertex  $P$  is on the side of all faces of  $\tau$  except for the faces of  $QBY$ . Notice the following trivial observation.

**Lemma 4.15.** *If a face  $F$  is contained in a  $B$ -facet  $\tau$  and not contained in a coordinate hyperplane, then every apex of  $\tau$  on the side of  $F$  is contained in  $F$ .*

## 4.5 Sprouts and cancellation of contributions

Let  $C$  be a (not necessarily convex) polyhedral cone in the star of  $\tau^\circ$ , where  $\tau$  is a compact  $B$ -facet. In the case of a  $B_2$ -facet, we additionally assume that  $C$  is not intersected by the delimiter  $D_\tau$ . Let  $P$  and  $i$  be the apex and the preferred base of  $\tau$  on the side of  $C$ . Recall that the standard basis in  $\mathbb{R}^n$  is denoted by  $e_1, \dots, e_n$ .

**Definition 4.16.** The union of  $C \cap \text{rel.int} P^\circ$  and all two-dimensional cones, generated by  $e_i$  and a point of  $C \cap \partial P^\circ$ , is denoted by  $S_{C,\tau}$  and is called the sprout of  $C$ . The vertex  $P$  is denoted by  $R_{C,\tau}$  and is called the root of  $C$ .

**Remark 4.17.** By the definition, if  $C$  is a closed  $n$ -dimensional polyhedral cone (in the sense of Definition 3.5), containing  $\tau^\circ$  in its interior, then so is the sprout. On the other hand, convexity of  $C$  does not imply convexity of the sprout, see the picture below.

**Example 4.18.** Figure 5 gives the simplest example of a cone  $C$  and its sprout in the star of the ray  $\tau^\circ$ .

**Lemma 4.19.** *The contribution of a sprout is simple:*

$$Z(S_{C,\tau}) = \int_{S_{C,\tau}} \exp(-\langle u, R_{C,\tau} \rangle s - \langle u, \mathbf{1} \rangle) du.$$




Figure 5: the sprout of the cone  $C$

*Proof.* The contribution of  $C \cap \text{rel.int}P^\circ$  equals the integral of  $\exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle)$  by definition. Triangulating the set  $C \cap \partial P^\circ$  and, for every simplicial cone  $T$  of this triangulation, applying Lemma 3.3 to the contribution of the cone generated by  $T$  and  $e_i$ , we conclude that the rest of  $S_{C,\tau}$  also contributes the integral of  $\exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle)$ .  $\square$

**Lemma 4.20.** *Let  $\tau$  be a  $B_2$ -facet, and assume without loss of generality that the projection of  $\tau$  along the  $(e_{n-1}, e_n)$ -coordinate plane is the  $(n-2)$ -dimensional standard simplex. Let  $r$  be a ray in the dual cone to the non-simplicial non- $V$ -facet of  $\tau$ . Then the contribution of the  $(n-1)$ -dimensional (relatively open) cone  $S_{r,\tau}$  generated by  $r, e_1, \dots, e_{n-2}$  (which we call the delimiter sprout of  $r$ ) is equal to 0.*

*Proof.* For  $i \in \{1, \dots, n-2\}$  we denote the length of the edge of  $\tau$  in the preimage of  $e_i$  by  $g_i$ . For a subset  $I \subset \{1, \dots, n-2\}$ , denote by  $C_I$  the cone generated by  $\tau^\circ$  and  $e_i$  ( $i \in I$ ), and by  $\tilde{C}_I$  the one spanned by  $C_I$  and  $r$ . Then the contribution of the delimiter sprout splits into those of the cones  $C_I$  for  $I \subset \{1, \dots, n-2\}$  and the cones  $\tilde{C}_I$  for  $I \subsetneq \{1, \dots, n-2\}$ . Note that the normalized volume of the non-simplicial non- $V$ -facet  $\tau_0$  of  $\tau$  is equal to  $\sum_{i=1}^{n-2} g_i$ . We have also a similar description for the normalized volume of any face of  $\tau_0$ . Then the sum of the contributions of the cones  $\tilde{C}_I$  over  $I \subsetneq \{1, \dots, n-2\}$  is equal to 0 (see Lemma 3.3 for a similar calculation). Similarly we can show that the corresponding sum for the cones  $C_I$  over  $I \subset \{1, \dots, n-2\}$  is 0.  $\square$

**Lemma 4.21.** *The edges of a sprout or a delimiter sprout  $S_{C,\tau}$  are either edges of the cone  $(\partial C) \cap \sigma^\circ$  for some face  $\sigma \prec \tau$ , or coordinate rays.*

This directly follows from the definition of a sprout.

## 4.6 Critical cones

We now introduce and study the key notion of critical faces for a candidate pole  $s_0 \neq -1$  of the zeta function  $Z_{\text{top},f}(s)$ . Similarly to Lemma 3.4, we have the following lemma.

**Lemma 4.22.** *For  $s_0 \neq -1$  and any vertex  $P$  of a  $B$ -facet with a base hyperplane  $\{x_i = 0\}$ , the equation  $\langle u, P \rangle s_0 + \langle u, \mathbf{1} \rangle = 0$  is not satisfied by the coordinate vector  $u = e_i$ . In particular, this equation is non-trivial and thus defines a hyperplane  $L_P$  in  $\mathbb{R}^n$ .*

**Definition 4.23.** For a face  $F$ , we define the critical set  $K_F \subset F^\circ$  in  $F^\circ$  to be the closure of  $\cup_P(\text{rel.int}F^\circ \cap L_P)$ , where in the union  $\cup_P$  the vertex  $P \prec \Gamma_+(f)$  ranges through the apices on the side of  $F$  for  $B$ -facets containing  $F$  (see Definition 4.14).

**Lemma 4.24.** 1. If a face  $F$  is contained in a coordinate plane, then  $K_F$  is a finite union of hyperplanes in  $F^\circ$ .

2. If a face  $F$  is not contained in a hyperplane, then  $K_F$  is given by one equation  $\langle u, Q \rangle_{s_0} + \langle u, \mathbf{1} \rangle = 0$ , where  $Q$  is an arbitrary point of  $F$ . In particular, either  $K_F$  is a hyperplane in  $F^\circ$ , or  $K_F = F^\circ$ .

*Proof.* 1) If  $F$  is in the coordinate plane  $x_i = 0$ , then  $e_i \in F^\circ$ . If  $F$  is in a  $B$ -facet  $\tau$ , then the set  $L_P$  does not contain  $e_i \in F^\circ$  by Lemma 4.22, so no  $L_P$  can contain the whole  $F^\circ$ .

2) If  $F$  is not in a coordinate plane, then any  $P$  in the definition of the critical set is contained in  $F$  by Lemma 4.15, so  $K_F$  is given by the equation  $\langle u, P \rangle_{s_0} + \langle u, \mathbf{1} \rangle = 0$ . The set defined by this equation in  $F^\circ$  does not change if we substitute  $P$  with any other point  $Q \in F$ , because  $\langle Q - P, u \rangle = 0$  for  $u \in F^\circ$ .  $\square$

**Definition 4.25.** We say that a face  $F \prec \Gamma_+(f)$  and its dual cone  $F^\circ \in \Sigma_0$  are critical (for the candidate pole  $s_0$ ) if  $K_F = F^\circ$ .

**Lemma 4.26.** A critical face is not contained in a coordinate plane. Equivalently, a critical cone is disjoint from coordinate planes.

This follows from Lemma 4.24(1).

**Proposition 4.27.** Assume that a candidate pole  $s_0$  is contributed only by  $B$ -facets. In this case, the following conditions are equivalent:

1. The face  $F$  is critical, that is,  $F$  is contained in a  $B$ -facet, and, for its apex  $P$  on the side of  $F$ , the plane  $L_P$  contains the cone  $F^\circ$ .
2. Every facet containing  $F$  is a  $B$ -facet, and, for every apex  $P$  on the side of  $F$ , the plane  $L_P$  contains the cone  $F^\circ$ .
3. Every facet containing  $F$  contributes the pole  $s_0$  to the topological  $\zeta$ -function.

The second condition will be mostly used in practice, and the third one is especially simple (and relates the first two).

*Proof.* Condition (3) follows from (1), because, for every facet  $\tau \supset F$  and its normal covector  $u$  and a point  $Q \in F$  we have  $\langle u, Q \rangle_{s_0} + \langle u, \mathbf{1} \rangle = 0$  by Lemma 4.24(2), which is applicable because  $F$  is not in a hyperplane by Lemma 4.26. Thus  $\tau$  contributes the pole  $s_0$ .

Condition (2) follows from (3), because every contributing  $\tau$  is a  $B$ -facet by the assumptions of the proposition, and the face  $F \subset \tau$  contains an apex of  $\tau$  on the side of  $F$  by Lemma 4.15, thus the equality

$$\langle u, P \rangle_{s_0} + \langle u, \mathbf{1} \rangle = 0$$

holds for the normal covector  $u$  to every facet  $\tau' \supset F \ni P$ , i.e. for the generating vector of every edge of  $F^\circ$ , thus it holds also for every  $u \in F^\circ$ .

Finally, Condition (1) follows from (2) tautologically.  $\square$

**Corollary 4.28.** A face of a critical cone in the dual fan  $\Sigma_0$  is a critical cone. Equivalently, a face containing a critical face is critical itself.

Note that, however, for non-critical faces  $F \subset G$ , it is not in general true that  $K_G \cap F^\circ \subset K_F$ .

It is now a crucial observation that, under the assumptions of Theorem 4.3, every critical face  $F$  is a  $B_1$ -pyramid (by Lemma 4.7 and Proposition 4.27), and all  $B$ -facets, containing  $F$ , have the same preferred base  $b_F$  and the same apex  $P_F$  on the side of  $F$ .

## 4.7 A tubular neighborhood of the critical subfan

We are now ready to prove Theorem 4.3. Recall that by this time we have

- chosen once and for all a preferred base of every  $B_1$ -facet and every  $B_1$ -facet of every  $B_2$ -facet in the Newton polyhedron  $\Gamma_+(f)$ ,
- depending on this choice, called some cones critical in the dual fan  $\Sigma_0$ ,
- defined the delimiter hyperplane  $D_\sigma$  for every cone  $\sigma$ , dual to a  $B_2$ -facet (see Definition 4.11).

Choose once and for all an affine structure on the projectivization  $\mathbb{P}\mathbb{R}_+^n$ . For every cone  $C$  and fan  $\Sigma$ , denote their projectivizations by  $\mathbb{P}C$  and  $\mathbb{P}\Sigma$  respectively. In this subsection, we refer to cones and their projectivizations interchangeably whenever it causes no confusion.

According to Corollary 4.28, the set of projectivized critical cones in the dual fan  $\Sigma_0$  is a polyhedral complex  $\Sigma_c$ , closed with respect to taking faces and intersections and disjoint from the boundary of  $\mathbb{P}\mathbb{R}_+^n$  by Lemma 4.26. We shall construct a generic piecewise linear tubular neighborhood of  $\Sigma_c$  in  $\mathbb{P}\mathbb{R}_+^n$ , whose boundary is transversal to the projectivized critical sets of non-critical cones.

Recall that two piecewise linear sets in  $\mathbb{R}^n$  are said to be transversal in  $U \subset \mathbb{R}^n$ , if they can be triangulated so that the relatively open simplices from these two triangulations are pairwise transversal at every intersection point in  $U$ .

Recall that the corner locus of a continuous piecewise linear function is the (piecewise linear) set of all points at which the function is not smooth.

For every projectivized closed cone  $\sigma \in \Sigma_c$ , choose a convex piecewise linear (in the sense of the selected affine structure) “distance function”  $d_\sigma : \mathbb{P}\mathbb{R}_+^n \rightarrow \mathbb{R}_+$ , satisfying the following properties:

- 1) It vanishes on  $\sigma$  and is strictly positive outside of it.
- 2) For every non-critical cone  $\sigma' \in \Sigma_0$ , the corner locus of  $d_\sigma$  is transversal to the projectivized critical set  $\mathbb{P}K_{\sigma'}$  in the complement to  $\sigma$ .
- 3) Moreover, for every non-critical cone  $\sigma' \in \Sigma_0$  and its edge  $\sigma''$  dual to a  $B_2$ -facet, the corner locus of  $d_\sigma$  is transversal to the projectivized delimiter  $\mathbb{P}D_{\sigma''} \cap \sigma'$  and the critical set  $\mathbb{P}D_{\sigma''} \cap K_{\sigma'}$  in the complement to  $\sigma$ .

In the sense of this distance, we shall consider  $\varepsilon$ -neighborhoods  $B_\sigma(\varepsilon) = \{u \mid d_\sigma(u) < \varepsilon\}$  and their boundaries  $S_\sigma(\varepsilon)$ , which are all piecewise-linear sets.

Associating positive numbers  $\varepsilon_\sigma$  to all  $\sigma \in \Sigma_c$ , introduce the following sets:

- the open neighborhood  $U = \bigcup_{\sigma \in \Sigma_c} B_\sigma(\varepsilon_\sigma)$  of the critical complex  $\Sigma_c$ ;
- for every  $\sigma \in \Sigma_c$ , the set  $U_\sigma = B_\sigma(\varepsilon_\sigma) \setminus \bigcup_{\sigma' \subsetneq \sigma} B_{\sigma'}(\varepsilon_{\sigma'})$ ;
- for every  $\sigma \in \Sigma_c$  dual to a  $B_2$ -facet, the *delimiter disk*  $DD_\sigma = U_\sigma \cap \mathbb{P}D_\sigma$ .

**Lemma 4.29.** *One can choose the numbers  $\varepsilon_\sigma$  so that*

1. *for every non-critical projectivized cone  $\sigma' \in \mathbb{P}\Sigma_0$ , no vertex of the boundary of  $U_\sigma \cap \sigma'$  and  $DD_\sigma \cap \sigma'$  is contained in the critical set  $\mathbb{P}K_{\sigma'} \cap \text{rel.int}\sigma'$ ;*

2.  $U_\sigma$  is contained in the star of  $\sigma$ ;
3.  $U_\sigma \cap U_\delta = \emptyset$  unless  $\sigma = \delta$ , and  $\bar{U}_\sigma \cap \bar{U}_\delta = \emptyset$  unless  $\sigma \subset \delta$  or vice versa;
4.  $DD_\sigma$  divides  $U_\sigma$  into two connected components.

*Proof.* All of these properties are satisfied if the tuple  $(\varepsilon_\sigma)$  is chosen generically (i.e. avoiding finitely many hyperplanes in the space of all such tuples) and rapidly decreasing (i.e.  $\varepsilon_\sigma \ll \varepsilon_\delta \ll 1$  for all  $\delta \subset \sigma$ ).  $\square$

**Definition 4.30.** Under the assumptions of Theorem 4.3, the *vertex function*  $v : U \setminus (\text{delimiter disks}) \rightarrow (\text{vertices of } \Gamma_+(f))$  is defined on every  $U_\sigma$  as follows:

- if  $\sigma \in \Sigma_c$  is dual to a non- $B_2$  critical face  $F$ , then we define  $v(\cdot)$  on  $U_\sigma$  as the apex  $P_F$ .
- if  $\sigma \in \Sigma_c$  is dual to a  $B_2$ -facet  $\tau$ , then we define  $v(\cdot)$  on each of the two components of  $U_\sigma \setminus DD_\sigma$  as the apex of  $\tau$  on the side of this component.

Note that the vertex function is locally constant on its domain. We now use this observation to consistently substitute every piece of the neighborhood  $U$  by an appropriate sprout. Recall that we refer to cones and their projectivizations interchangeably, and, in particular,  $S_{W,\tau}$  for the projectivization  $W$  of a cone  $C$  is another notation for the sprout  $S_{C,\tau}$ .

- If  $\sigma \in \Sigma_c$  is dual to a face  $F$  that is not a  $B_2$ -facet, then the vertex function  $v$  equals a constant  $P$  on it, so we define  $V_{\sigma,0}$  as the sprout  $S_{U_\sigma,\tau}$  (see Definition 4.16) for any  $B$ -facet  $\tau \supset F$ . Note that neither the sprout  $V_{\sigma,0} := S_{U_\sigma,\tau}$  nor its root  $R_{\sigma,0} := R_{U_\sigma,\tau} = P$  depend on the choice of  $\tau$ .

- If  $\sigma \in \Sigma_c$  is dual to a  $B_2$ -facet  $\tau$  with the apices  $P_{\pm 1}$ , then we define  $V_{\sigma,\pm 1}$  as the sprout  $S_{\{v=P_{\pm 1}\} \cap U_\sigma,\tau}$  and the root  $R_{\sigma,\pm 1}$  as  $R_{\{v=P_{\pm 1}\} \cap U_\sigma,\tau} = P_{\pm 1}$ . Also, choosing a ray  $r$  in the dual cone  $\sigma'$  of the non-simplicial non- $V$ -facet of  $\tau$  outside of  $K_{\sigma'}$ , we define  $V_{\sigma,0}$  as  $S_{r,\tau}$  (see Lemma 4.20), leaving  $R_{\sigma,0}$  undefined.

Define the *sprouting*  $S_P \subset \mathbb{R}^n$  of a vertex  $P$  as the union of all  $V_{\sigma,*}$  ( $* = 0, \pm 1$ ), such that  $R_{\sigma,i} = P$ .

**Lemma 4.31.** 1. The contribution  $Z(S_P)(s)$  of the sprouting  $S_P$  to the zeta function  $Z_{\text{top},f}(s)$  is equal to

$$\int_{S_P} \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du$$

modulo a function that has no pole at  $s_0$ .

2. No edge of the boundary of  $S_P$  is critical for  $(s_0, P)$ .

*Proof.* The part (1) follows from Lemma 4.19. To deduce (2), it is enough (by Lemma 4.21) to show that every vertex of  $(\partial U_\sigma) \cap \sigma'$  and  $(\partial DD_\sigma) \cap \sigma'$  for every projectivized cone  $\sigma' \supset \sigma$  is either not in  $\mathbb{P}K_{\sigma'}$ , or not a projectivized edge of  $\partial S_P$ . For non-critical  $\sigma'$ , this follows from Lemma 4.29(1), and every critical  $\sigma'$  is either in the projectivized interior of  $S_P$ , or disjoint from its closure, or intersects its boundary at an interior point of its facet  $DD_{\sigma'}$  (thus no vertex in  $\sigma'$  can be a projectivized edge of  $\partial S_P$ ).  $\square$

*Proof of Theorem 4.3.* By the preceding lemma and Lemma 3.6, the contribution of every  $S_P$  to the topological zeta function of  $f$  has no pole at  $s_0$ . By Lemma 4.20, the same is true for  $V_{\sigma,0}$  for every  $B_2$ -facet  $\sigma$ . Since

$$V = \bigcup_{\sigma, * = 0, \pm 1} V_{\sigma,*}$$

contains all the critical cones in its interior, for every cone  $\sigma' \in \Sigma_0$  the set  $\sigma' \setminus V$  has edges of two types: either edges of  $(\partial V) \cap \sigma'$  or non-critical 1-dimensional cones of  $\Sigma_0$ . The edges of the first kind are not in the critical set of any cone by Lemma 4.31(2), and for the second kind the same holds by definition. Thus the contribution of  $\sigma' \setminus V$  to the topological zeta function has no pole at  $s_0$  as well by Lemma 3.6. We have subdivided  $\mathbb{R}_+^n$  into the pieces

- $S_P$  for some vertices  $P$ ,
- $V_{\tau^c,0}$  for some  $B_2$ -facets  $\tau$ ,
- $\sigma' \setminus V$  for some non-critical cones  $\sigma'$ ,

so that none of them contributes to the pole  $s_0$ . □

## 5 Fake poles of the topological zeta function in dimension 4

Throughout this section we work in dimension  $n = 4$ . In the first subsection, we classify all non-contributing configurations of  $B$ -faces. The proof of the classification theorem occupies the subsequent two subsections and makes use of the tools introduced in the preceding two sections. In the last subsection, we show that every non- $B$ -facet contains a non- $B$ -simplex and discuss a possible general definition of  $B$ -facets in arbitrary dimension. Both of the mentioned results will be used in the last section to prove the monodromy conjecture for all non-degenerate singularities of functions of 4 variables.

### 5.1 The main theorem

We shall prove that if a candidate pole of the topological zeta function  $Z_{\text{top},f}(s)$  is contributed only by  $B$ -facets, then it is fake, with one exception:

**Definition 5.1.** A border is a triangular face of  $\Gamma_+(f)$ , such that, up to a reordering of the coordinates, its vertices are of the form  $A = (1, 1, *, *)$ ,  $B = (0, 0, *, *)$  and  $C = (0, 0, *, *)$ , and the two facets containing it are  $B$ -facets with the vertex  $A$  and the bases in the coordinate hyperplanes  $\{x_1 = 0\}$  and  $\{x_2 = 0\}$  respectively.

The border  $ABC$  is said to be a  $B$ -border unless (up to reordering  $B$  and  $C$  and the last two coordinates) we have  $A = (1, 1, 0, a)$ ,  $B = (0, 0, 0, b)$  and  $C = (0, 0, 1, c)$ . In the latter case (implying, in particular, that the edge  $BC$  is itself a  $B$ -edge in the coordinate plane  $O_{34}$ ) the border  $ABC$  is said to be a  $B^2$ -border (see Figure 6). The vertex  $A$  is called the apex of the border, and the first two coordinates are called its bases.

Let  $f$  be a non-degenerate polynomial on  $\mathbb{C}^4$  with the Newton polyhedron  $\Gamma_+(f) \subset \mathbb{R}^4$  and  $\Sigma_0$  its dual fan.




Figure 6: a  $B^2$ -border and a  $B$ -border

**Theorem 5.2.** *Assume that  $f$  does not have a Morse singularity at the origin  $0 \in \mathbb{C}^4$ , that a candidate pole  $s_0 \neq -1$  of the topological zeta function  $Z_{\text{top},f}(s)$  is contributed only by  $B$ -facets, and that no two of them contain the same  $B$ -border (although they may contain the same  $B^2$ -border). Then  $s_0$  is fake i.e. not an actual pole of  $Z_{\text{top},f}(s)$ .*

The rest of this subsection is devoted to the proof of this theorem.

The first difference with the setting of Theorem 4.3 is that we have to prove that  $B^2$ -borders do not contribute to candidate poles. For this purpose, we shall extend the notion of a sprout (Definition 4.16) to them. Let  $C$  be a (not necessarily convex) cone in the star of  $\tau^\circ$ , where  $\tau$  is a border with bases  $i$  and  $j$  and the apex  $P$ .

**Definition 5.3.** The cone  $C$  is said to be border-convex, if every 3-dimensional plane through  $e_i$  and  $e_j$  intersects  $C$  by a convex cone.

**Definition 5.4.** The union of  $C \cap \text{rel.int}P^\circ$  and all three-dimensional cones, generated by the coordinate vectors  $e_i, e_j$  and  $u \in C \cap \partial P^\circ$ , is denoted by  $S_{C,\tau}$  and is called the border sprout of  $C$ . The vertex  $P$  is denoted by  $R_{C,\tau}$  and is called the root of  $C$ .

The definition implies the following property.

**Lemma 5.5.** *The edges of a border sprout of a cone  $C$  are either edges of  $(\partial C) \cap \sigma'$  for some  $\sigma' \supset \sigma$ , or coordinate rays.*

**Lemma 5.6.** *Let  $C$  be a border-convex cone in the star of the dual cone to a  $B^2$ -border  $\tau$ , and assume that both of its adjacent  $B_1$ -facets contribute to the same candidate pole  $s_0$ .*

*Then the contribution of the border sprout  $Z(S_{C,\tau})$  is equal to*

$$\int_{S_{C,\tau}} \exp(-\langle u, R_{C,\tau} \rangle s - \langle u, \mathbf{1} \rangle) du$$

*modulo a function that has no pole at  $s = s_0$ .*

*Proof.* Let  $ABD$  be a  $B^2$ -border with coordinates  $A = (1, 1, 0, a)$ ,  $B = (0, 0, 0, b)$  and  $D = (0, 0, 1, d)$ . The fact that the adjacent  $B_1$ -facets  $\tau_1$  and  $\tau_2$  contribute the same candidate pole means that the line  $s \cdot A + \mathbf{1}$  intersects the vector spans of  $\tau_1 - A$  and  $\tau_2 - A$  at the same point  $s_0 \cdot A + \mathbf{1}$ . Thus this point is a linear combination of the vectors  $DB$  and  $AD$ :

$$DB = \beta(s_0 \cdot A + \mathbf{1}) + \alpha \cdot AD.$$

Solving this system of equations for  $\alpha, \beta$  and  $s_0$ , we find

$$s_0 = (a + d - 2b - 1)/b, \quad \alpha = -s_0 - 1, \quad \beta = 1.$$

At the same time, for the future reference, we interpret the latter equalities as the computation of the ratio of  $DB$  and  $s_0 \cdot A + \mathbf{1}$  as linear functions on the dual space  $(\mathbb{R}^4)^*$  restricted on the hyperplanes  $AD^\perp$  and  $AB^\perp$ :

$$\frac{DB|_{AB^\perp}}{(s_0 \cdot A + \mathbf{1})|_{AB^\perp}} = \beta = 1, \quad \frac{DB|_{AD^\perp}}{(s_0 \cdot A + \mathbf{1})|_{AD^\perp}} = \beta/(1 - \alpha) = 1/(2 + s_0). \quad (*)$$

We now first prove the lemma for a very special choice of the cone  $C$ .

Choose vectors  $u$ ,  $u_B$  and  $u_D$  in the interior of the cones  $ABD^\circ$ ,  $AB^\circ$  and  $BD^\circ$  respectively. Recall that the standard basis is denoted by  $e_1, e_2, e_3, e_4$ . Denote the closure of the union of the simplicial cones  $\langle u, u_B, e_1, e_2 \rangle$  and  $\langle u, u_D, e_1, e_2 \rangle$  by  $N_{ABD}$ .

Then the statement of the lemma is valid for  $C = N_{ABD}$ . To prove this, represent  $Z(N_{ABD})(s) - \int_{N_{ABD}} \exp(-s\langle v, A \rangle - \langle v, \mathbf{1} \rangle) dv$  by Lemma 2.2 as

$$\begin{aligned} & \frac{|\det(e_1, e_2, u, u_D)|}{(s \cdot A + \mathbf{1})|_u \cdot (s \cdot A + \mathbf{1})|_{u_D}} + \frac{|\det(e_1, e_2, u, u_B)|}{(s \cdot A + \mathbf{1})|_u \cdot (s \cdot A + \mathbf{1})|_{u_B}} - \frac{s \cdot |e_1 \wedge e_2 \wedge u|}{(s+1) \cdot (s \cdot A + \mathbf{1})|_u} \\ & \frac{|\det(e_1, e_2, u, u_D)|}{(s+1)^2 \cdot (s \cdot A + \mathbf{1})|_u \cdot (s \cdot A + \mathbf{1})|_{u_D}} - \frac{|\det(e_1, e_2, u, u_B)|}{(s+1)^2 \cdot (s \cdot A + \mathbf{1})|_u \cdot (s \cdot A + \mathbf{1})|_{u_B}}, \quad (**) \end{aligned}$$

where  $|\cdot|$  in the third term stands for the lattice length. Collecting similar terms and taking into account  $u \cdot DB = 0$  (which in coordinates  $v = (v_1, v_2, v_3, v_4)$  reads as  $v_3 = v_4 \cdot (b-d)$ ), we can rewrite  $(**)$  as the uninteresting factor  $\frac{s \cdot v_4}{(s+1)^2 \cdot (s \cdot A + \mathbf{1})|_u}$  times

$$\frac{(s+2)|DB|_{u_D}|}{(s \cdot A + \mathbf{1})|_{u_D}} + \frac{(s+2)|DB|_{u_B}|}{(s \cdot A + \mathbf{1})|_{u_B}} - (s+1) \quad (*)$$

Since  $u_B$  and  $u_D$  are chosen to be support vectors of the edges  $AB$  and  $AD$  respectively, we have  $u_B \cdot A = u_B \cdot B < u_B \cdot D$  and  $u_D \cdot A = u_D \cdot D < u_D \cdot B$ , so  $|DB|_{u_B}| = DB|_{u_B}$  and  $|DB|_{u_D}| = -DB|_{u_D}$ . Applying this and  $(*)$ , we conclude that  $(*)$  for  $s = s_0$  equals  $-(s+2)/(s+2) + (s+2) - (s+1) = 0$ .

The border sprout  $S_{C,\tau}$  of an arbitrary border-convex  $C$  can be represented as the union (with disjoint interiors) of  $N_{ABC}$  and the sprouts of the form  $S_{C^{(k)},\tau_k}$ ,  $k = 1, 2$ , for certain cones  $C^{(k)}$ . Thus the statement of the lemma is valid for it, applying the preceding computation to  $N_{ABC}$  and Lemma 4.19 to  $S_{C^{(k)},\tau_k}$ .  $\square$

**Remark 5.7.** Most of borders do not satisfy this property.

## 5.2 Very critical cones

In contrast to the setting of Theorem 4.3, we cannot in general choose preferred bases and apices of  $B_1$ -faces consistently, so we now choose them arbitrarily once and for all in this section. Recall that we fix the choice of apices  $P_\tau$  and preferred bases  $b_\tau$  for all  $B_1$ -facets  $\tau$  and for all  $B_1$ -facets of  $B_2$ -facets  $\tau$ , and then use them to define preferred bases and apices of  $B_2$ -facets on the side of a given face or cone, see Subsection 4.4 for details.

As a result, in contrast to the setting of Theorem 4.3, different  $B_1$ -facets, containing a given critical face, may have different preferred bases. Such critical faces are said to be very critical. They are studied in this subsection.

**Definition 5.8.** The preferred base  $b_F = b_{F^\circ} \subset \{1, 2, 3, 4\}$  of a critical face  $F \prec \Gamma_+(f)$  and its dual cone  $F^\circ \in \Sigma_0$  is the set of the preferred bases on the side of  $F$  for all  $B$ -facets containing  $F$ . We say that the face  $F$  and its dual cone  $F^\circ \in \Sigma_0$  are very critical if  $|b_F| \geq 2$ .

**Example 5.9.** The projectivized pictures of the Newton polyhedron  $\Gamma_+(f)$  on Figure 7 below give some examples of (2-dimensional) very critical faces (hatched), provided that all the facets on the pictures are  $B$ -facets contributing the same candidate pole  $s_0$ . The apices  $B_1$ -faces are bold points, and the apices of the  $B_2$ -facet are the end points of the bold segment.




Figure 7: very critical faces

This subsection is devoted to the study of very critical faces depending on their dimension. First of all, by Lemma 4.22 there is no critical vertex, and a facet  $\tau$  is critical if and only if the candidate pole  $s_0$  is contributed by  $\tau$  (and thus  $\tau$  is a  $B$ -facet in the assumptions of Theorem 5.2). A 2-dimensional face  $F \prec \Gamma_+(f)$  is critical if and only if it separates two  $B$ -facets  $\tau_1$  and  $\tau_2$ , contributing  $s_0$ . This critical face  $F$  is very critical, if moreover we have  $b_{\tau_1}^F \neq b_{\tau_2}^F$ .

**Definition 5.10.** Let  $F$  be a 2-dimensional very critical face separating two  $B$ -facets  $\tau_1$  and  $\tau_2$ . Then we define the  $F$ -delimiter  $D_F = D_{F^\circ} \subset \mathbb{R}^4$  to be the vector subspace in  $\mathbb{R}^4$  generated by the  $b_{\tau_1}^F$  and  $b_{\tau_2}^F$ 'th coordinate lines in  $\mathbb{R}^4$  and a ray  $R \subset \text{rel.int}F^\circ$ .

The delimiter is said to be generic, if, for every non-critical face  $G \subset F$ , the critical set  $D_F \cap K_G$  is nowhere dense in  $D_F \cap \text{rel.int}G^\circ$ .

**Lemma 5.11.** *Under the assumptions of Theorem 5.2, any 2-dimensional very critical face  $F$  is a triangle. If it is a border, then the delimiter  $D_F$  is the affine span of the dual cone of the  $V$ -edge of  $F$ . Otherwise the affine span  $\text{aff}(F^\circ)$  of  $F^\circ$  is transversal to the delimiter  $D_F$ . In particular, in both cases  $D_F$  is a hyperplane in  $\mathbb{R}^4$ .*

*Proof.* The face  $F$  is a triangle by Lemma 4.7. Assume that  $\text{aff}(F^\circ)$  is not transversal to the delimiter. Then for the coordinate plane  $H = \{x_{b_{\tau_1}^F} = x_{b_{\tau_2}^F} = 0\} \subset \mathbb{R}^4$  we have  $\dim\{\text{aff}(F) \cap H\} \geq 1$ . This implies that  $\text{aff}(F)$  is not transversal to  $H$ , and hence their intersection is a  $V$ -edge in both  $\tau_1$  and  $\tau_2$ . Then  $F$  is a border.  $\square$

**Corollary 5.12.** *All but finitely many possible delimiters of a non-border very critical face are generic.*

An edge  $F \prec \Gamma_+(f)$  is critical, if and only if all of the facets containing it are  $B$ -facets  $\tau$  contributing to the candidate pole  $s_0$ . Since we have  $\dim F^\circ = 3$  in this case, moreover, for any apex  $P$  of such  $\tau$  on the side of  $F$ , the critical hyperplane  $L_P$  coincides with  $\text{aff}(F^\circ)$ .

**Proposition 5.13.** *Under the assumption of Theorem 5.2, no edge of  $\Gamma_+(f)$  is very critical.*

*Proof.* Assume that an edge  $P_1P_2$  is very critical. Then, by Proposition 4.27(2), there exist  $B$ -facets  $\tau_1$  and  $\tau_2$  containing it with the apices  $Q_1$  and  $Q_2$  on the side of  $P_1P_2$ , and their critical hyperplanes  $L_{Q_i} : \langle u, Q_i \rangle_{s_0} + \langle u, \mathbf{1} \rangle = 0$

both coincide with the hyperplane  $\langle u, P_1 - P_2 \rangle = 0$  generated by the dual cone  $(P_1P_2)^\circ$ . Moreover, since  $P_1P_2$  is not in a coordinate plane, we have  $P_1 = Q_1$  and  $P_2 = Q_2$  or vice versa. Thus the vectors  $P_1s_0 + \mathbf{1}$  and  $P_2s_0 + \mathbf{1}$  are parallel.

On the other hand, it cannot happen that one of the points  $P_1$  and  $P_2$  is in the apex of  $\tau_1$  and in the base of  $\tau_2$ , and the other one is in the apex of  $\tau_2$  and in the base of  $\tau_1$ . Otherwise, reordering coordinates if necessary, we would have  $P_1 = (1, 0, *, *)$ ,  $P_2 = (0, 1, *, *)$ ,  $P_1s_0 + \mathbf{1} = (1 + s_0, 1, *, *)$ ,  $P_2s_0 + \mathbf{1} = (1, 1 + s_0, *, *)$ . Since the last two vectors are parallel and  $s_0 \neq 0$ , we have  $s_0 = -2$ ,  $P_1s_0 + \mathbf{1} = -(P_2s_0 + \mathbf{1})$  and hence  $P_1 + P_2 = \mathbf{1}$ . Thus, up to reordering the coordinates and  $P_i$ 's, we have  $P_1 = (1, 0, 0, 0)$  (i.e.  $f$  has no singularity at the origin) or  $P_1 = (1, 0, 1, 0)$  and  $P_2 = (0, 1, 0, 1)$  and hence  $f$  has a Morse singularity at the origin by Lemma 4.8.

If one of  $P_i$ 's is in the apex of all  $B$ -facets  $\tau_j$ 's containing  $P_1P_2$ , and one of  $\tau_j$ 's is  $B_2$ , then the other facet containing its quadrilateral face  $F$  also contains  $P_1P_2$ , and thus is also  $B_2$  (it cannot be  $B_1$ , because it has a quadrilateral face  $F$  outside the coordinate hyperplanes). Then we would have two  $B_2$ -facets with a common quadrilateral face  $F$  (see Lemma 4.7 for a contradiction).

Thus all of  $\tau_j$ 's are  $B_1$ -facets. As we have seen above, one of  $P_1$  and  $P_2$  is equal to the apices of all  $\tau_j$ 's, and the other one is in the bases of all  $\tau_j$ 's. Then, among  $\tau_j$ 's, we can find at least two pairs of  $B_1$ -facets with a common apex, a common triangular face and different bases.

These two pairs surround two borders, and, since there are no  $B$ -borders by the assumptions of Theorem 5.2, they are  $B^2$ -borders. Two  $B^2$ -borders with a common apex  $A$ , intersecting in a common edge  $AC$  in the interior of  $\mathbb{R}_+^4$ , by their definition have  $C = (0, 0, 1, 1)$  and  $A = (1, 1, 0, 0)$ , which by Lemma 4.8 implies that the singularity  $f$  is Morse non-degenerate.  $\square$

### 5.3 A tubular neighborhood of the critical subfan revisited

We are now ready to prove Theorem 5.2. Recall that by this time we have

- chosen once and for all a preferred base of every  $B_1$ -facet and every  $B_2$ -facet in the Newton polyhedron  $\Gamma_+(f)$ ,
- depending on this choice, called some cones critical in the dual fan  $\Sigma_0$  (see Definitions 4.25 and 5.8 respectively),
- chosen once and for all a generic delimiter hyperplane  $D_\sigma$  for every 2-dimensional very critical cone  $\sigma$ , dual to a non-border (see Definition 5.10), and defined the delimiter hyperplane  $D_\sigma$  for every 1-dimensional very critical cone  $\sigma$ , dual to a  $B_2$ -facet (see Definition 4.11).

**Definition 5.14.** These two kinds of faces and their dual cones will be called *delimited*.

Recall that, similarly to Subsection 4.7, we choose once and for all an affine structure on the projectivization  $\mathbb{P}\mathbb{R}_+^n$  and refer to cones and their projectivizations interchangeably whenever it causes no confusion.

According to Corollary 4.28, the set of projectivized critical cones in the dual fan  $\Sigma_0$  is a closed polyhedral complex  $\Sigma_c$ , disjoint from the boundary of  $\mathbb{P}\mathbb{R}_+^n$  by Lemma 4.26. We shall construct a generic piecewise linear tubular neighborhood of  $\Sigma_c$ , whose boundary is transversal to the critical sets of non-critical cones and delimiters of very critical cones.

For every projectivized closed cone  $\sigma \in \Sigma_c$ , choose a convex piecewise linear (with respect to the selected affine structure) “distance function”  $d_\sigma : \mathbb{P}\mathbb{R}_+^n \rightarrow \mathbb{R}_+$ , satisfying the following properties:

- 1) It vanishes on  $\sigma$  and is strictly positive outside of it.
- 2) For every non-critical cone  $\sigma' \in \Sigma_0$ , the corner locus of  $d_\sigma$  is transversal to the projectivized critical set  $\mathbb{P}K_{\sigma'}$  in the complement to  $\sigma$ .
- 3) Moreover, for every non-critical cone  $\sigma'$  and every its delimited face  $\sigma'' \subset \sigma'$ , the corner locus of  $d_\sigma$  is transversal to the projectivized delimiter  $\mathbb{P}D_{\sigma''} \cap \sigma'$  and the critical set  $\mathbb{P}D_{\sigma''} \cap K_{\sigma'}$  in the complement to  $\sigma$ .

For this distance, we shall consider  $\varepsilon$ -neighborhoods  $B_\sigma(\varepsilon) = \{u \mid d_\sigma(u) < \varepsilon\}$  and their boundaries  $S_\sigma(\varepsilon)$ , which are all piecewise-linear sets.

Associating positive numbers  $\varepsilon_\sigma$  to all  $\sigma \in \Sigma_c$ , introduce the following sets:

- the open neighborhood  $U = \bigcup_{\sigma \in \Sigma_c} B_\sigma(\varepsilon_\sigma)$  of the critical complex  $\Sigma_c$ ;
- for every  $\sigma \in \Sigma_c$ , the set  $U_\sigma = B_\sigma(\varepsilon_\sigma) \setminus \bigcup_{\sigma' \subsetneq \sigma} B_{\sigma'}(\varepsilon_{\sigma'})$ ;
- for every delimited  $\sigma \in \Sigma_c$ , the *delimiter disk*  $DD_\sigma = U_\sigma \cap \mathbb{P}D_\sigma$ .

**Lemma 5.15.** *One can choose the numbers  $\varepsilon_\sigma$  so that*

1. *For every projectivized non-critical cone  $\sigma' \in \mathbb{P}\Sigma_0$ , no vertex of the boundary of  $U_\sigma \cap \sigma'$  and  $DD_\sigma \cap \sigma'$  is contained in the critical set  $\mathbb{P}K_{\sigma'} \cap \text{rel.int}\sigma'$ ;*
2.  *$U_\sigma$  is contained in the star of  $\sigma$  and is border-convex (Definition 5.3) if  $\sigma$  is dual to a border;*
3.  *$U_\sigma \cap U_\delta = \emptyset$  unless  $\sigma = \delta$ , and  $\bar{U}_\sigma \cap \bar{U}_\delta = \emptyset$  unless  $\sigma \subset \delta$  or vice versa;*
4.  *$DD_\sigma$  divides  $U_\sigma$  into two connected components.*

*Proof.* All of these properties are satisfied if the tuple  $(\varepsilon_\sigma)$  is chosen generically (i.e. avoiding finitely many hyperplanes in the space of all such tuples) and rapidly decreasing (i.e.  $\varepsilon_\sigma \ll \varepsilon_\delta \ll 1$  for all  $\delta \subset \sigma$ ).  $\square$

**Definition 5.16.** The *vertex function*  $v : U \setminus (\text{delimiter disks}) \rightarrow (\text{vertices of } \Gamma_+(f))$  is defined on every  $U_\sigma$  as follows:

1. if a projectivized cone  $\sigma \in \Sigma_c$  is dual to a non-delimited face  $F \subset \Gamma_+(f)$ , then all facets containing  $F$  are  $B$ -facets, and their apices on the side of  $F$  are all equal to the same vertex  $P$ . Then we define  $v(\cdot)$  on  $F^\circ$  as  $P$ .
2. if  $\sigma \in \Sigma_c$  is dual to a delimited triangle  $F$ , separating two  $B$ -facets  $\tau_1$  and  $\tau_2$ , then we define  $v(\cdot)$  on each of the two components of  $U_\sigma \setminus DD_\sigma$  as the apex of the corresponding facet  $\tau_i$  on the side of this component.

3. if  $\sigma \in \Sigma_c$  is dual to a  $B_2$ -facet  $\tau$ , then we define  $v(\cdot)$  on each of the two components of  $U_\sigma \setminus DD_\sigma$  as the apex of  $\tau$  on the side of this component.

Note that the vertex function is locally constant on its domain. We now use this observation to consistently substitute every piece of the neighborhood  $U$  by an appropriate sprout. Recall that we refer to cones and their projectivizations interchangeably, and, in particular,  $S_{W,\tau}$  for the projectivization  $W$  of a cone  $C$  is another notation for the sprout  $S_{C,\tau}$ .

– If  $\sigma \in \Sigma_c$  is neither delimited nor dual to a border, then the vertex function  $v$  equals a constant  $P$  on  $\sigma$ , so we define  $V_{\sigma,0}$  as the sprout  $S_{U_\sigma,\tau}$  (see Definition 4.16) for any  $B$ -facet  $\tau$  containing the dual face of  $\sigma$ . Note that neither the sprout  $V_{\sigma,0} := S_{U_\sigma,\tau}$  nor its root  $R_{\sigma,0} := R_{U_\sigma,\tau} = P$  depend on the choice of  $\tau$ .

– If  $\sigma \in \Sigma_c$  is dual to a border, then the vertex function  $v$  equals a constant  $P$  on  $\sigma$ , and we define  $V_{\sigma,0}$  as the border sprout  $S_{U_\sigma,\sigma}$  (see Definition 5.4) and the root  $R_{\sigma,0}$  as  $R_{U_\sigma,\tau} = P$ .

– If  $\sigma \in \Sigma_c$  is dual to a very critical triangle, separating two  $B$ -facets  $\tau_{\pm 1}$  with vertices  $P_{\pm 1}$  on the side of  $\sigma$ , then we define  $V_{\sigma,\pm 1}$  as the sprout  $S_{\{v=P_{\pm 1}\} \cap U_\sigma, \tau_{\pm 1}}$  and the root  $R_{\sigma,\pm 1}$  as  $R_{U_\sigma, \tau_{\pm 1}} = P_{\pm 1}$ .

– If  $\sigma \in \Sigma_c$  is dual to a  $B_2$ -facet  $\tau$  with the apices  $P_{\pm 1}$ , then we define  $V_{\sigma,\pm 1}$  as the sprout  $S_{\{v=P_{\pm 1}\} \cap U_\sigma, \tau}$  and the root  $R_{\sigma,\pm 1}$  as  $R_{\{v=P_{\pm 1}\} \cap U_\sigma, \tau} = P_{\pm 1}$ . Also, choosing a ray  $r$  in the dual cone  $\sigma'$  to the quadrilateral non- $V$ -face of  $\tau$  outside of  $K_{\sigma'}$ , we define  $V_{\sigma,0}$  as the delimiter sprout  $S_{r,\tau}$  (see Lemma 4.20), leaving  $R_{\sigma,0}$  undefined.

Define the *sprouting*  $S_P$  of a vertex  $P$  as the union of all the sprouts  $V_{\sigma,*}$  ( $* = 0, \pm 1$ ) such that  $R_{\sigma,i} = P$ .

We have introduced the same system of notation as in Section 4.7, giving it meaning in a more general setting (most notably, admitting  $B^2$ -borders). With this wider meaning of notation at hand, the proof of Theorem 5.2 almost literally repeats the one for Theorem 4.3 (we repeat it for the convenience of the reader).

**Lemma 5.17.** 1. *The contribution of  $S_P$  equals*

$$\int_{S_P} \exp(-\langle u, P \rangle s - \langle u, \mathbf{1} \rangle) du$$

*modulo a function that has no pole at  $s_0$ .*

2. *No edge of the boundary of  $S_P$  is critical for  $(s_0, P)$ .*

*Proof.* The part (1) follows from Lemmas 4.19 and 5.6. To deduce (2), it is enough (by Lemma 4.21) to show that every vertex of  $(\partial U_\sigma) \cap \sigma'$  and  $(\partial DD_\sigma) \cap \sigma'$  for every projectivized cone  $\sigma' \supset \sigma$  is either not in  $\mathbb{P}K_{\sigma'}$ , or not a projectivized edge of  $\partial S_P$ . For non-critical  $\sigma'$ , this follows from Lemma 4.29(1), and every critical  $\sigma'$  is either in the projectivized interior of  $S_P$ , or disjoint from its closure, or intersects its boundary at an interior point of its facet  $DD_{\sigma'}$  (thus no vertex in  $\sigma'$  can be a projectivized edge of  $\partial S_P$ ).  $\square$

*Proof of Theorem 5.2.* By the preceding lemma and Lemma 3.6, the contribution of every  $S_P$  to the topological  $\zeta$ -function of  $f$  has no pole at  $s_0$ . By Lemma 4.20, the same

is true for  $V_{\sigma,0}$  for every  $B_2$ -facet  $\sigma$ . Since

$$V = \bigcup_{\sigma, * = 0, \pm 1} V_{\sigma,*}$$

contains all the critical cones in its interior, for every cone  $\sigma' \in \Sigma_0$  the set  $\sigma' \setminus V$  has edges of two types: either edges of  $(\partial V) \cap \sigma'$  or non-critical 1-dimensional cones of  $\Sigma_0$ . The edges of the first kind are not in the critical set of any cone by Lemma 4.31(2), and for the second kind the same holds by definition. Thus the contribution of  $\sigma' \setminus V$  to the topological zeta function has no pole at  $s_0$  as well by Lemma 3.6. We have subdivided  $\mathbb{R}_+^n$  into pieces

- $S_P$  for some vertices  $P$ ,
- $V_{\tau^c,0}$  for some  $B_2$ -facets  $\tau$ ,
- $\sigma' \setminus V$  for some non-critical cones  $\sigma'$ ,

so that none of them contributes the pole  $s_0$ . □

## 5.4 Generalizing the notion of $B$ -facets

We have seen that a  $B_1$  or  $B_2$ -facet alone never contributes its candidate pole. In Section 8 we shall prove a somewhat complementary fact:

*For  $n = 4$ , all other facets do contribute their (nearby) monodromy eigenvalues.* (\*)

(See Section 8 for a precise statement.) This dichotomy is central for the proof of the monodromy conjecture for non-degenerate singularities.

However, for  $n > 4$ , in order to keep the fact (\*) true, we should exclude from our consideration  $B$ -facets in a certain more general sense than the one assumed in Definition 3.10. What is the proper general notion of a  $B$ -facet in arbitrary dimension? A possible answer is given below by the lemma that we need in order to prove the fact (\*).

**Lemma 5.18.** *For  $n = 4$  if a compact facet  $\tau \prec \Gamma_+(f)$  is not a  $B$ -facet, then it splits into lattice simplices (with no new vertices) so that one of the simplices is not of type  $B_1$ .*

This observation suggests the following definition.

**Definition 5.19.** A facet is called a *general  $B$ -facet*, if it does not contain an  $(n - 1)$ -dimensional  $B_1$ -simplex.

The preceding discussion makes us believe that the proof of the monodromy conjecture for non-degenerate singularities in any dimension might look as follows.

**Plan 5.20.** 1. *Classify general  $B$ -facets in the sense of Definition 5.19 (that is, generalize Lemma 5.18 to an arbitrary dimension).*

2. *Generalize Theorem 4.3 to all combinatorial types of general  $B$ -facets.*

3. *Generalize the technique of Section 8 and, ultimately, the fact (\*) to all facets that are different from general  $B$ -facets.*

*Proof of Lemma 5.18.* The facet  $\tau$  contains a face  $F$  not contained in a coordinate hyperplane. Note the following facts about  $F$ :

**(1):**  $F$  has at most 4 vertices. Otherwise it contains a triangle whose sides are not in coordinate hyperplanes, and the union of this triangle and any vertex of  $\tau \setminus F$  gives a non- $B_1$ -simplex in  $\tau$ .

**(2):** If  $F$  is a quadrilateral, then some pair of its opposite edges are contained in coordinate hyperplanes, say,  $\{x_1 = 0\}$  and  $\{x_2 = 0\}$ , otherwise we get the same contradiction as in **(1)**. In this case, if a vertex of  $F$  at the hyperplane  $\{x_1 = 0\}$  has  $x_2 > 1$ , then this vertex, the two vertices of  $F \cap \{x_2 = 0\}$  and any other vertex of  $\tau \setminus F$  form a non- $B_1$ -simplex in  $\tau$ . Thus both vertices of  $F$  in the hyperplane  $\{x_1 = 0\}$  have  $x_2 = 1$  and vice versa. Thus  $\tau$  is a  $B_2$ -facet.

**(3):** If  $F$  is a triangle, then at least one of its edges is contained in a coordinate hyperplane, otherwise we get the same contradiction as in **(1)**.

**(3.1):** If the triangle  $F$  has exactly one edge in a coordinate hyperplane, say,  $\{x_1 = 0\}$ , then the coordinate  $x_1$  of the other vertex of  $F$  equals 1, otherwise  $F$  together with any vertex from  $\tau \setminus F$  form a non- $B_1$ -simplex in  $\tau$ . Also in this case, all other vertices of  $\tau$  should be in the hyperplane  $\{x_1 = 0\}$ , because otherwise such vertex together with  $F$  form a non- $B_1$ -simplex in  $\tau$ . Thus  $\tau$  is a  $B_1$ -pyramid for  $x_1$ .

**(3.2):** If the triangle  $F$  has all three edges in coordinate hyperplanes, then denote by  $V$  the set of the vertices of  $\tau$  outside  $F$ .

**(3.2.1):** If  $V$  is contained in a coordinate hyperplane  $L$ , containing one of the edges of  $F$ , then  $\tau$  has one vertex  $v$  outside  $L$ . So, depending on the distance of  $v$  to  $L$ , the facet  $\tau$  either contains a non- $B_1$ -simplex, or is itself a  $B_1$ -pyramid with the base  $L$ .

**(3.2.2):** Otherwise  $V$  contains three points in the three coordinate 2-planes containing the vertices of  $F$ , or contains two points in two of the three coordinate hyperplanes containing the edges of  $F$ , or contains two points in a coordinate 2-plane and a coordinate hyperplane transversal to it. In each of these three cases,  $\tau$  contains a non- $B$ -simplex.

**(3.3):** The only remaining case is that  $F$  is a triangle, exactly two of whose faces are in coordinate hyperplanes. Since its third edge is not in a coordinate hyperplane, then it should be an edge of another 2-dimensional face  $G$  of  $\tau$  not contained in a coordinate hyperplane.

**(3.3.1):** If  $G$  is also a triangle, exactly two of whose faces are in coordinate hyperplanes, then

the convex hull of  $F \cup G$  has a triangular face  $F'$ , whose edges are in three different coordinate hyperplanes. So this case can be done in the same way as **(3.2)** (although  $F'$  is not necessarily a face of  $\tau$ , we can still consider the set  $V$  of all the vertices of  $\tau$  outside  $F'$  and proceed as in **(3.2)**).

**(3.3.2):** Otherwise,  $G$  is of one of the types **(2)** or **(3.1)**, and thus  $\tau$  is  $B_1$  or  $B_2$  as shown in the corresponding paragraphs.  $\square$

## 6 Eigenvalues of monodromy and corners

In the first subsection, we formulate the main result of this section: certain configurations of  $V$ -faces (so called corners) always contribute a non-zero multiplicity of the expected sign to the corresponding monodromy eigenvalue. As a corollary, we prove the monodromy conjecture for a large class of Newton-non-degenerate singularities in arbitrary dimension.

The rest of the section is devoted to the proof of the main result. In particular, in the second subsection we introduce the notion of a hypermodular function, which may be of independent interest for convex geometry and analysis.

### 6.1 Motivation and results

Let  $f(x_1, \dots, x_n)$  be a polynomial on  $\mathbb{C}^n$  such that  $f(0) = 0$ . For lattice simplices  $\tau$  contained in compact facets of  $\Gamma_+(f)$  we define their  $V$ -faces and polynomials  $\zeta_\tau(t) = (1 - t^{N(\tau)})^{\text{Vol}_{\mathbb{Z}}(\tau)} \in \mathbb{C}[t]$  in the same way as for faces of  $\Gamma_+(f)$ .

Let us first observe the following fact.

**Proposition 6.1.** *Let  $\tau \prec \Gamma_+(f)$  be a compact facet such that  $\gamma = \tau \cap \{x_i = 0\}$  is its facet. Then  $\zeta_\tau/\zeta_\gamma \in \mathbb{C}(t)$  is a polynomial of  $t$ . If we assume moreover that  $\tau$  is not a  $B_1$ -pyramid for the variable  $x_i$ , then the complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C}$$

*is a root of the polynomial.*

*Proof.* By Lemma 2.7 we can easily prove that  $\zeta_\tau/\zeta_\gamma \in \mathbb{C}(t)$  is a polynomial. Let us prove the remaining assertion. If  $\tau$  is not a pyramid over  $\gamma = \tau \cap \{x_i = 0\}$ , then we have  $\text{Vol}_{\mathbb{Z}}(\tau) > \text{Vol}_{\mathbb{Z}}(\gamma)$  and the assertion is obvious. So it suffices to consider the case where  $\tau$  is a pyramid over  $\gamma = \tau \cap \{x_i = 0\}$  but its unique vertex  $P \prec \tau$  such that  $P \notin \gamma$  has height  $h \geq 2$  from the hyperplane  $\{x_i = 0\} \subset \mathbb{R}_+^n$ . In this case, we define two hyperplanes  $H_\tau$  and  $L_\tau$  in  $\mathbb{R}^n$  by

$$\begin{aligned} H_\tau &= \{v \in \mathbb{R}^n \mid \langle a(\tau), v \rangle = N(\tau)\}, \\ L_\tau &= \{v \in \mathbb{R}^n \mid \langle a(\tau), v \rangle = \langle a(\tau), \mathbf{1} \rangle = \nu(\tau)\}. \end{aligned}$$

Note that  $P \in \tau \subset H_\tau$  and  $L_\tau$  is the hyperplane passing through the point  $(1, 1, \dots, 1) \in \mathbb{R}_+^n$  and parallel to  $H_\tau$ . Namely  $H_\tau$  is the affine span  $\text{aff}(\tau) \simeq \mathbb{R}^{n-1}$  of  $\tau$ . Moreover the affine subspace  $L_\tau \cap \{x_i = 0\} \subset \mathbb{R}^n$  is parallel to the affine span  $H_\tau \cap \{x_i = 0\} \subset \mathbb{R}^n$  of  $\gamma = \tau \cap \{x_i = 0\}$ . This implies that  $\lambda = \exp(-2\pi i \nu(\tau)/N(\tau)) \in \mathbb{C}$  is a root of  $\zeta_\tau(t)$  if and only if  $L_\tau \cap \{x_i = 0\}$  is rational i.e.  $L_\tau \cap \{x_i = 0\} \cap \mathbb{Z}^n \neq \emptyset$ . On the other hand, it is easy to see that the affine subspace  $H_\tau \cap \{x_i = h - 1\} \subset \mathbb{R}^n$  is a parallel translation of  $L_\tau \cap \{x_i = 0\}$  by a lattice vector. Hence if  $L_\tau \cap \{x_i = 0\}$  is rational, then  $H_\tau \cap \{x_i = h - 1\} \cap \mathbb{Z}^n \neq \emptyset$  and the lattice height of the pyramid  $\tau$  from its base  $\gamma = \tau \cap \{x_i = 0\}$  is  $h \geq 2$  i.e.  $\text{Vol}_{\mathbb{Z}}(\tau) \geq 2\text{Vol}_{\mathbb{Z}}(\gamma)$ . It follows that the polynomial  $\zeta_\tau/\zeta_\gamma$  is divisible by the factor  $t - \lambda$ . This completes the proof.  $\square$

Motivated by this proposition, we introduce the following definitions.

**Definition 6.2.** Let  $\tau$  be a  $k$ -dimensional lattice  $V$ -simplex contained in a compact facet of  $\Gamma_+(f)$ .

1. We say that  $\tau$  has a (possibly empty) corner  $\tau_0 \prec \tau$  of codimension  $r$  if  $\dim\tau - \dim\tau_0 = k - \dim\tau_0 = r$  and any face  $\sigma$  of  $\tau$  containing it is a  $V$ -face.
2. If  $\tau$  has a (possibly empty) corner  $\tau_0 \prec \tau$  of codimension  $r$ , then we set

$$F_{\tau, \tau_0}(t) = \prod_{\sigma: \tau_0 \prec \sigma \prec \tau, \sigma \neq \emptyset} \left\{ \zeta_\sigma(t) \right\}^{(-1)^{k - \dim\sigma}} \in \mathbb{C}(t).$$

**Remark 6.3.** Every  $(n - 1)$ -dimensional lattice simplex  $\tau$  contained in a compact facet of  $\Gamma_+(f)$  has a unique corner of maximal codimension, which we will denote by  $\mathcal{C}_\tau$ . We will also write shortly  $F_\tau(t)$  for  $F_{\tau, \mathcal{C}_\tau}(t)$ .

In the next subsection we will prove the following result.

**Theorem 6.4.** 1. Let  $\tau$  be a  $k$ -dimensional lattice  $V$ -simplex contained in a compact facet of  $\Gamma_+(f)$ . Assume that for some  $r \geq 1$  it has a non-empty corner  $\tau_0$  of codimension  $r$ , then  $F_{\tau, \tau_0}(t) \in \mathbb{C}(t)$  is a polynomial of  $t$ .

2. If  $k = n - 1$  and if moreover  $\tau$  is not a  $B_1$ -pyramid, then the complex number

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C}$$

is a root of the polynomial  $F_\tau(t)$ .

We can generalize Theorem 6.4 slightly to allow also simplices with empty corners as follows. If an  $(n - 1)$ -dimensional lattice simplex  $\tau$  contained in a compact facet of  $\Gamma_+(f)$  has an empty corner  $\tau_0 = \emptyset$ , then we have the expression  $\tau = A_1 A_2 \cdots A_n$  such that for any  $1 \leq i \leq n$  the vertex  $A_i$  is on the positive part of the  $i$ -th coordinate axis of  $\mathbb{R}^n$ .

**Proposition 6.5.** Let  $\tau$  be an  $(n - 1)$ -dimensional lattice simplex contained in a compact facet of  $\Gamma_+(f)$ . Assume that it has an empty corner  $\tau_0 = \emptyset$ , then

$$F_\tau(t) \cdot (1 - t)^{(-1)^n} \in \mathbb{C}(t)$$

is a polynomial. If we assume moreover that  $\tau$  is not a  $B_1$ -simplex, then the complex number

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C}$$

is a root of the polynomial.

*Proof.* By the embedding  $\mathbb{R}^n \hookrightarrow \mathbb{R}^n \times \mathbb{R}$ ,  $v \mapsto (v, 0)$  we regard  $\tau$  as a lattice simplex in  $\mathbb{R}^n \times \mathbb{R}$  and set  $Q(0, 1) \in \mathbb{R}^n \times \mathbb{R}$ . Let  $\tau'$  be the convex hull of  $\{Q\} \cup \tau$  in  $\mathbb{R}^n \times \mathbb{R}$ . In this situation, it is easy to see that we have an equality

$$F_{\tau', Q}(t) = F_\tau(t) \cdot (1 - t)^{(-1)^n}$$

from which the first assertion immediately follows by Theorem 6.4. Since we have  $N(\tau') = N(\tau)$  and  $\nu(\tau') = \nu(\tau) + N(\tau)$ , the second assertion also follows from Theorem 6.4.  $\square$

Together with Theorem 4.3, following the strategy of Lemahieu-Van Proeyen [16] we can now confirm the monodromy conjecture for non-degenerate hypersurfaces in many cases also for  $n \geq 4$ . Let  $\tau_1, \dots, \tau_k \prec \Gamma_+(f)$  be the compact facets of  $\Gamma_+(f)$ . Then we say that the Newton polytope of  $f$  has a good pavement by lattice simplices if for any  $1 \leq i \leq k$  there exists a decomposition  $\tau_i = \cup_{j=1}^{n_i} \tau_{ij}$  of  $\tau_i$  into  $(n-1)$ -dimensional lattice simplices  $\tau_{ij}$  for which the following conditions are satisfied.

(i) For any  $1 \leq i \leq k$  and  $1 \leq j \leq n_i$  the lattice simplex  $\tau_{ij}$  has no V-face or it has a (non-empty) corner which is contained in any V-face of  $\tau_{ij}$ .

(ii) If  $\tau_{ij} \neq \tau_{i'j'}$  then they have no common V-face. Moreover any V-face of  $\Gamma_+(f)$  is decomposed into those of the lattice simplices  $\tau_{ij}$ .

(iii) For any  $1 \leq i \leq k$  the facet  $\tau_i$  is a  $B$ -facet or there exists a lattice simplex  $\tau_{ij}$  in it which is not a  $B_1$ -pyramid.

**Theorem 6.6.** *Assume that  $f$  is non-degenerate and the Newton polytope of  $f$  has a good pavement by lattice simplices. Let  $s_0 \in \mathbb{C}$  be a pole of  $Z_{\text{top},f}(s)$  which is contributed only by compact facets of  $\Gamma_+(f)$ , and those of them that are  $B$ -facets are consistent in the sense of Definition 4.2. Then the complex number  $\exp(2\pi i s_0) \in \mathbb{C}$  is an eigenvalue of the monodromy of  $f$  at the origin  $0 \in \mathbb{C}^n$ .*

*Proof.* If  $s_0 = -1$  the assertion is trivial. Otherwise, by Theorem 4.3 the pole  $s_0 \neq -1$  is contributed also by a non- $B$ -facet  $\tau_i$  of  $\Gamma_+(f)$ . Moreover by Theorem 2.4 and the conditions (i), (ii) we have

$$\zeta_{f,0}(t) = \left\{ \prod_{i=1}^k \prod_{j=1}^{n_i} F_{\tau_{ij}}(t) \right\}^{(-1)^{n-1}}.$$

Then the complex number  $\exp(2\pi i s_0) \in \mathbb{C}$  is an eigenvalue of the monodromy of  $f$  at the origin  $0 \in \mathbb{C}^n$  by Theorem 6.4 applied to the condition (iii).  $\square$

**Example 6.7.** *We consider the hypersurface*

$$H : f(x_1, x_2, x_3, x_4, x_5) = x_1^8 + x_2^6 + x_3^{10} + x_4^{12} + x_5^9 + x_1^2 x_3^3 + x_1 x_2 x_4 + x_1 x_2 x_5^3 + x_2^2 x_3 = 0$$

*which is non-degenerate at the origin in  $\mathbb{C}^5$ . We denote the vertices of  $\Gamma_+(f)$  by*

$$A = (8, 0, 0, 0, 0), B = (0, 6, 0, 0, 0), C = (0, 0, 10, 0, 0), D = (0, 0, 0, 12, 0), E = (0, 0, 0, 0, 9),$$

$$F = (2, 0, 3, 0, 0), G = (1, 1, 0, 1, 0), H = (1, 1, 0, 0, 3), I = (0, 2, 1, 0, 0).$$

*The Newton polyhedron  $\Gamma_+(f)$  has 10 compact facets, which are*

$$\tau_1 = EFGHI, \tau_2 = CEFGI, \tau_3 = CDEFG, \tau_4 = BEGHI, \tau_5 = CDEGI,$$

$$\tau_6 = BDEGI, \tau_7 = ABGHI, \tau_8 = AEF GH, \tau_9 = ADEFG, \tau_{10} = AFGHI.$$

*Then we find that*

$$\zeta_{f,0}(t) = \prod_{i=1}^{10} F_{\tau_i}(t) F_{CDE}(t) F_{AEF}(t) F_{BEI}(t) F_{CEI,CI}(t) F_{CEF,CF}(t).$$

*It follows immediately by Theorem 6.4 that the monodromy conjecture holds for  $H$  at the origin. Notice that we did not take  $F_{CEI}(t) = (\zeta_{CEI}(t)\zeta_C(t))(\zeta_{CE}(t)\zeta_{CI}(t))$  nor  $F_{CEF}(t) = (\zeta_{CEF}(t)\zeta_C(t))(\zeta_{CE}(t)\zeta_{CF}(t))$  as we have already the contributions of the V-faces  $CE$  and  $C$  in  $F_{CDE}(t) = (\zeta_{CDE}(t)\zeta_C(t)\zeta_D(t)\zeta_E(t))(\zeta_{CD}(t)\zeta_{CE}(t)\zeta_{DE}(t))$ .  $\square$*

## 6.2 Hypermodular functions

For the proof of Theorem 6.4 we shall introduce some new notions and their basic properties. Let  $S$  be a finite set and denote its power set by  $2^S$ . Namely elements of  $2^S$  are subsets  $I \subset S$  of  $S$ . Then for a function  $\phi : 2^S \rightarrow \mathbb{Z}$  we define new ones  $\phi^\downarrow, \phi^\uparrow : 2^S \rightarrow \mathbb{Z}$  by

$$\phi^\downarrow(I) = \sum_{J \subset I} \phi(J), \quad \phi^\uparrow(I) = \sum_{J \subset I} (-1)^{|I|-|J|} \phi(J).$$

We call  $\phi^\downarrow$  (resp.  $\phi^\uparrow$ ) the antiderivative (resp. derivative) of  $\phi$ . Then we can easily check that  $\phi^{\uparrow\downarrow} = \phi^{\downarrow\uparrow} = \phi$ .

**Definition 6.8.** 1. We say that the function  $\phi$  is hypermodular if  $\phi^\uparrow(I) \geq 0$  for any subset  $I \subset S$ .

2. The function  $\phi$  is called strictly hypermodular if it is hypermodular and  $\phi^\uparrow(S) > 0$ .

**Lemma 6.9.** *The product of two hypermodular functions  $\phi, \psi : 2^S \rightarrow \mathbb{Z}$  is hypermodular. Moreover it is strictly hypermodular if and only if there exist subsets  $I, J \subset S$  of  $S$  such that  $I \cup J = S$  and both  $\phi^\uparrow(I)$  and  $\psi^\uparrow(J)$  are strictly positive.*

*Proof.* For any subset  $R \subset S$  of  $S$  we have

$$\begin{aligned} (\phi\psi)^\uparrow(R) &= (\phi^{\uparrow\downarrow}\psi^{\uparrow\downarrow})^\uparrow(R) = \sum_{I \cup J \subset R} (-1)^{|R|-|I|} \phi^\uparrow(I) \psi^\uparrow(J) \\ &= \sum_{I \cup J = R} \phi^\uparrow(I) \psi^\uparrow(J). \end{aligned}$$

Then the assertion immediately follows.  $\square$

## 6.3 Reduction to the case $k = n - 1$ and $r = n - 1$

We can obviously suppose that  $k = n - 1$  as the computations are made in the coordinate hyperplane containing  $\tau$ . We now explain how to reduce the proof of Theorem 6.4 to the case  $r = n - 1$ . For simplicity assume that the corner  $\gamma \prec \tau$  of the simplex  $\tau \subset \partial\Gamma_+(f)$  is defined by  $\gamma = \tau \cap \{x_1 = x_2 = \dots = x_r = 0\}$ . We set

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C}.$$

As in the proof of Proposition 6.1 we define two parallel affine hyperplanes  $H_\tau$  and  $L_\tau$  in  $\mathbb{R}^n$  by

$$H_\tau = \{x \in \mathbb{R}^n \mid \langle a(\tau), x \rangle = N(\tau)\},$$

$$L_\tau = \{x \in \mathbb{R}^n \mid \langle a(\tau), x \rangle = \langle a(\tau), \mathbf{1} \rangle = \nu(\tau)\}.$$

Let  $W = \{x_1 = x_2 = \dots = x_r = 0\} \simeq \mathbb{R}^{n-r} \subset \mathbb{R}^n$  be the linear subspace of  $\mathbb{R}^n$  spanned by  $\gamma$ . Similarly, for a face  $\sigma$  of  $\tau$  containing  $\gamma$  let  $W_\sigma \simeq \mathbb{R}^{\dim\sigma+1} \subset \mathbb{R}^n$  be the linear subspace of  $\mathbb{R}^n$  spanned by  $\sigma$ . Then  $\zeta_\sigma(\lambda) = 0$  if and only if the affine hyperplane  $L_\tau \cap W_\sigma \subset W_\sigma$  of  $W_\sigma$  is rational i.e.  $L_\tau \cap W_\sigma \cap \mathbb{Z}^n \neq \emptyset$ . Let  $\Phi_0 : W \xrightarrow{\sim} W$  be a unimodular transformation of  $W$  such that  $\Phi_0(\gamma) \subset W \cap \{x_n = k\}$  for some  $k \in \mathbb{Z}_{>0}$ . Then we can easily extend it

to a unimodular transformation  $\Phi : \mathbb{R}^n \xrightarrow{\sim} \mathbb{R}^n$  of  $\mathbb{R}^n$  which preserves  $W_\sigma$  for any  $\sigma \prec \tau$  containing  $\gamma$  and the point

$$\mathbf{1} = \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{R}^n.$$

We can choose such  $\Phi$  so that the heights of  $\tau$  and  $\Phi(\tau)$  from each coordinate hyperplane in  $\mathbb{R}^n$  containing  $W$  are the same. Indeed, for the invertible matrix  $A_0 \in \mathrm{GL}_{n-r}(\mathbb{Z})$  representing  $\Phi_0 : W \xrightarrow{\sim} W$  it suffices to define  $\Phi : \mathbb{R}^n \xrightarrow{\sim} \mathbb{R}^n$  by taking an invertible matrix  $A \in \mathrm{GL}_n(\mathbb{Z})$  of the form

$$A = \left( \begin{array}{c|c} I_r & 0 \\ \hline * & A_0 \end{array} \right) \in \mathrm{GL}_n(\mathbb{Z})$$

such that  $A\mathbf{1} = \mathbf{1}$ , where  $I_r \in \mathrm{GL}_r(\mathbb{Z})$  stands for the identity matrix of size  $r$ . By this construction of  $\Phi$ ,  $\tau$  is a  $B_1$ -pyramid if and only if  $\Phi(\tau)$  is so. Set  $\tau' = \Phi(\tau)$  and define two parallel affine hyperplanes  $H_{\tau'}$  and  $L_{\tau'}$  in  $\mathbb{R}^n$  similarly to the case of  $\tau$  so that we have  $\Phi(H_\tau) = H_{\tau'}$ . Since  $\Phi(L_\tau)$  is parallel to  $\Phi(H_\tau) = H_{\tau'}$  and passes through the point  $\Phi(\mathbf{1}) = \mathbf{1} \in \mathbb{R}^n$ , we have also  $\Phi(L_\tau) = L_{\tau'}$ . Now let us set

$$\lambda' = \exp\left(-2\pi i \frac{\nu(\tau')}{N(\tau')}\right) \in \mathbb{C}.$$

Then for any  $\sigma \prec \tau$  containing  $\gamma$  we have  $\zeta_\sigma(\lambda) = 0 \iff \zeta_{\Phi(\sigma)}(\lambda') = 0$ . Since the lattice distance  $N(\sigma) > 0$  of  $\sigma$  (resp.  $N(\Phi(\sigma)) > 0$  of  $\Phi(\sigma)$ ) from the origin  $0 \in W_\sigma \simeq \mathbb{R}^{\dim\sigma+1}$  is equal to the number of rational hyperplanes in  $W_\sigma \simeq \mathbb{R}^{\dim\sigma+1}$  parallel to  $\sigma$  (resp.  $\Phi(\sigma)$ ) between  $\mathrm{aff}(\sigma)$  (resp.  $\mathrm{aff}(\Phi(\sigma))$ ) and the origin, we have  $N(\sigma) = N(\Phi(\sigma))$  and hence  $\zeta_\sigma(t) \equiv \zeta_{\Phi(\sigma)}(t)$ . Then we obtain an equality  $F_\tau(t) = F_{\tau'}(t)$ , where we slightly generalized Definition 6.2 in an obvious way to define  $F_{\tau'}(t)$ . Hence, to prove Theorem 6.4 we may assume that the corner  $\gamma$  of  $\tau$  is contained in  $W \cap \{x_n = k\}$  for some  $k \in \mathbb{Z}_{>0}$ . Let  $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^{r+1}$ ,  $x \mapsto (x_1, \dots, x_r, x_n)$  be the projection. Then by the definition of normalized volumes, for any face  $\sigma$  of  $\tau$  containing the corner  $\gamma \subset W \cap \{x_n = k\}$  we have  $\mathrm{Vol}_{\mathbb{Z}}(\sigma) = \mathrm{Vol}_{\mathbb{Z}}(\pi(\sigma)) \cdot \mathrm{Vol}_{\mathbb{Z}}(\gamma)$  and hence  $\zeta_\sigma(t) = \{\zeta_{\pi(\sigma)}(t)\}^{\mathrm{Vol}_{\mathbb{Z}}(\gamma)}$ . We thus obtain an equality  $F_\tau(t) = \{F_{\pi(\tau)}(t)\}^{\mathrm{Vol}_{\mathbb{Z}}(\gamma)}$ . Moreover we have  $N(\tau) = N(\pi(\tau))$  and  $\nu(\tau) = \nu(\pi(\tau))$ . This implies that we have only to consider the case  $r = n - 1$ .

## 6.4 The proof of the case $r = n - 1$

We have reduced our proof to the case where  $r = n - 1$ , a vertex  $Q$  of our simplex  $\tau = QA_1A_2 \cdots A_{n-1}$  has the form  $Q = (0, 0, \dots, 0, k)$  for some  $k \in \mathbb{Z}_{>0}$  and its edges are given by

$$\overrightarrow{QA_1} = \begin{pmatrix} a_1 \\ 0 \\ \vdots \\ 0 \\ b_1 \end{pmatrix}, \quad \overrightarrow{QA_2} = \begin{pmatrix} 0 \\ a_2 \\ \vdots \\ 0 \\ b_2 \end{pmatrix}, \quad \dots, \quad \overrightarrow{QA_{n-1}} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ a_{n-1} \\ b_{n-1} \end{pmatrix},$$

where  $a_1, a_2, \dots, a_{n-1} \in \mathbb{Z}_{>0}$  and  $b_1, b_2, \dots, b_{n-1} \in \mathbb{Z}$ . We set

$$D = \prod_{i=1}^{n-1} a_i, \quad K_i = \frac{b_i}{a_i} \cdot D \quad (1 \leq i \leq n-1)$$

and  $K = \sum_{i=1}^{n-1} K_i$ . Moreover for a subset  $I \subset S = \{1, 2, \dots, n-1\}$  we denote by  $\tau_I \prec \tau$  the face of  $\tau$  whose vertices are  $Q$  and  $A_i$  ( $i \in I$ ) and set

$$D_I = \prod_{i \in I} a_i, \quad \gcd_I = \text{GCD}\left(D, K_i \ (i \in I)\right).$$

Then it is easy to see that the  $|I|$ -dimensional normalized volume  $\text{Vol}_{\mathbb{Z}}(\tau_I)$  of  $\tau_I$  is given by the formula

$$\text{Vol}_{\mathbb{Z}}(\tau_I) = \gcd_I \cdot \frac{D_I}{D}$$

and hence we have

$$N(\tau_I) = \frac{D_I}{\text{Vol}_{\mathbb{Z}}(\tau_I)} \cdot k = \frac{D}{\gcd_I} \cdot k.$$

In particular, we obtain

$$N(\tau) = \frac{D}{\gcd_S} \cdot k.$$

For a subset  $I \subset S = \{1, 2, \dots, n-1\}$  we set

$$\zeta_I(t) = \left\{1 - t^{N(\tau_I)}\right\}^{\text{Vol}_{\mathbb{Z}}(\tau_I)} \in \mathbb{C}[t]$$

so that we have the equality

$$F_{\tau}(t) = \prod_{I \subset S} \left\{ \zeta_I(t) \right\}^{(-1)^{n-1-|I|}}.$$

**Lemma 6.10.** *The complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C}$$

*is a root of the polynomial  $\zeta_I(t)$  if and only if  $\gcd_I | K$ .*

*Proof.* Since the primitive conormal vector of the  $(n-1)$ -dimensional simplex  $\tau$  is equal to

$$a(\tau) = \frac{1}{\gcd_S} \begin{pmatrix} -K_1 \\ -K_2 \\ \vdots \\ -K_{n-1} \\ D \end{pmatrix},$$

we have

$$\nu(\tau) = \frac{D - K}{\gcd_S}$$

and

$$\frac{\nu(\tau)N(\tau_I)}{N(\tau)} = \frac{D - K}{\gcd_I}. \quad (6.1)$$

Note that  $\lambda$  is a root of  $\zeta_I(t)$  if and only if  $\nu(\tau)N(\tau_I)/N(\tau)$  is an integer. Then the assertion follows immediately from (6.1) and the fact  $\gcd_I|D$ . This completes the proof.  $\square$

By this lemma the multiplicity of  $t - \lambda$  in the rational function  $F_\tau(t)$  is equal to

$$\sum_{I: \gcd_I|K} (-1)^{n-1-|I|} \gcd_I \cdot \frac{D_I}{D}.$$

Similarly we obtain the following result.

**Lemma 6.11.** *For any  $m \in \mathbb{Z}$  the complex number  $\exp(2\pi im/N(\tau)) \in \mathbb{C}$  is a root of the polynomial  $\zeta_I(t)$  if and only if  $\gcd_I|(m \cdot \gcd_S)$ .*

**Proposition 6.12.** *The function  $F_\tau(t)$  is a polynomial in  $t$ .*

*Proof.* By Lemma 6.11 it suffices to show that for any  $m \in \mathbb{Z}$  the alternating sum

$$G_m = \sum_{I: \gcd_I|(m \cdot \gcd_S)} (-1)^{n-1-|I|} \gcd_I \cdot \frac{D_I}{D}$$

is non-negative. Fix  $m \in \mathbb{Z}$  and for a prime number  $p$  denote its multiplicities in the prime decompositions of  $a_i, b_i$  and  $m$  by  $\alpha(p)_i, \beta(p)_i$  and  $\delta(p)$  respectively. We set

$$\gamma(p) = \delta(p) + \min_{1 \leq i \leq n-1} \{\beta(p)_i - \alpha(p)_i, 0\}$$

and define a function  $\phi_p : 2^S \rightarrow \mathbb{Z}$  by

$$\phi_p(I) = \begin{cases} p^{\min_{i \in I} \{\beta(p)_i - \alpha(p)_i, 0\} + \sum_{i \in I} \alpha(p)_i} & \left( \min_{i \in I} \{\beta(p)_i - \alpha(p)_i, 0\} \leq \gamma(p) \right), \\ 0 & \text{(otherwise)}. \end{cases}$$

Then it is easy to see that for the function  $\phi = \prod_{p: \text{prime}} \phi_p : 2^S \rightarrow \mathbb{Z}$  we have

$$\phi^\uparrow(S) = G_m.$$

By Lemma 6.9 we have only to prove that for any prime number  $p$  the function  $\phi_p : 2^S \rightarrow \mathbb{Z}$  is hypermodular. For this purpose, we reorder the pairs  $(a_i, b_i)$  ( $1 \leq i \leq n-1$ ) so that we have

$$\beta(p)_1 - \alpha(p)_1 \leq \beta(p)_2 - \alpha(p)_2 \leq \cdots \leq \beta(p)_{n-1} - \alpha(p)_{n-1}.$$

Fix a subset  $I = \{i_1, i_2, i_3, \dots\} \subset S = \{1, 2, \dots, n-1\}$  ( $i_1 < i_2 < i_3 < \dots$ ) of  $S$ . We will show the non-negativity of the alternating sum

$$\phi_p^\uparrow(I) = \sum_{J \subset I} (-1)^{|I|-|J|} \phi_p(J). \quad (6.2)$$

We define  $q \geq 0$  to be the maximal number such that  $\beta(p)_{i_q} - \alpha(p)_{i_q} < 0$  (resp.  $\beta(p)_{i_q} - \alpha(p)_{i_q} \leq \gamma(p)$ ) in the case  $\gamma(p) \geq 0$  (resp.  $\gamma(p) < 0$ ). First let us consider the case  $\gamma(p) \geq 0$ . Then for  $1 \leq l \leq q$  the part of the alternating sum (6.2) over the subsets  $J \subset I$  such that  $\min J = i_l$  is equal to

$$(-1)^{l-1} p^{\beta(p)_{i_l}} \prod_{j>l} (p^{\alpha(p)_{i_j}} - 1).$$

Moreover the remaining part of (6.2) is equal to

$$(-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1).$$

We thus obtain the equality

$$\begin{aligned} \phi_p^\uparrow(I) &= p^{\beta(p)_{i_1}} \prod_{j>1} (p^{\alpha(p)_{i_j}} - 1) - p^{\beta(p)_{i_2}} \prod_{j>2} (p^{\alpha(p)_{i_j}} - 1) + \dots \\ &\dots + (-1)^{q-1} p^{\beta(p)_{i_q}} \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1) + (-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1). \end{aligned}$$

Note that for any  $1 \leq j \leq q$  we have  $\beta(p)_{i_j} - \alpha(p)_{i_j} < 0$  and obtain an inequality

$$p^{\beta(p)_{i_{j-1}}} (p^{\alpha(p)_{i_j}} - 1) \geq p^{\alpha(p)_{i_j}} - 1 \geq p^{\beta(p)_{i_j}}. \quad (6.3)$$

Thus, subdividing the terms in the above expression of  $\phi_p^\uparrow(I)$  into pairs, we get the desired non-negativity  $\phi_p^\uparrow(I) \geq 0$ . Finally let us consider the case  $\gamma(p) < 0$ . In this case, we have the following expression of  $\phi_p^\uparrow(I)$ :

$$\phi_p^\uparrow(I) = p^{\beta(p)_{i_1}} \prod_{j>1} (p^{\alpha(p)_{i_j}} - 1) - p^{\beta(p)_{i_2}} \prod_{j>2} (p^{\alpha(p)_{i_j}} - 1) + \dots + (-1)^{q-1} p^{\beta(p)_{i_q}} \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1).$$

Then by using the inequality (6.3) we can prove the non-negativity  $\phi_p^\uparrow(I) \geq 0$  as in the previous case  $\gamma(p) \geq 0$ . This completes the proof.  $\square$

**Proposition 6.13.** *Assume that  $\tau$  is not a  $B_1$ -simplex. Then the complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C}$$

*is a root of the polynomial  $F_\tau(t)$ .*

*Proof.* By Lemma 6.10 it suffices to show that the alternating sum

$$G = \sum_{I: \gcd_I | K} (-1)^{n-1-|I|} \gcd_I \cdot \frac{D_I}{D}$$

is positive. For a prime number  $p$  denote its multiplicities in the prime decompositions of  $a_i, b_i$  and  $K$  by  $\alpha(p)_i, \beta(p)_i$  and  $\kappa(p)$  respectively. We set

$$\mu(p) = \kappa(p) - \sum_{i=1}^{n-1} \alpha(p)_i$$

and define a function  $\psi_p : 2^S \longrightarrow \mathbb{Z}$  by

$$\psi_p(I) = \begin{cases} p^{\min_{i \in I} \{\beta(p)_i - \alpha(p)_i, 0\} + \sum_{i \in I} \alpha(p)_i} & \left( \min_{i \in I} \{\beta(p)_i - \alpha(p)_i, 0\} \leq \mu(p) \right), \\ 0 & \text{(otherwise)}. \end{cases}$$

Then it is easy to see that for the function  $\psi = \prod_{p: \text{prime}} \psi_p : 2^S \longrightarrow \mathbb{Z}$  we have

$$\psi^\uparrow(S) = G.$$

Now let us set  $S_p = \{1 \leq i \leq n-1 \mid \alpha(p)_i = 0\}$  and  $I_p = S \setminus S_p = \{1 \leq i \leq n-1 \mid \alpha(p)_i > 0\}$ . By our assumption we have  $a_i > 1$  for any  $1 \leq i \leq n-1$  and hence  $\cup_{p: \text{prime}} I_p = S$ . By Lemma 6.9, in order to show the positivity  $\psi^\uparrow(S) > 0$  it suffices to prove that for any prime number  $p$  we have  $\psi_p^\uparrow(I_p) > 0$ . As in the proof of Proposition 6.12 we reorder the pairs  $(a_i, b_i)$  ( $1 \leq i \leq n-1$ ) so that we have

$$\beta(p)_1 - \alpha(p)_1 \leq \beta(p)_2 - \alpha(p)_2 \leq \dots \leq \beta(p)_{n-1} - \alpha(p)_{n-1}$$

and  $\alpha(p)_i \geq \alpha(p)_{i+1}$  whenever  $\beta(p)_i - \alpha(p)_i = \beta(p)_{i+1} - \alpha(p)_{i+1}$ . Moreover we set  $I_p = \{i_1, i_2, i_3, \dots\}$  ( $i_1 < i_2 < i_3 < \dots$ ). We define  $q \geq 0$  to be the maximal number such that  $\beta(p)_{i_q} - \alpha(p)_{i_q} < 0$  (resp.  $\beta(p)_{i_q} - \alpha(p)_{i_q} \leq \mu(p)$ ) in the case  $\mu(p) \geq 0$  (resp.  $\mu(p) < 0$ ). Then we have the same expressions of  $\psi_p^\uparrow(I_p) > 0$  as in the proof of Proposition 6.12. In the case  $\mu(p) \geq 0$  we have

$$\begin{aligned} \psi_p^\uparrow(I_p) &= p^{\beta(p)_{i_1}} \prod_{j>1} (p^{\alpha(p)_{i_j}} - 1) - p^{\beta(p)_{i_2}} \prod_{j>2} (p^{\alpha(p)_{i_j}} - 1) + \dots \\ &\dots + (-1)^{q-1} p^{\beta(p)_{i_q}} \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1) + (-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1). \end{aligned} \quad (6.4)$$

In the case  $\mu(p) < 0$  we have

$$\psi_p^\uparrow(I_p) = p^{\beta(p)_{i_1}} \prod_{j>1} (p^{\alpha(p)_{i_j}} - 1) - p^{\beta(p)_{i_2}} \prod_{j>2} (p^{\alpha(p)_{i_j}} - 1) + \dots + (-1)^{q-1} p^{\beta(p)_{i_q}} \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1). \quad (6.5)$$

By the definitions of  $I_p = \{i_1, i_2, i_3, \dots\} \subset S$  and  $q \geq 0$  we have  $i \in I_p$  for any  $i \leq i_q$ . Eventually we find that  $i_j = j$  for any  $j \leq q$ . First let us consider the case  $I_p = \emptyset$ . Then we have

$$\psi_p(I_p) = \begin{cases} p^0 = 1 & (\mu(p) \geq 0), \\ 0 & (\mu(p) < 0). \end{cases}$$

In the case  $\mu(p) \geq 0$  we thus obtain the positivity  $\psi_p^\uparrow(I_p) > 0$ . But in the case  $\mu(p) < 0$  the condition  $I_p = \emptyset$  implies  $q = 0$  and such a case cannot occur by the following lemma.

**Lemma 6.14.** *The case  $q = 0$  and  $\mu(p) < 0$  cannot occur.*

*Proof.* Assume that  $q = 0$  and  $\mu(p) < 0$ . By the definition of  $\mu(p)$  we have

$$\text{mult}_p(K) = \text{mult}_p(D \cdot p^{\mu(p)}). \quad (6.6)$$

On the other hand, it follows from the condition  $\mu(p) < 0$  and  $q = 0$  that

$$\mu(p) < \beta(p)_{i_1} - \alpha(p)_{i_1} \leq \beta(p)_{i_2} - \alpha(p)_{i_2} \leq \cdots \cdots \cdots$$

Namely for any  $i \in I_p$  we have  $\mu(p) < \beta(p)_i - \alpha(p)_i$  and hence

$$\text{mult}_p(K_i) > \text{mult}_p(D \cdot p^{\mu(p)}).$$

Moreover for any  $i \in S_p = S \setminus I_p$  we have

$$\text{mult}_p(K_i) \geq \text{mult}_p(D) > \text{mult}_p(D \cdot p^{\mu(p)}),$$

where we used the condition  $\mu(p) < 0$  in the second inequality. We thus obtain the inequality

$$\text{mult}_p(K) = \text{mult}_p\left(\sum_{i \in S} K_i\right) > \text{mult}_p(D \cdot p^{\mu(p)})$$

which contradicts (6.6). □

By this lemma, it remains for us to treat the case  $I_p \neq \emptyset$ . From now on, we assume that  $I_p \neq \emptyset$ . Note that the inequality (6.3) becomes an equality only in the case  $p = 2$ ,  $\beta(p)_{i_{j-1}} = \beta(p)_{i_j} = 0$  and  $\alpha(p)_{i_j} = 1$ . By Lemma 6.14 this means that the sums (6.4) and (6.5) may be zero only in the following two cases:

**Case 1:**  $p = 2$ ,  $\mu(p) \geq 0$ ,  $q = 2m + 1$  for  $m \geq 0$  and  $(\alpha(p)_1, \beta(p)_1) = (a, 0)$  for  $a > 0$ ,  $(\alpha(p)_2, \beta(p)_2) = \cdots \cdots \cdots = (\alpha(p)_q, \beta(p)_q) = (1, 0)$ .

**Case 2:**  $p = 2$ ,  $\mu(p) < 0$ ,  $q = 2m$  for  $m \geq 1$  and  $(\alpha(p)_1, \beta(p)_1) = (a, 0)$  for  $a > 0$ ,  $(\alpha(p)_2, \beta(p)_2) = \cdots \cdots \cdots = (\alpha(p)_q, \beta(p)_q) = (1, 0)$ .

Indeed, in the case  $\mu(p) \geq 0$  and  $q = 2m$  for  $m \geq 0$ , if  $q < |I_p|$  the last term  $(-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1)$  of the alternating sum (6.4) is positive. Even if  $q = |I_p|$  we still have the positivity

$$(-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1) = \psi_p(\emptyset) = 1 > 0.$$

Let us show that none of the above two cases can occur.

**Case 1:** Set  $\alpha(p) = \sum_{i \in S} \alpha(p)_i$ . Then  $2^{\alpha(p)} | D$  and for any  $i \in S_2$  we have  $2^{\alpha(p)} | K_i$ . We thus obtain the equality

$$\begin{aligned} K &\equiv 2^{\alpha(p)-a} \cdot \text{odd} + (q-1) \cdot 2^{\alpha(p)-1} \cdot \text{odd} + \sum_{j>q} 2^{\alpha(p)+\beta(p)_{i_j}-\alpha(p)_{i_j}} \\ &\equiv 2^{\alpha(p)-a} \cdot \text{odd} + (q-1) \cdot 2^{\alpha(p)-1} \cdot \text{odd} \equiv 2^{\alpha(p)-a} \cdot \text{odd} \end{aligned}$$

of mod  $2^{\alpha(p)}$ , where we used also the fact that  $\beta(p)_{i_j} - \alpha(p)_{i_j} \geq 0$  for any  $j > q$ . We conclude that  $2^{\alpha(p)}$  does not divide  $K$ , which contradicts our assumption  $\mu(p) \geq 0$ .

**Case 2:** By the condition  $q \geq 2$  we have  $-1 = \beta(p)_{i_2} - \alpha(p)_{i_2} \leq \mu(p)$ . Then by  $\mu(p) < 0$  we obtain  $\mu(p) = -1$ . As in **Case 1**, by using the fact that  $q-1$  is odd and  $\mu(p) = -1$ , if  $a = 1$  we obtain the equality

$$K \equiv \sum_{j>q} 2^{\alpha(p)+\beta(p)_{i_j}-\alpha(p)_{i_j}} \equiv 0$$

of mod  $2^{\alpha(p)}$ . But this result  $2^{\alpha(p)}|K$  contradicts our assumption  $\mu(p) < 0$ . If  $a > 1$  we obtain the equality

$$K \equiv 2^{\alpha(p)-a} \cdot \text{odd}$$

of mod  $2^{\alpha(p)-1}$ . But it also contradicts  $\mu(p) = -1$ .

This completes the proof.  $\square$

## 7 On non-convenient Newton polyhedra

When dealing with a singularity  $(f, 0)$  with non-convenient Newton polyhedron  $\Gamma_+(f)$ , it happens already in dimension 2 and 3 that one has to search for the monodromy eigenvalue at some point of the hypersurface  $f = 0$  close to the origin.

**Definition 7.1** (cf. [11]). Let  $f : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$  be a germ of a holomorphic function. For all sufficiently small  $x_0 \in \mathbb{C}^n$ , the *nearby singularity* germ

$$f_{x_0} : (\mathbb{C}^n, x_0) \rightarrow (\mathbb{C}, 0), \quad f_{x_0}(x) = f(x_0 + x),$$

is well defined. We shall refer to the roots and poles of the monodromy  $\zeta$ -function of the latter germ as *nearby monodromy eigenvalues* of  $f$ .

### 7.1 Nearby singularities at coordinate lines

At almost all points on the hypersurface that are contained in a coordinate axis, the compact part of the Newton polyhedron coincides with the compact part of the projection of  $\Gamma_+(f)$  along that coordinate axis. Hence the monodromy zeta function can be computed by the same Varchenko formula in one dimension less, at least when the nearby singularity germ still is non-degenerate.

In this subsection, we prove the following generalization of [16, Lemma 9].

**Proposition 7.2.** *Assume that  $f$  is non-degenerate at the origin  $0 \in \mathbb{C}^n$ , then except for finitely many  $c \in \mathbb{C}$  the polynomial  $f(x_1, x_2, \dots, x_{n-1}, x_n + c)$  is non-degenerate at the origin  $0 \in \mathbb{C}^n$ .*

*Proof.* Let  $\tau' \prec \Gamma_+(f)$  be a face of  $\Gamma_+(f)$  which is non-compact for the variable  $x_n$  and denote its image by the projection  $\mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$  by  $\sigma \subset \mathbb{R}^{n-1}$ . Assume that  $\sigma$  is compact. Here we shall treat only the case where  $\tau'$  is a facet and hence  $\dim \sigma = n - 2$ . The other cases can be treated similarly. By a unimodular transformation of  $\mathbb{R}^n$  we regard  $\tau'$  as a lattice polytope in its affine span  $\text{aff}(\tau') \simeq \mathbb{R}^{n-1}$  and the  $\tau'$ -part  $f_{\tau'}$  of  $f$  as a Laurent polynomial on  $T' = (\mathbb{C}^*)^{n-1}$ . Then by our assumption for any compact face  $\tau$  of  $\tau'$  the hypersurface  $\{f_{\tau} = 0\} \subset T'$  is smooth and reduced. Moreover the  $\sigma$ -part of the polynomial  $f(x_1, x_2, \dots, x_{n-1}, x_n + c)$  is naturally identified with  $f_{\tau'}(x_1, \dots, x_{n-2}, c)$ . Therefore, in order to prove the assertion, it suffices to show that except for finitely many  $c \in \mathbb{C}$  the hypersurface

$$W_c = \{(x_1, \dots, x_{n-2}) \mid f_{\tau'}(x_1, \dots, x_{n-2}, c) = 0\} \subset (\mathbb{C}^*)^{n-2}$$

in  $T' \cap \{x_{n-1} = c\} \simeq (\mathbb{C}^*)^{n-2}$  is smooth and reduced. Let  $h : T' = (\mathbb{C}^*)^{n-1} \rightarrow \mathbb{C}$  be the function defined by  $h(x_1, \dots, x_{n-1}) = x_{n-1}$ . Then the set of  $c \in \mathbb{C}$  for which

$W_c \subset (\mathbb{C}^*)^{n-2}$  is not smooth or not reduced is contained in the discriminant variety of the map  $h|_{\{f_{\tau'}=0\}} : \{f_{\tau'}=0\} \rightarrow \mathbb{C}$ . For  $\varepsilon > 0$  let  $B(0; \varepsilon)^* = \{c \in \mathbb{C} \mid 0 < |c| < \varepsilon\}$  be the punctured disk centered at the origin  $0 \in \mathbb{C}$ . Then there exists a sufficiently small  $0 < \varepsilon \ll 1$  such that the hypersurface  $W_c \subset (\mathbb{C}^*)^{n-2}$  is smooth and reduced for any  $c \in B(0; \varepsilon)^*$ . Indeed, let  $\Delta = \tau' \cap \{x_n \leq l\} \subset \tau'$  ( $l \gg 0$ ) be the truncation of  $\tau'$ . Let  $\Sigma_0$  be the dual fan of the  $(n-1)$ -dimensional polytope  $\Delta$  in  $\mathbb{R}^{n-1}$  and  $\Sigma$  its smooth subdivision. We denote by  $X_\Sigma$  the toric variety associated to  $\Sigma$  (see [12] and [22] etc.). Then  $X_\Sigma$  is a smooth compactification of  $T' = (\mathbb{C}^*)^{n-1}$ . Recall that  $T' = (\mathbb{C}^*)^{n-1}$  acts naturally on  $X_\Sigma$  and the  $T'$ -orbits in it are parametrized by the cones in the smooth fan  $\Sigma$ . For a cone  $C \in \Sigma$  denote by  $T_C \simeq (\mathbb{C}^*)^{n-1-\dim C} \subset X_\Sigma$  the  $T'$ -orbit associated to  $C$ . By our assumption above, if  $C \in \Sigma$  corresponds to a compact face  $\tau$  of  $\tau'$  then the hypersurface  $W = \{f_{\tau'}=0\} \subset X_\Sigma$  intersects  $T_C \subset X_\Sigma$  transversally. We denote the meromorphic extension of  $h : T' = (\mathbb{C}^*)^{n-1} \rightarrow \mathbb{C}$  to  $X_\Sigma$  by the same letter  $h$ . Note that  $h$  has no point of indeterminacy on the whole  $X_\Sigma$  (because it is a monomial). Then as  $|c| \rightarrow 0$  the level set  $h^{-1}(c) \subset X_\Sigma$  of  $h$  tends to the union of the  $T'$ -orbits which correspond to the compact faces of  $\tau'$ . More precisely, if a cone  $C \in \Sigma$  corresponds to a compact face of  $\tau'$  then there exists an affine chart  $\mathbb{C}_y^{n-1}$  of  $X_\Sigma$  on which

$$T_C = \left\{ y = (y_1, \dots, y_{n-1}) \mid y_i = 0 \ (1 \leq i \leq \dim C), \ y_i \neq 0 \ (\dim C + 1 \leq i \leq n-1) \right\}$$

and  $h(y) = y_1^{m_1} y_2^{m_2} \cdots y_k^{m_k}$  ( $m_i \in \mathbb{Z}_{>0}$ ) for some  $k \geq 1$ . By this explicit description of  $h$  we see that for  $0 < |c| \ll 1$  the hypersurface  $h^{-1}(c)$  intersects  $W$  transversally. It follows that

$$W_c = W \cap h^{-1}(c) \cap T' \subset h^{-1}(c) \cap T' \simeq (\mathbb{C}^*)^{n-2}$$

is smooth and reduced for  $0 < |c| \ll 1$ . This completes the proof.  $\square$

## 7.2 Nearby singularities outside coordinate lines

The following example shows that from dimension four on, one might not always find the eigenvalue of monodromy corresponding to a pole of the topological zeta function at a point on a coordinate axis.

**Example 7.3.** Consider the polynomial  $f(x_1, x_2, x_3, x_4) = x_3^6 + x_2^4 x_3^5 + x_1^2 x_2^3 x_3^2 + x_2^3 x_3^2 x_4^2$ , which is non-degenerate at the origin. One finds that  $-1/3$  is a pole of  $Z_{\text{top}, f}(s)$ , contributed by the only compact facet of  $\Gamma_+(f)$ . For the zeta function of monodromy at the origin one finds

$$\zeta_{f,0}(t) = \frac{1-t^6}{1-t^{24}},$$

and so  $e^{-2i\pi/3}$  is not a zero or pole of this function. One can check that there does not exist a point  $b = (c_1, c_2, c_3, c_4)$  in  $\{f=0\}$  such that the compact part of the Newton polyhedron  $\Gamma_+(g)$  of  $g(x_1, x_2, x_3, x_4) = f(x_1 - c_1, x_2 - c_2, x_3 - c_3, x_4 - c_4)$  is a projection of  $\Gamma_+(f)$  along one or several coordinate axes and  $e^{-2i\pi/3}$  is a zero or pole of  $\zeta_{f,b}(t)$ . However,  $e^{-2i\pi/3}$  is an eigenvalue of monodromy at the points of the curve  $\mathcal{C} = \{(c, 0, 0, -ic), c \in \mathbb{C}\} \subset \{f=0\}$ .  $\square$

In this particular example, one can check that the singularity is still non-degenerate at the points of the curve  $\mathcal{C}$  where we found the eigenvalue of monodromy. However, this will not always be the case, as shown in the next example.

**Example 7.4.** We consider  $g(x_1, x_2, x_3, x_4) = x_3^6 + x_2^4 x_3^5 + x_1^2 x_2^{13} x_3^2 + x_2^{13} x_3^2 x_4^2 + 2x_1^{100} x_2^7 x_3^4 x_4^{100} + x_1^{200} x_2^{14} x_3^2 x_4^{200}$ , which up to terms of higher order equals the polynomial  $f$  considered in Example 7.3. The polynomial  $g$  is non-degenerate at the origin, and its Newton polyhedron at the origin has one compact facet, spanned by the vertices  $A(0, 13, 2, 2), B(2, 13, 2, 0), C(0, 4, 5, 0)$  and  $D(0, 0, 6, 0)$ . It contributes the candidate pole  $-1/3$  which is a pole of  $Z_{top,g}(s)$ . For the zeta function of monodromy at the origin one finds

$$\zeta_{g,0}(t) = \frac{1 - t^6}{1 - t^{24}}.$$

In Example 7.3 we found the eigenvalue of monodromy  $e^{-2\pi i/3}$  back at the points  $(c, 0, 0, \pm ic), c \in \mathbb{C}$ . In the translated local coordinates at the point  $(c, 0, 0, \pm ic)$ , the principal part of  $g$  will be

$$x_3^6 + x_2^4 x_3^5 + 2c^{200} x_2^7 x_3^4 + c^{400} x_2^{14} x_3^2 + 2cx_1 x_2^{13} x_3^2 - 2icx_2^{13} x_3^2 x_4,$$

having a degenerate edge  $x_3^6 + 2c^{200} x_2^7 x_3^4 + c^{400} x_2^{14} x_3^2$ . □

This example shows we have to quit the non-degenerate setting to prove the monodromy conjecture for non-degenerate singularities in dimension 4 and higher. It also shows that the Newton polyhedron of a singularity at an adjacent singular point may depend not only on the Newton polyhedron of the initial singularity, but also on higher order terms.

This obstacle motivated the first author to introduce the notion of tropical monodromy eigenvalues (see [11]). The main result in [11] makes it possible to find some of the nearby monodromy eigenvalues outside the coordinate axes, given only the Newton polyhedron of a non-degenerate singularity at the origin. We recall this result and restrict to dimension four from now on.

Assume that  $f(x) \in \mathbb{C}[x_1, \dots, x_4]$  is non-degenerate at the origin  $0 \in \mathbb{C}^4$ . Pick some pole of the topological zeta function  $s_0$  and denote the corresponding candidate eigenvalue of the Milnor monodromy by  $t_0 = \exp(2\pi i s_0)$ .

We suppose that there is a V-vertex  $A$  contained in an unbounded face, contributing to the eigenvalue  $t_0$ . With no loss in generality, assume that  $A$  is on the coordinate axis  $O_1$ .

Let  $I \subset \{1, \dots, 4\}$ , we will denote  $\pi_I = \pi_{\{x_i\}_{i \in I}}$  for the projection map

$$\begin{aligned} \mathbb{R}^4 &\longrightarrow \mathbb{R}^{4-|I|} \\ (x_1, x_2, x_3, x_4) &\longmapsto (\hat{x}_1, \hat{x}_2, \hat{x}_3, \hat{x}_4), \end{aligned}$$

where  $\hat{x}_i$  means that  $x_i$  is removed if and only if  $i \in I$ . When  $I = \{x_i\}$  is a singleton, we will also write  $\pi_{x_i}$ .

**Theorem 7.5.** [11, Cor. 6.15] Let  $\pi_I : \mathbb{Z}^4 \rightarrow \mathbb{Z}^2$  be the projection map. Let  $N$  be the projection of the Newton polyhedron  $\Gamma_+(f)$  under  $\pi_I$ . Let a compact edge  $E$  of the polygon  $N$  be the projection of a two-dimensional compact face  $F$  of  $\Gamma_+(f)$ . Denote the lattice distance from  $E$  to the origin by  $d$  and some root of the polynomial  $t^d - 1$  by  $t_0$ . If  $E$  is not a segment of lattice length 1 such that exactly one of its end points is a V-vertex of  $N$  contributing to  $t_0$  and such that another end point has a unique preimage in  $F$ , then  $t_0$  is a monodromy eigenvalue of the germ of  $f$  at a non-zero point of  $\{f = 0\}$ .

More specifically, when  $I = \{x_1, x_2\}$ , then there exists a curve  $C_{E,N,t_0}$  through the origin in the coordinate plane  $x_3 = x_4 = 0$  (and outside the axes  $O_1$  and  $O_2$ ) such that  $t_0$  is a monodromy eigenvalue of the germ of  $f$  at a generic point of  $C_{E,N,t_0}$ .

**Remark 7.6.** 1. If the face  $F$  contains a V-vertex on a coordinate axis out of  $I$  contributing to  $t_0$ , then the condition in Theorem 7.5 is fulfilled.

2. The nearby monodromy eigenvalues provided by this theorem are called *tropical*, because the proof of their existence in [11] is based on the calculus of so called tropical characteristic classes.

For instance, this theorem allows to find the ‘‘complicated’’ nearby monodromy eigenvalue in Example 7.4.

## 8 The monodromy conjecture for $n = 4$

Assume that  $f(x) \in \mathbb{C}[x_1, \dots, x_4]$  is non-degenerate at the origin  $0 \in \mathbb{C}^4$ . Pick some pole of the topological zeta function  $s_0$  and denote the corresponding candidate eigenvalue of the Milnor monodromy by  $t_0 = \exp(2\pi i s_0)$ . Our aim is to prove that, once Theorem 5.2 does not guarantee that  $s_0$  is fake,  $t_0$  is a root or pole of the monodromy  $\zeta$ -function of a singularity of  $f$  at some point near the origin.

In the second subsection we will see how [11] allows us to isolate many cases of the combinatorial structure of the Newton polyhedron  $\Gamma_+(f)$ , which ensure that  $t_0$  is a nearby monodromy eigenvalue outside the origin.

In the third subsection, we continue this work in the presence of a triangulation of  $\Gamma_+(f)$ . In the fourth subsection, assuming that  $\Gamma_+(f)$  does not fall within the scope of the second subsection, we subdivide the  $V$ -faces of  $\Gamma_+(f)$  into groups such that each of them ‘contributes’ a nonnegative multiplicity to  $t_0$  as an eigenvalue of the monodromy  $\zeta$ -function at the origin, in the sense of the following definition.

**Definition 8.1.** Recall that a  $V$ -face  $F$  is said to contribute to the eigenvalue  $t_0$  if  $t_0^{N(F)} = 1$ . The number  $(-1)^{\text{codim}F-1} \text{Vol}_{\mathbb{Z}}(F)$  for a contributing  $F$  and 0 for a non-contributing  $F$  is called the contribution of  $F$ . The sum of contributions of all faces from some set of faces  $S$  is called the contribution of  $S$ .

Finally, in the last subsection, we show that once one of the aforementioned groups violates the assumptions of Theorem 5.2, its contribution is strictly positive. This proves the monodromy conjecture for non-degenerate singularities of 4 variables.

The aforementioned subdivision of facets into groups is based on the fact that every  $V$ -simplex  $F$  has a unique minimal corner (possibly equal to  $F$ ), and the following notion.

**Definition 8.2.** For any triangulation of the union of compact faces of  $\Gamma_+(f)$ , the family of a  $V$ -simplex  $F$  of this triangulation is the set of all faces of  $F$ , containing its minimal corner (note that all of them are  $V$ -faces by definition of the corner). A family is said to be trivial if it consists of one element. The dimension of the maximal simplex in a family is also referred to as the dimension of the family.

Before implementing this plan, we devote the first subsection to a sandbox three-dimensional version of this story (first, in order to illustrate a significantly more complicated four-dimensional case beforehand, and, secondly, because we shall need this three-dimensional statement anyway). Although the three-dimensional result is essentially covered by [16], the logic of our reasoning is different, and this difference becomes important in higher dimensions. Indeed, the result in [11] makes us to approach the monodromy conjecture for non-isolated singularities first by searching for the monodromy eigenvalue outside the origin and then, if necessary, at the origin (having already excluded many combinatorial possibilities for the structure of the Newton polytope).

## 8.1 Two- and three-dimensional case

We intentionally formulate things in a more complicated way than we could for two or three variables, in order to keep all the wording consistent with the four-dimensional case.

**Theorem 8.3.** *For every non-degenerate  $f \in \mathbb{C}[x, y]$ , if the family of a  $V$ -simplex  $F$  contributes to the candidate eigenvalue  $t_0$ , then  $t_0$  is a nearby monodromy eigenvalue for  $f$ , unless the compact  $V$ -simplices containing  $F$  form a non-trivial family.*

*Proof.* Step 1: looking for the monodromy eigenvalue outside the origin.

If  $\Gamma_+(f)$  has a  $V$ -vertex that contributes to the eigenvalue  $t_0$  and is not contained in a compact  $V$ -edge, then  $t_0$  is obviously a nearby monodromy eigenvalue of  $f$  at a point of a coordinate axis.

Step 2: splitting  $\Gamma_+(f)$  into families otherwise.

Define the register of  $\Gamma_+(f)$  as the set of families of all  $V$ -edges. Neglecting the cases covered by Step 1, we notice that every  $V$ -simplex enters exactly one family in the register. We first suppose that there is more than one family in the register. Notice that then there are no families with an empty corner, so the contribution of every family to the multiplicity of  $t_0$  is non-negative. If  $F$  is a  $V$ -edge, then the contribution is moreover positive for the family of  $F$ . If  $F$  is a  $V$ -vertex, then again the contribution of the family in the register to which belongs  $F$  is positive, unless the compact  $V$ -simplices containing  $F$  form a non-trivial family.

Suppose now that there is only one family in the register. If  $F$  is a  $V$ -edge, then again we are done. If  $F$  is a  $V$ -vertex contained in a  $V$ -edge having no other  $V$ -vertex, then the compact  $V$ -simplices containing  $F$  form a non-trivial family. If  $F$  is a  $V$ -vertex that belongs to the family of a  $V$ -edge  $\tau$  with exactly two  $V$ -vertices, then by Theorem 6.4  $t_0$  is a nearby monodromy eigenvalue of  $f$ , unless  $\tau$  is a  $B_1$ -facet. However, then by Proposition 3.7  $s_0$  is not a pole of the topological zeta function of  $f$ .  $\square$

**Theorem 8.4.** *For every non-degenerate  $f \in \mathbb{C}[x, y, z]$  and a triangulation of the compact faces of its Newton polyhedron, if the family of an even-dimensional  $V$ -simplex  $F$  of this triangulation contributes to the candidate eigenvalue  $t_0$ , then  $t_0$  is a nearby monodromy eigenvalue for  $f$ , unless all compact  $V$ -simplices containing  $F$  are contained in the same non-trivial family.*

*Proof.* Step 1: looking for the monodromy eigenvalue outside the origin.

If  $\Gamma_+(f)$  has a  $V$ -edge  $G$ , whose family contributes to the eigenvalue  $t_0$ , and which is not contained in a compact  $V$ -triangle, then such face is contained in a non-compact

face, parallel to a coordinate axis (say,  $O_1$ ). Then  $G$  is also a  $V$ -edge of the projection of  $\Gamma_+(f)$  along  $O_1$ , satisfying the assumption of the preceding theorem (recall Proposition 7.2), so  $t_0$  is a nearby monodromy eigenvalues of  $f$  at a point of axis  $O_1$ .

Step 2: splitting  $\Gamma_+(f)$  into families otherwise.

Define the register of  $\Gamma_+(f)$  as the set of

- families of all  $V$ -triangles,
- families of all  $V$ -edges outside the aforementioned families,
- families of all  $V$ -vertices outside the aforementioned families.

Every  $V$ -simplex enters exactly one family in the register. Neglecting the cases covered by Step 1, there are no families of  $V$ -edges that are not contained in  $V$ -triangles.

By Theorem 6.4 the contribution of every even-dimensional family to the multiplicity of  $t_0$  is non-negative. If  $F$  is a  $V$ -facet, then the family of  $F$  is in the register with a positive contribution. If  $F$  is a  $V$ -vertex, then the family in the register to which belongs  $F$  has a positive contribution, unless all compact  $V$ -simplices containing  $F$  are contained in the same non-trivial family. □

## 8.2 Dimension 4 : Looking for the eigenvalue $t_0$ outside the origin : part 1.

We suppose that there is a  $V$ -vertex  $A$  contributing to the eigenvalue  $t_0$ , and we assume that  $A$  is on the coordinate axis  $O_1$ .

In this subsection we start to exploit as much as possible Proposition 7.2 and Theorem 7.5 to derive properties on the combinatorial structure of the Newton polyhedron locally around  $A$ .

**Definition 8.5.** The link  $L_A$  is the subdivision of the triangle  $T_A$  defined by

$$x_1 = 0, x_2 + x_3 + x_4 = \varepsilon, x_2 \geq 0, x_3 \geq 0, x_4 \geq 0$$

into the isomorphic images of faces of  $\Gamma_+(f)$  intersected with the hyperplane  $x_2 + x_3 + x_4 = \varepsilon$  under the projection  $\pi_{x_1}$ , for  $\varepsilon > 0$  small enough. (The link does not depend on the choice of  $\varepsilon > 0$  in the sense that the links for all  $\varepsilon > 0$  small enough are affine isomorphic to each other.)

The image of a face  $F \subset \Gamma_+(f)$  containing  $A$  in the link  $L_A$  is referred to as the link of  $F$  in  $L_A$ .

For example, the link of a vertex of a Newton polyhedron is shown in bold on Figure 8.

**Proposition 8.6.** *The union of the relative interiors of the links of bounded faces in  $L_A$  is closed and contractable.*

*Proof.* We denote  $A(a, 0, 0, 0)$ . By taking a slice of the Newton polyhedron  $\Gamma_+(f)$  with the hyperplane  $x_1 = a - \epsilon$ , it is sufficient to prove that the union of the compact faces  $\mathcal{C}$  of every Newton polyhedron  $N$  is contractable.

Let  $N_\epsilon$  be the Minkowski sum of  $N$  and a ball of radius  $\epsilon$ . The union of its compact faces  $\mathcal{C}_\epsilon$  (i.e. the set of those boundary points that do not belong to a ray of boundary




Figure 8: links of Newton polyhedra

points) is a topological disc, because the homeomorphism with the standard simplex is provided by the Gauss-Bonnet map (sending every boundary point to its unit exterior normal vector). Now the family of sets  $\mathcal{C}_\epsilon$  is a family of vanishing neighborhoods of  $\mathcal{C}$ , so the contractibility of all  $\mathcal{C}_\epsilon$  implies the contractibility of  $\mathcal{C}$ .  $\square$

Recall that a ray  $r \in \mathbb{R}^n$  is said to belong to the recession cone of a set  $S \in \mathbb{R}^n$ , if the Minkowsky sum of  $r$  and  $S$  equals  $S$ .

**Definition 8.7.** A face of  $\Gamma_+(f)$  is called an at least  $I$ -face,  $I \subset \{1, 2, 3, 4\}$ , if its recession cone contains the positive coordinate axes  $O_i$ ,  $i \in I$ , and is called an (exactly)  $I$ -face, if no larger  $I$  satisfies this condition. This terminology transfers to the corresponding pieces of the link  $L_A$ .

For example, on Figure 8, unbounded facets of the Newton polyhedron are grey, and a 1-piece of the link, corresponding to an unbounded 1-facet, is shown in bold dashes.

**Definition 8.8.** Let  $v_i$  be the vertex of the triangle  $T_A$  opposite to its edge  $x_i = 0$ . Let  $l$  be a line separating  $v_i$  from the other vertices of the pieces of  $L_A$ . Then the subdivision of the segment  $l \cap T_A$  into its intersections with the pieces of  $L_A$  is independent of the choice of  $l$  (up to a projective transformation) and is called the link of the vertex  $v_i$  in the link  $L_A$ .

**Remark 8.9.** Let  $v_i$  be the vertex of the triangle  $T_A$  opposite to its edge  $x_i = 0$ . Denote by  $N_i$  the projection  $\pi_{x_i}\Gamma_+(f)$  along  $O_i$ .

1. If the vertex  $v_i$  as a piece of the subdivision  $L_A$  corresponds to a bounded edge of  $\Gamma_+(f)$ , then no pieces of the link  $L_A$  correspond to at least  $i$ -faces. In this case the point  $\pi_{x_i}(A)$  is not a vertex of  $N_i$ .
2. Otherwise, the pieces of the link  $L_A$ , containing  $v_i$ , are exactly the pieces corresponding to at least  $i$ -faces. Then the point  $\pi_{x_i}(A)$  is a vertex  $A_i$ , whose link is (projectively) isomorphic to a subdivision of the link of  $v_i$  in  $L_A$ .

See Figure 8 for three-dimensional examples of both cases. The preceding proposition extends to  $I$ -faces as follows. We shall refer to the links of  $I$ -faces in the link  $L_A$  as  $I$ -pieces of the link  $L_A$ .

**Corollary 8.10.** *The union of the relative interiors for all exactly  $I$ -pieces of the link  $L_A$  is contractible, and the union of the relative interiors for all at most  $I$ -pieces of the link  $L_A$  is closed.*

We get the first result about the combinatorial configuration locally at the  $V$ -vertex  $A$ .

**Lemma 8.11.** *If  $A$  contributes to the monodromy eigenvalue  $t_0$ , and  $t_0$  is not a nearby eigenvalue outside the origin, then there are two possibilities:*

- *either  $A$  is contained in a unique facet outside the coordinate planes, and this facet is an  $\{i, j\}$ -facet,*
- *or  $A$  is contained in no  $I$ -faces for  $|I| > 1$ , and in at most one  $i$ -facet for every  $i \in \{2, 3, 4\}$ .*

*Proof.* We discuss only faces containing  $A$  and only pieces of the link  $L_A$ .

(0) If there is an  $\{i, j, k\}$ -facet, then  $t_0$  is a nearby monodromy eigenvalue at every point of the  $\{i, j, k\}$ -coordinate hyperplane.

(1) If there is more than one two-dimensional at least  $i$ -piece in the link, then, by Remark 8.9, the vertex  $\pi_{x_i}(A)$  of the polyhedron  $N_i$  satisfies the assumption of Theorem 8.4 (for any triangulation of compact faces of  $N_i$ ).

(2) If there is an  $\{i, j\}$ -piece in the link with at least one edge, not contained in the boundary of  $T_A$  and disjoint from its vertices, then the two-dimensional face  $F \subset \Gamma_+(f)$ , corresponding to this edge, satisfies the assumptions of Theorem 7.5 by Remark 7.6.

(3) If there is an  $\{i, j\}$ -piece in the link with no edge, not contained in the boundary of  $T_A$  and disjoint from its vertices, and it is not a unique two-dimensional piece in the link  $L_A$ , then we arrive at the situation (1) (possibly with  $j$  instead of  $i$ ).  $\square$

### 8.3 Dimension 4 : Looking for the eigenvalue $t_0$ outside the origin : part 2.

In this subsection we continue exploiting Proposition 7.2 to get further information on the combinatorial structure of the link  $L_A$ . We shall thus investigate in particular the cases when  $t_0$  is a monodromy eigenvalue of the singularity of  $f$  at some point of the  $i$ -th coordinate axis (although not at the origin). The Newton polyhedron of such singularity equals  $\mathbb{R}_+ \times N_i$  (see Remark 8.9 for notation), so we could apply the 3-dimensional Theorem 8.4 to its analysis. However, for this theorem, we need triangulations of the bounded faces of the polyhedra  $N_i$ .

The preceding Lemma 8.11 will play a crucial role. Indeed, under its assumption, every triangulation  $\mathcal{T}$  of the compact faces of the Newton polyhedron  $\Gamma_+(f)$  ‘naturally’ (see Remark 8.12) induces a triangulation of the link  $L_A$  of a  $V$ -vertex  $A$ , contributing to the eigenvalue  $t_0$ :

1. Assuming  $A = (*, 0, 0, 0)$ , take the isomorphic images of simplices of  $\mathcal{T}$  intersected with the hyperplane  $x_2 + x_3 + x_4 = \varepsilon$  under the projection  $\pi_{x_1}$  for small  $\varepsilon$ .
2. Subdivide every  $i$ -piece (Definition 8.7) of the link  $L_A$  by several segments from the vertex  $v_i$ , so that the resulting pieces together with the ones from (1) form a triangulation of  $T_A$ .

This triangulation will be denoted by  $\tilde{L}_A$ . (If the link  $L_A$  contains a unique two-dimensional piece, corresponding to an  $\{i, j\}$ -facet, we triangulate it trivially.)

**Remark 8.12.** 1. The triangulation  $\tilde{L}_A$  is natural in the sense that, in the notation of Remark 8.9, it agrees with the corresponding triangulations of the projection polyhedra  $N_i$ . More specifically, every compact face of  $N_i$  is the projection of one compact face of  $\Gamma_+(f)$ , so the triangulation  $\mathcal{T}$  of  $\Gamma_+(f)$  induces a triangulation  $\mathcal{T}_i$  of the compact faces of  $N_i$ . Assume that the link  $L_A$  contains an  $i$ -piece, then the  $\mathcal{T}_i$ -triangulated link of  $A_i$  in  $N_i$  is affinely isomorphic to the triangulated link of  $v_i$  in  $\tilde{L}_A$ . This is an important refinement of Remark 8.9, as we shall see later in Lemma 8.15.

2. No triangulation of the link  $L_A$  may be natural in the above sense in the presence of  $\{i, j\}$ -pieces with edges outside the boundary of  $T_A$ . So we really have to work under the assumptions of Lemma 8.11 in this subsection. In particular, we have no natural notion of a link triangulation  $\tilde{L}_A$  associated to  $\mathcal{T}$  for a  $V$ -vertex  $A$  that does not contribute to the eigenvalue  $t_0$ .

In what follows, we refer to the simplices in the triangulation  $\mathcal{T}$  as  $V$ -simplices or just  $V$ -faces, because we shall not be interested in faces of  $\Gamma_+(f)$  in the usual sense anymore.

We will now continue to study how the combinatorics of  $\Gamma_+(f)$  assures the existence of a nearby monodromy eigenvalue  $t_0$  outside the origin, but, this time, taking into account the chosen triangulation  $\mathcal{T}$ .

We will choose once and for all a triangulation  $\mathcal{T}$  and corresponding link triangulations  $\tilde{L}_A$  according to the following lemma.

**Lemma 8.13.** *If  $s_0$  is a pole of the topological zeta function, then there exists a triangulation  $\mathcal{T}$  (with no new vertices) of the Newton polyhedron  $\Gamma_+(f)$ , such that either it has a non- $B$ -simplex (see Definition 3.10) contributing to the eigenvalue  $t_0 = \exp 2\pi i s_0$ , or a  $B$ -border, whose  $V$ -edge contributes to the eigenvalue  $t_0$ .*

*Proof.* Since  $\Gamma_+(f)$  does not satisfy the assumptions of Theorem 5.2 for  $s_0$ , either it contains a  $B$ -border contributing to  $t_0$ , or a non- $B$ -facet, contributing  $s_0$ . In the first case, the  $V$ -edge of the border contributes to the sought eigenvalue (see the proof of Lemma 8.30). In the second case, triangulate the contributing non- $B$ -facet by Lemma 5.18 (so that one of the resulting non- $B$ -simplices contributes to the sought eigenvalue) and extend this triangulation arbitrarily to the whole  $\Gamma_+(f)$ .  $\square$

The absence of the nearby monodromy eigenvalue  $t_0$  outside the origin imposes lots of restrictions on combinatorics of the Newton polyhedron  $\Gamma_+(f)$  and the triangulation  $\mathcal{T}$ . Let us use these restrictions to make some crucial conclusions about the combinatorics of the triangulated links of the  $V$ -vertices.

**Definition 8.14.** The vertex  $v_i$  of the triangle  $T_A$  opposite to the edge  $x_i = 0$  is said to be the  $i$ -th corner of the link  $L_A$ , if the link triangulation  $\tilde{L}_A$  contains a piece of the form  $v_i BC$ , where  $B$  and  $C$  are points on the two edges of  $T_A$  containing  $v_i$ . Its star is the set of four pieces  $v_i BC, v_i B, v_i C, v_i$ .

**Lemma 8.15.** *If  $A$  contributes to the eigenvalue  $t_0$  and  $t_0$  is not a nearby eigenvalue outside the origin, then for every  $i \in \{2, 3, 4\}$ , either the edge of  $\Gamma_+(f)$  corresponding to the vertex  $v_i \in T_A$  is bounded and is not a corner of a  $V$ -facet, or the link  $L_A$  has the  $i$ -th corner (which may correspond to an  $i$ -edge or to a bounded  $V$ -face of  $\Gamma_+(f)$ ). In particular, there are six alternatives for the vertex  $A$  in this situation:*

- The vertex  $A$  is a corner of a  $V$ -facet;




Figure 9: examples of links of vertices in the setting of Lemma 8.15

- The vertex  $A$  is a corner of a  $V$ -edge and is contained in a unique facet, which is an  $\{i, j\}$ -facet.
- The vertex  $A$  is contained in a  $V$ -edge that is not a corner of a  $V$ -facet, and the link  $L_A$  has two corners.
- The vertex  $A$  is contained in two  $V$ -edges that are not corners of  $V$ -facets, and the link  $L_A$  has one corner.
- The vertex  $L_A$  is contained in three  $V$ -edges that are not corners of  $V$ -facets.
- The link  $L_A$  has three corners.

*Proof.* If we exclude the first two cases from our consideration, then, by Lemma 8.11, there is at most one two-dimensional piece in the link  $L_A$  for  $i \in \{2, 3, 4\}$ . However, it still could contain more than one two-dimensional  $i$ -piece of the triangulated link  $\tilde{L}_A$ . To exclude this possibility, note that, in the latter case, by Remark 8.12(1) and using its notation, the vertex  $A_i$  of the polyhedron  $N_i$  with the triangulation  $\mathcal{T}_i$  satisfies the assumption of Theorem 8.4.  $\square$

We will need the following combinatorial observation, applicable to the conclusion of Lemma 8.15. Let  $\tilde{L}$  be a triangulation of a triangle  $T$ .

**Definition 8.16.** A triangle of  $\tilde{L}$  is said to be interior, if none of its edges is contained in the edges of  $T$ , and no one of its vertices is contained in the vertices of  $T$ . An interior triangle of  $\tilde{L}$  is said to be inscribed, if its three vertices are in the interior of the three edges of  $T$ . A vertex of  $T$  is called a corner of the triangulation  $\tilde{L}$ , if it is a vertex of only one of the triangles in the triangulation.

**Lemma 8.17.** 1. If the triangulation  $\tilde{L}$  of a triangle  $T$  has three corners, then either it has an inscribed triangle, or it has at least three interior triangles.

2. If the triangulation  $\tilde{L}$  has a corner, and no edge of the triangulation connects a vertex of  $T$  with its opposite edge, then  $T$  has an interior triangle.

Informally speaking, this means that, in the setting of Lemma 8.15, the link looks similarly to one of the six examples on Figure 9. (The pieces of the link that may correspond to  $i$ -facets for some  $i$  are shown in grey.)

*Proof.* The proof of Part 1 proceeds by induction on the number of triangles in the triangulation. If any corner triangle can be glued with its (unique) adjacent triangle into a larger triangle, then glue and apply the induction hypothesis. Otherwise each of the three corner triangles are adjacent to some interior triangles  $T_i$ . If  $T_i = T_j$ , then its vertices are in the interior of the three edges of  $T$ , and otherwise  $T_i$  are three different interior triangles.

Part 2 is proved in the same way.  $\square$

From Proposition 7.2 we can also deduce the following result.

**Lemma 8.18.** *If a  $V$ -triangle  $F \subset \Gamma_+(f)$  is not contained in a  $V$ -facet, and its family (see Definition 8.2) contributes to the eigenvalue  $t_0$ , then  $t_0$  is a nearby monodromy eigenvalue outside the origin.*

*Proof.* Under these assumptions, if  $F$  is in the  $i$ -th coordinate hyperplane, then  $\pi_{x_i}F$  is a facet of  $N_i = \pi_{x_i}\Gamma_+(f)$ , whose family contributes to  $t_0$ , so  $t_0$  is a nearby monodromy eigenvalue at a point of the  $i$ -th coordinate hyperplane by Theorem 8.4.  $\square$

## 8.4 Monodromy conjecture for non-degenerate singularities of four variables

We now prove the monodromy conjecture for non-degenerate singularities of four variables similarly to Theorem 8.4. Recall our setting.

Let  $f \in \mathbb{C}[x_1, x_2, x_3, x_4]$  be non-degenerate at the origin. Let  $s_0$  be a pole of the topological zeta function of  $f$  and set  $t_0 = \exp(2\pi i s_0)$ . Analogously to the proof of Theorem 8.4, we choose once and for all a triangulation  $\mathcal{T}$  of the Newton polyhedron  $\Gamma_+(f)$  in accordance with Lemma 8.13, and we will define the register of  $\Gamma_+(f)$  as a disjoint union of certain groups of families. It will be practical to work with what we call extended families.

**Definition 8.19.** The extended family of a  $V$ -simplex  $F$  is the set of all  $V$ -faces from the family of  $F$  and, in the case of  $\dim F = 3$ , also the  $V$ -vertex of  $F$ , if it is a maximal by inclusion  $V$ -face of  $F$  (note that, in this case,  $F$  has no other  $V$ -faces except for, maybe, the facet of  $F$  opposite to the  $V$ -vertex).

**Remark 8.20.** A  $V$ -vertex  $A$  is in the extended family of a  $V$ -tetrahedron  $\tau$  if and only if the image of  $\tau$  in the triangulated link  $\tilde{L}_A$  is an interior triangle or coincides with  $T_A$ .

**Definition 8.21.** The register  $\mathcal{R}$  of  $\Gamma_+(f)$  (depending on the chosen triangulation) is the set of the following extended families:

- extended families of all 3-dimensional  $V$ -simplices;
- extended families of all 2-dimensional  $V$ -simplices that do not enter the aforementioned extended families;
- extended families of all 1-dimensional  $V$ -simplices that do not enter the aforementioned extended families.

**Remark 8.22.** Notice that every positive-dimensional  $V$ -simplex enters exactly one extended family of the register. This is obvious for simplices of dimensions 2 and 3, and a  $V$ -edge  $E$  may enter families of two different  $V$ -triangles, but in this case, by the subsequent Lemma 8.23, both of these triangles are themselves in families of  $V$ -tetrahedra  $\tau_1$  and  $\tau_2$ , so their own families are not in the register. Thus the extended family of  $E$  is itself in the register for  $\tau_1 \neq \tau_2$ , and  $E$  is in the extended family of  $\tau_1 = \tau_2$  otherwise.

**Lemma 8.23.** *If a bounded  $(n - 3)$ -dimensional  $V$ -face  $\nu$  of a Newton polyhedron in  $\mathbb{R}^n$  is contained in two bounded  $(n - 2)$ -dimensional  $V$ -faces  $\nu_1$  and  $\nu_2$ , then each of these faces is contained in a bounded  $V$ -facet.*

*Proof.* This is obvious for  $n = 3$ , and the general case reduced to  $n = 3$  by taking the projection of the Newton polyhedron along the affine span of  $\nu$ .  $\square$

We will now prove the monodromy conjecture for non-degenerate singularities of four variables, modulo several lemmas in the next subsection regarding certain exotic families.

**Theorem 8.24.** *Let  $f \in \mathbb{C}[x_1, x_2, x_3, x_4]$  be non-degenerate at the origin. Let  $s_0 \neq -1$  be a pole of the topological zeta function of  $f$  and set  $t_0 = \exp(2\pi i s_0)$ . If  $t_0$  is not a (tropical) nearby monodromy eigenvalue outside the origin, then  $t_0$  is a root of the monodromy zeta function of  $f$  at the origin, and hence a monodromy eigenvalue.*

*Proof.* Let  $\mathcal{T}$  be a triangulation of the Newton polyhedron  $\Gamma_+(f)$ , according to Lemma 8.13, and induce from  $\mathcal{T}$  the corresponding link triangulations  $\tilde{L}_v$  for every  $V$ -vertex  $v$ , contributing to the eigenvalue  $t_0$  (see the beginning of the preceding subsection).

For every  $V$ -vertex  $v$ , denote by  $r(v)$  the number of extended families containing  $v$  in the register  $\mathcal{R}$ .

By Remark 8.22, we represent the multiplicity of the candidate root  $t_0 = \exp(2\pi i s_0)$  of the monodromy zeta function as

$$\sum_{\mathcal{F}} (\text{contribution of } \mathcal{F} \text{ to the multiplicity of } t_0) + \sum_v (r(v) - 1), \quad (*)$$

where  $\mathcal{F}$  runs over the register, and  $v$  runs over  $V$ -vertices, contributing to  $t_0$ .

We will prove that every term in every sum of  $(*)$  is non-negative, and, moreover, at least one term in  $(*)$  is strictly positive.

In our setting we have :

**1.** The contribution of every extended family  $\mathcal{F} \in \mathcal{R}$  to the multiplicity of  $t_0$  is non-negative. For 3-dimensional extended families, this follows from Theorem 6.4 and Lemma 8.28 in the next subsection. Note that there are no 2-dimensional families contributing to  $t_0$  by Lemma 8.18. As  $s_0 \neq -1$ , the contribution of every 1-dimensional family to  $t_0$  is also non-negative (see for example the proof of [16, Prop. 5]).

**2.** For every  $V$ -vertex  $A$ , contributing to the eigenvalue  $t_0$ , we have  $r(A) \geq 1$ . This is because, by Lemmas 8.15 and 8.17(1), the vertex  $A$  is contained in one of the following:

- a  $V$ -tetrahedron, for which  $A$  is a corner;
- a  $V$ -edge, whose extended family is in the register;
- a  $V$ -tetrahedron, whose image in the triangulated link  $\tilde{L}_A$  is interior, and whose extended family thus contains  $A$ .

**3.** The contribution of at least one odd-dimensional extended family  $\mathcal{F} \in \mathcal{R}$  is positive, or  $r(v) > 1$  for some  $V$ -vertex  $v$ . To see this, consider the following possible cases:

– If Lemma 8.13 provides a  $B$ -border  $AA_1A_2$  contained in two  $V$ -tetrahedra, whose candidate pole of the topological zeta function equals  $s_0$ , then we have three subcases for its  $V$ -edge  $A_1A_2$ :

- – The contribution of the extended family of the edge  $A_1A_2$  is positive.
- – Assume for every  $A_i$  the following: if it is a  $V$ -vertex, contributing to  $t_0$ , then the edge  $AA_i$  is in a coordinate plane. Under this assumption, the contribution of the family of the  $V$ -edge of this border is positive by Lemma 8.30 in the next subsection.

– – Assume that some  $A_i$  breaks the preceding assumption, then we have again two subcases:

--- The triangulated link  $\tilde{L}_{A_i}$  has exactly two corners, then Lemma 8.17(2) applies to  $\tilde{L}_{A_i}$  and provides an interior triangle. The  $V$ -vertex  $A_i$  is then contained in the extended families of the  $V$ -edge  $A_1A_2$  and the  $V$ -tetrahedron, corresponding to the interior triangle, so  $r(A_i) > 1$ .

--- The triangulated link  $\tilde{L}_{A_i}$  has at most one corner, then  $A_i$  is contained by Lemma 8.15 in the extended families of two  $V$ -edges, which are not corners, hence in the register, so  $r(A_i) > 1$ .

– If Lemma 8.13 provides a contributing  $V$ -tetrahedron, whose extended family coincides with the usual family, then the contribution of this family is positive by Theorem 6.4.

– If Lemma 8.13 provides a  $V$ -tetrahedron  $F$ , whose candidate pole of the topological zeta function is  $s_0$ , and whose extended family consists of the family of  $F$  and one additional contributing  $V$ -vertex  $A$ , then we have two subcases for the triangulated link  $\tilde{L}_A$  by Lemma 8.15:

– – One of the vertices of  $\tilde{L}_A$  corresponds to a  $V$ -edge, whose extended family contains  $A$  and is contained in the register. In this case  $r(A) \geq 2$ , because  $A$  is also in the extended family of  $F$ .

– – There are three corners in  $\tilde{L}_A$ . This case subdivides into the following subcases by Lemma 8.17(1):

– – – There are three interior triangles in  $\tilde{L}_A$ . Then  $r(A) > 2 > 1$ .

– – – There is an inscribed triangle in  $\tilde{L}_A$ , and it is not the image of  $F$ . Then  $r(A) > 1$ .

– – – There is an inscribed triangle in  $\tilde{L}_A$ , and it is the image of  $F$ . Then, Lemma 8.26 or 8.29 from the next subsection applies to  $F$ , so its extended family has a positive contribution to the multiplicity of  $t_0$ .

We conclude that the number  $t_0$  is a root of the monodromy zeta function and, in particular, a monodromy eigenvalue of the singularity  $f$  at the origin.  $\square$

## 8.5 Exotic families

In the course of the proof of the monodromy conjecture for  $n = 4$ , we encountered certain exotic families of  $V$ -faces, whose contributions to the multiplicity of the corresponding monodromy eigenvalue ought to be non-zero. Their contributions are estimated in this subsection.

For the most part (Lemmas 8.26–8.29), we will study the extended family of a facet  $\tau$  if it does not coincide with the family of  $\tau$  (i. e. consists of a  $V$ -vertex  $A \prec \tau$  and possibly its opposite triangular face whenever it is a  $V$ -face), and moreover  $\tau$  defines an inscribed triangle (Definition 8.16) in the link of  $A$ .

We will need the following observation from the integer linear algebra. Let us first fix notation. For a square matrix  $A$  of size  $n \times n$  and for  $I, J \subset \{1, \dots, n\}$  with same cardinality, we denote  $A_J^I$  for the minor of  $A$  removing the rows with index in the set  $I$  and removing the columns with index in the set  $J$ . When  $I = \{x_i\}$  is a singleton, we will also write  $A_J^{x_i}$  and idem for  $J$ .

**Lemma 8.25.** *Let  $A \in M_n(\mathbb{Z})$  be a square matrix of size  $n \times n$  with integer entries. Let  $P = {}^t(P_1, \dots, P_n)$  be a primitive vector of size  $n$  such that  $AP = 0$ . Then  $P_n$  divides the minors  $A_n^i$ , for  $1 \leq i \leq n$ .*

*Proof.* We proceed by induction on  $n$ . One easily verifies that the statement is true when  $n = 2$ . We now take a matrix  $A \in M_n(\mathbb{Z})$  of size  $n > 2$ . By elementary row operations over  $\mathbb{Z}$ , we transform the matrix  $A$  in a matrix  $B$  having first column  ${}^t(k, 0, \dots, 0)$ , with  $k \in \mathbb{Z}$ . Then it still holds that  $BP = 0$ . We set  $d = \gcd(P_2, \dots, P_n)$ . As  $P$  is a primitive vector, we deduce that  $d$  divides  $k$ . By the induction hypothesis we have that  $P_n/d$  divides the minors  $B_{\{1,n\}}^{\{1,i\}}$ , for all  $i \in \{2, \dots, n\}$ . Hence  $P_n$  divides  $kB_{\{1,n\}}^{\{1,i\}} = B_n^i$ , for all  $i \in \{2, \dots, n\}$ . As  $B_n^1 = 0$ , we get that  $P_n$  divides  $B_n^i$ , for all  $i \in \{1, \dots, n\}$  and so  $P_n$  also divides the original minors  $A_n^i$ , for all  $i \in \{1, \dots, n\}$ .  $\square$

We now start to study the contribution of the extended family of a  $V$ -tetrahedron  $\tau = ABCD$  if it does not coincide with the family of  $\tau$  (i.e. if the only proper  $V$ -faces of  $\tau$  are a vertex  $A$  and possibly its opposite triangle).

**Lemma 8.26.** *Let  $\tau = ABCD$  be a 3-dimensional lattice simplex in a compact facet of  $\Gamma_+(f)$  such that  $v = A(0, 0, 0, \alpha)$  is its only proper  $V$ -face contributing to  $t_0 = e^{-2\pi i v(\tau)/N(\tau)}$ . If  $B(\beta_1, \beta_2, \beta_3, \beta_4)$ ,  $C(\gamma_1, \gamma_2, \gamma_3, \gamma_4)$  and  $D(\delta_1, \delta_2, \delta_3, \delta_4)$  with  $\beta_2 = \gamma_3 = \delta_1 = 0$ , then  $\text{Vol}_{\mathbb{Z}}(\tau) \neq 1$ .*

*Proof.* Assuming to the contrary that  $\text{Vol}_{\mathbb{Z}}(\tau) = 1$ , the vector product of  $AB, AC, AD$  is a primitive vector  $(a, b, c, d)$ , normal to  $\tau$ :

$$\text{aff}(\tau) \leftrightarrow ax_1 + bx_2 + cx_3 + dx_4 = N, \text{ with}$$

$$\begin{aligned} a &= \gamma_2\delta_3(\alpha - \beta_4) + \delta_2\beta_3(\alpha - \gamma_4) - \beta_3\gamma_2(\alpha - \delta_4) \\ b &= \delta_3\beta_1(\alpha - \gamma_4) + \beta_3\gamma_1(\alpha - \delta_4) - \gamma_1\delta_3(\alpha - \beta_4) \\ c &= \beta_1\gamma_2(\alpha - \delta_4) + \gamma_1\delta_2(\alpha - \beta_4) - \delta_2\beta_1(\alpha - \gamma_4) \\ d &= \beta_1\gamma_2\delta_3 + \gamma_1\delta_2\beta_3 \\ N &= \alpha d. \end{aligned}$$

Since  $A$  contributes to  $t_0$ , we have  $d \mid a + b + c + d$ . Let  $k \in \mathbb{Z}$  be such that  $a + b + c + d = (-k + 1)d$  and let  $M$  be the matrix

$$\begin{pmatrix} \beta_1 & 0 & \beta_3 & \beta_4 - \alpha \\ \gamma_1 & \gamma_2 & 0 & \gamma_4 - \alpha \\ 0 & \delta_2 & \delta_3 & \delta_4 - \alpha \\ 1 & 1 & 1 & k \end{pmatrix}.$$

Then  $M^t v = {}^t 0$ . By Lemma 8.25 it follows that  $d$  divides the minors  $M_4^i$ ,  $1 \leq i \leq 4$ , which are

$$\begin{aligned} M_4^1 &= \gamma_1\delta_2 + \gamma_2\delta_3 - \gamma_1\delta_3 \\ M_4^2 &= \delta_3\beta_1 + \delta_2\beta_3 - \delta_2\beta_1 \\ M_4^3 &= \beta_1\gamma_2 + \beta_3\gamma_1 - \beta_3\gamma_2 \\ M_4^4 &= d. \end{aligned}$$

As  $A$  is a maximal (by inclusion) proper  $V$ -face of  $\tau$ , none of the numbers  $\beta_1, \beta_3, \gamma_1, \gamma_2, \delta_2, \delta_3$  equals 0, and then obviously  $M_4^1, M_4^2$ , and  $M_4^3$  are strictly less than  $d$ . As

$$\frac{M_4^1}{\gamma_1\delta_3} + \frac{M_4^2}{\delta_2\beta_1} + \frac{M_4^3}{\beta_3\gamma_2} = \left(\frac{\delta_2}{\delta_3} + \frac{\delta_3}{\delta_2}\right) + \left(\frac{\gamma_2}{\gamma_1} + \frac{\gamma_1}{\gamma_2}\right) + \left(\frac{\beta_3}{\beta_1} + \frac{\beta_1}{\beta_3}\right) - 3 > 0,$$

it follows that at least one of the minors  $M_4^1, M_4^2$ , or  $M_4^3$  is strictly positive, which contradicts the fact that  $d$  divides the minors  $M_4^i$ ,  $1 \leq i \leq 4$ .  $\square$

**Lemma 8.27.** *Let  $\tau = APQR$  with  $A(2, 0, 0, 0), P(0, 0, p_2, p_3), Q(0, q_1, 0, q_3), R(0, r_1, r_2, 0)$  be a 3-dimensional lattice simplex in a compact facet of  $\Gamma_+(f)$  with  $t_0 = e^{-2\pi i v(\tau)/N(\tau)} \neq 1$ . Assume that  $v = A$  and  $\sigma = PQR$  are the only proper contributing  $V$ -faces to  $t_0$  in  $\tau$ . Then  $(\text{Vol}_{\mathbb{Z}}(\sigma), \text{Vol}_{\mathbb{Z}}(\tau)) \neq (1, 2)$ .*

*Proof.* Suppose that  $\text{Vol}_{\mathbb{Z}}(\tau) = 2$  and  $\text{Vol}_{\mathbb{Z}}(\sigma) = 1$ . Let

$$\text{aff}(\tau) : ax_1 + bx_2 + cx_3 + dx_4 = N(\tau) = 2a$$

be the equation of  $\text{aff}(\tau)$  with  $\gcd(a, b, c, d) = 1$ .

The condition  $\text{Vol}_{\mathbb{Z}}(\tau) = 2\text{Vol}_{\mathbb{Z}}(\sigma)$  implies that  $\gcd(b, c, d) = 1$ .

Since the  $V$ -face  $v = A$  contributes to  $t_0$ , we have also  $a|(b+c+d)$ . Then by  $N(\tau) = 2a$ , we obtain  $t_0 = 1$  (which is excluded) or  $t_0 = -1$ . We study what happens when  $t_0 = -1$ . This implies that  $2a|(b+c+d)$ . As  $\text{Vol}_{\mathbb{Z}}(\tau) = 2$ , the even integers  $2a, 2b, 2c$  and  $2d$  are the  $3 \times 3$  minors of the matrix

$$\begin{pmatrix} -2 & 0 & p_2 & p_3 \\ -2 & q_1 & 0 & q_3 \\ -2 & r_1 & r_2 & 0 \end{pmatrix},$$

and hence the expressions for  $a, b, c$  and  $d$  become

$$\begin{aligned} a &= \frac{q_1 r_2 p_3 + r_1 p_2 q_3}{2}, & b &= r_2 p_3 + p_2 q_3 - q_3 r_2, \\ c &= p_3 q_1 + q_3 r_1 - r_1 p_3, & d &= q_1 r_2 + r_1 p_2 - p_2 q_1. \end{aligned}$$

As in the proof of Lemma 8.26, for the integer  $k = (b+c+d)/a$  we can show that the vector  $(-k, 1, 1, 1)$  is a rational linear combination of  $\overrightarrow{AP}, \overrightarrow{AQ}$  and  $\overrightarrow{AR}$ . As  $\text{Vol}_{\mathbb{Z}}(\tau) = 2$ , the cone generated by  $\overrightarrow{AP}, \overrightarrow{AQ}$  and  $\overrightarrow{AR}$  has multiplicity two and so in particular there should be an integer solution  $(x, y, z) \in \mathbb{Z}^3$  for the equation

$$\begin{pmatrix} 0 & q_1 & r_1 \\ p_2 & 0 & r_2 \\ p_3 & q_3 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix}. \quad (8.1)$$

We define a matrix  $M$  by

$$M = \begin{pmatrix} 0 & q_1 & r_1 \\ p_2 & 0 & r_2 \\ p_3 & q_3 & 0 \end{pmatrix}.$$

Then by Cramer's rule we find that

$$x = \frac{2(r_2 q_1 + r_1 q_3 - r_2 q_3)}{\det(M)}, \quad y = \frac{2(r_2 p_3 + r_1 p_2 - r_1 p_3)}{\det(M)},$$

$$\text{and } z = \frac{2(q_1 p_3 + p_2 q_3 - p_2 q_1)}{\det(M)}.$$

Now we study the possible signs of  $x, y$  and  $z$ . If  $p_2 \geq p_3$  and  $q_1 \geq q_3$ , then  $y > 0$  and  $x > 0$ . If  $p_2 \geq p_3$  and  $q_1 \leq q_3$ , then  $y > 0$  and  $z > 0$ . If  $p_2 \leq p_3$  and  $r_1 \leq r_2$ , then  $z > 0$  and  $y > 0$  and so on. Thus we find that at least two of the integers  $x, y$  and  $z$  are always positive. By permuting them, we may assume that  $x > 0$  and  $y > 0$ . As none of  $p_2, p_3, q_1, q_3, r_1, r_2$  is equal to 0, the equation  $p_3x + q_3y = 2$  obtained by (8.1) implies that  $p_3 = q_3 = 1$  and  $x = y = 1$ . Consequently we get

$$a = \frac{q_1r_2 + r_1p_2}{2}, \quad b = p_2, \quad c = q_1, \quad d = q_1r_2 + r_1p_2 - p_2q_1$$

and  $\det(M) = q_1r_2 + r_1p_2$ . As we supposed that  $2a|(b+c+d)$ , we have

$$(q_1r_2 + r_1p_2)|(p_2 + q_1 + q_1r_2 + r_1p_2 - p_2q_1) \iff \det(M)|(p_2 + q_1 - p_2q_1)$$

and  $z$  is an even integer. Hence, again by (8.1) and by using that  $x = y = 1$ , we find that  $p_2$  and  $q_1$  should be even. Then we have

$$\gcd(b, c, d) = \gcd(p_2, q_1, q_1r_2 + r_1p_2 - p_2q_1) \geq 2.$$

However, it contradicts  $\gcd(b, c, d) = 1$ . This completes the proof.  $\square$

Recall that, for a  $V$ -face  $\tau$ , we define

$$\zeta_\tau(t) = \left(1 - t^{N(\tau)}\right)^{\text{Vol}_{\mathbb{Z}}(\tau)} \in \mathbb{C}[t],$$

**Lemma 8.28.** *Let  $\tau = APQR$  be a 3-dimensional lattice simplex in a compact facet of  $\Gamma_+(f)$  such that  $v = A(\alpha, 0, 0, 0)$  and  $\sigma = PQR$  are  $V$ -faces. Then*

$$\frac{\zeta_\tau(t) \cdot (1-t)}{\zeta_v(t) \cdot \zeta_\sigma(t)}$$

*is a polynomial.*

*Proof.* If  $\text{Vol}_{\mathbb{Z}}(\tau) > \text{Vol}_{\mathbb{Z}}(\sigma)$  then the assertion is obvious. So suppose that  $\text{Vol}_{\mathbb{Z}}(\tau) = \text{Vol}_{\mathbb{Z}}(\sigma)$ . Let

$$\text{aff}(\tau) : ax_1 + bx_2 + cx_3 + dx_4 = N(\tau)$$

be the equation of  $\text{aff}(\tau)$  with  $\gcd(a, b, c, d) = 1$ . Since we have

$$N(v) = \alpha = \frac{N(\tau)}{a}, \quad N(\sigma) = \frac{N(\tau)}{\gcd(b, c, d)}$$

and  $\gcd(a, b, c, d) = 1$ , the condition

$$\text{Vol}_{\mathbb{Z}}(\tau) = \text{Vol}_{\mathbb{Z}}(\sigma) \iff \gcd(N(v), N(\sigma)) = 1$$

implies that  $N(\tau) = a \cdot \gcd(b, c, d)$ ,  $N(\sigma) = a$  and  $N(v) = \alpha = \gcd(b, c, d)$ . Hence we get

$$\frac{\zeta_\tau(t)}{\zeta_v(t) \cdot \zeta_\sigma(t)} = \frac{(1 - t^{a \cdot \alpha})^{\text{Vol}_{\mathbb{Z}}(\sigma)}}{(1 - t^\alpha) \cdot (1 - t^a)^{\text{Vol}_{\mathbb{Z}}(\sigma)}}.$$

The only common zero of  $\zeta_v(t)$  and  $\zeta_\sigma(t)$  is equal to 1.  $\square$

**Lemma 8.29.** *Let  $\tau = APQR$  with  $A(\alpha, 0, 0, 0)$ ,  $P(0, 0, p_2, p_3)$ ,  $Q(0, q_1, 0, q_3)$ ,  $R(0, r_1, r_2, 0)$  be a 3-dimensional lattice simplex in a compact facet of  $\Gamma_+(f)$  that is not of type  $B_1$  ( $\iff \alpha \geq 2$ ). Assume that  $v = A$  and  $\sigma = PQR$  are its proper contributing  $V$ -faces to  $t_0 = e^{-2\pi i\nu(\tau)/N(\tau)} \neq 1$ . Then for the polynomial*

$$F(t) = \frac{\zeta_\tau(t) \cdot (1-t)}{\zeta_v(t) \cdot \zeta_\sigma(t)} \in \mathbb{C}[t]$$

(see Lemma 8.28) we have  $F(t_0) = 0$ .

*Proof.* Since the  $V$ -face  $\sigma$  contributes to  $t_0$ , by the proof of Proposition 6.1 we have  $\text{Vol}_{\mathbb{Z}}(\tau) = \alpha \cdot \text{Vol}_{\mathbb{Z}}(\sigma)$ . Let

$$\text{aff}(\tau) : ax_1 + bx_2 + cx_3 + dx_4 = N(\tau) = a \cdot \alpha$$

be the equation of  $\text{aff}(\tau)$  with  $\gcd(a, b, c, d) = 1$ . Then by the proof of Lemma 8.28 the condition

$$\text{Vol}_{\mathbb{Z}}(\tau) = \alpha \cdot \text{Vol}_{\mathbb{Z}}(\sigma) \iff \gcd(N(v), N(\sigma)) = \alpha$$

implies that  $N(\tau) = a \cdot \alpha \cdot \gcd(b, c, d)$ ,  $N(\sigma) = a \cdot \alpha$  and  $N(v) = \alpha = \alpha \cdot \gcd(b, c, d)$ . Hence we get  $\gcd(b, c, d) = 1$ ,  $N(\tau) = N(\sigma) = a \cdot \alpha$  and

$$F(t) = \frac{(1 - t^{a \cdot \alpha})^{\alpha \cdot \text{Vol}_{\mathbb{Z}}(\sigma)} \cdot (1-t)}{(1-t^\alpha) \cdot (1-t^{a \cdot \alpha})^{\text{Vol}_{\mathbb{Z}}(\sigma)}}.$$

It follows from the assumption  $\alpha \geq 2$  that we have  $F(t_0) = 0$  unless  $\alpha = 2$  and  $\text{Vol}_{\mathbb{Z}}(\sigma) = 1$ . As shown in Lemma 8.27 this can never occur.  $\square$

It now remains to study contributions of  $B$ -borders (Definition 5.1), generalizing the proof of Theorem 15 of [16].

**Lemma 8.30.** *Assume that  $f(x) \in \mathbb{C}[x_1, \dots, x_4]$  is non-degenerate at the origin  $0 \in \mathbb{C}^4$ . Let  $\tau_1$  and  $\tau_2$  be compact  $B$ -facets in  $\Gamma_+(f)$  intersecting in a  $B$ -border  $AA_1A_2$  with a  $V$ -edge  $A_1A_2$ , and contributing the same candidate pole  $s_0 \neq 1$ .*

*Assume additionally for every  $A_i$  the following: if it is a  $V$ -vertex, contributing to the eigenvalue  $t_0 = \exp(-2\pi i s_0)$ , then the edge  $AA_i$  is in a coordinate plane.*

*Then the contribution of the family of the  $V$ -edge  $A_1A_2$  of this border to the multiplicity of the corresponding eigenvalue  $t_0 = \exp(-2\pi i s_0)$  is positive, i.e. for the polynomial*

$$F(t) = \frac{\zeta_{A_1A_2}(t) \cdot (1-t)}{\zeta_{A_1}(t) \cdot \zeta_{A_2}(t)} \in \mathbb{C}[t]$$

we have  $F(t_0) = 0$ .

*Proof.* Without loss of generality, we can assume that both  $\tau_1$  and  $\tau_2$  are compact simplicial  $B_1$ -facets: indeed, if  $\tau_i$  is a  $B_2$ -facet, then it contains a  $B_1$ -tetrahedron with the same  $B$ -border, and we can consider this  $B_1$ -tetrahedron instead of  $\tau_i$ .

We redenote  $\tau_1 = ABCD$  and  $\tau_2 = ABCE$  such that  $A = (1, 1, *, *)$ ,  $B = (0, 0, \beta_3, \beta_4)$ ,  $C = (0, 0, \gamma_3, \gamma_4)$ ,  $D = (0, \delta_2, \delta_3, \delta_4)$  and  $E = (\varepsilon_1, 0, \varepsilon_3, \varepsilon_4)$ . We compute the candidate pole  $s_0$  contributed by  $\tau_1$  and  $\tau_2$ :

$$s_0 = \frac{(\gamma_4 - \beta_4)(\beta_3 - \alpha_3 + 1) + (\beta_3 - \gamma_3)(\alpha_4 - \beta_4 + 1)}{\beta_3(\gamma_4 - \beta_4) + \beta_4(\beta_3 - \gamma_3)}.$$

As the lattice index of the V-segment  $BC$  is equal to

$$\frac{\beta_3(\gamma_4 - \beta_4) + \beta_4(\beta_3 - \gamma_3)}{\gcd(\gamma_4 - \beta_4, \beta_3 - \gamma_3)},$$

the V-face  $BC$  contributes to  $t_0$ . If neither  $B$  nor  $C$  contribute to the eigenvalue  $t_0$ , then the statement is obvious by projecting along  $BC$ .

So first suppose that  $B$  is a contributing V-vertex and  $C$  is not. Then, by the additional assumption in the statement of the lemma, we have  $A = (1, 1, 0, \alpha_4)$ , so the affine space passing through the facet  $ABCD$  has as equation:

$$\begin{aligned} x_1 [\gamma_3(\delta_4 - \beta_4) - \delta_3(\gamma_4 - \beta_4) - \delta_2\gamma_3(\alpha_4 - \beta_4)] + x_2 [\delta_3(\gamma_4 - \beta_4) - \gamma_3(\delta_4 - \beta_4)] + \\ x_3 [\delta_2(\beta_4 - \gamma_4)] + x_4(\gamma_3\delta_2) = \beta_4\gamma_3\delta_2. \end{aligned}$$

The corresponding candidate pole is then

$$\frac{\gamma_3(\alpha_4 - \beta_4) + \gamma_4 - \beta_4 - \gamma_3}{\beta_4\gamma_3}.$$

As  $B$  also contributes to  $t_0$  (say  $\beta_3 = 0$ ), one has that  $\gamma_3$  divides  $\beta_4 - \gamma_4$ . Note that  $\gamma_3 > 1$  (otherwise we have a  $B^2$ -border). This implies that  $N(BC) < \beta_4\gamma_3$  and so  $\text{Vol}_{\mathbb{Z}}(BC) > 1$ , q.e.d.

We now suppose that both  $B$  and  $C$  are contributing V-vertices to  $t_0$ . Then, by the additional assumption in the statement of the lemma, we have  $A = (1, 1, 0, 0)$ , so as before one gets  $\text{Vol}_{\mathbb{Z}}(BC) > 1$ . As  $C$  contributes to  $t_0$ , one would have a cancelation if  $\text{Vol}_{\mathbb{Z}}(BC) = 2$ . Now

$$s_0 = \frac{-\gamma_3\beta_4 - \beta_4 - \gamma_3}{\beta_4\gamma_3}$$

and so  $\beta_4$  should divide  $\gamma_3$ . We have  $\text{Vol}_{\mathbb{Z}}(BC) = \gcd(\beta_4, \gamma_3) = \gamma_3$ , because  $B$  contributes to  $t_0$ . If  $\text{Vol}_{\mathbb{Z}}(BC) = 2$ , then  $\gamma_3 = 2$  and  $\beta_4|2$ . Note that  $\beta_4 > 1$  (otherwise we have a  $B^2$ -border), so  $t_0 = 1$ .

□

## References

- [1] A'Campo, N. A. "La fonction zêta d'une monodromie", *Comment. Math. Helv.*, 50 (1975): 233-248.
- [2] Artal Bartolo, E., Cassou-Noguès, P., Luengo, I. and Melle Hernández, A., "Monodromy conjecture for some surface singularities", *Ann. Scient. Ecole Norm. Sup.*, 35 (2002): 405-460.
- [3] Artal Bartolo, E., Cassou-Noguès, P., Luengo, I. and Melle Hernández, A., "Quasi-ordinary power series and their zeta functions", *Memoirs of the A.M.S.*, 178, no. 841 (2005).
- [4] Bories, B. and Veys, W. "Igusa's  $p$ -adic local zeta function and the monodromy conjecture for non-degenerated surface singularities", *Mem. Amer. Math. Soc.*, 242, no. 1145 (2016).

- [5] Denef, J. “Report on Igusa’s local zeta function”, Séminaire Bourbaki, Vol. 1990/91. *Astérisque*, No. 201-203 (1991): 359-386.
- [6] Denef, J. “Degree of local zeta functions and monodromy”, *Compos. Math.*, No. 89 (1994): 207-216.
- [7] Denef, J. and Loeser, F. “Caractéristique d’Euler-Poincaré, fonctions zêta locales et modifications analytiques ”, *J. Amer. Math. Soc.*, 5 (1992): 705-720.
- [8] Denef, J. and Loeser, F. “Motivic Igusa zeta functions”, *J. Alg. Geom.*, 7 (1998): 505-537.
- [9] Denef, J. and Loeser, F. “Geometry on arc spaces of algebraic varieties”, *Progr. Math.*, 201 (2001): 327-348.
- [10] Dimca, A. *Sheaves in topology*, Universitext, Springer-Verlag, Berlin, 2004.
- [11] Esterov, A. “Tropical nearby monodromy eigenvalues ”, arXiv:1807.00609, submitted.
- [12] Fulton, W. *Introduction to toric varieties*, Princeton University Press, 1993.
- [13] González Villa, M. and Lemahieu, A., “The monodromy conjecture for plane meromorphic functions”, *Bull. of the London Math. Soc.*, 46, no. 3 (2014): 441-453 doi:10.1112/blms/bdt098.
- [14] Hotta, R., Takeuchi, K. and Tanisaki, T., *D-modules, perverse sheaves and representation theory*, Progress in Math., Birkhäuser, Boston, 2008.
- [15] Kouchnirenko, A. G. “Polyèdres de Newton et nombres de Milnor”, *Invent. Math.*, 32 (1976): 1-31.
- [16] Lemahieu, A. and Van Proeyen, L. “Monodromy conjecture for nondegenerate surface singularities”, *Transactions of AMS*, Vol. 363, No. 9 (2011): 4801-4829.
- [17] Lemahieu, A. and Veys, W., “Zeta functions and monodromy for surfaces that are general for a toric idealistic cluster”, *Int. Math. Res. Not.*, Vol. 2009, No. 1 (2009): 11-62.
- [18] Loeser, F. “Fonctions d’Igusa  $p$ -adiques et polynômes de Bernstein”, *Amer. J. Math.*, 110 (1988): 1-22.
- [19] Loeser, F. “Fonctions d’Igusa  $p$ -adiques, polynômes de Bernstein et polyèdres de Newton”, *J. Reine Angew. Math.*, 412 (1990): 75-96.
- [20] A. Némethi and W. Veys, “Generalized monodromy conjecture in dimension two”, *Geometry and Topology*, 16, (2012): 155-217.
- [21] Nicaise, J. “An introduction to  $p$ -adic and motivic zeta functions and the monodromy conjecture”, Algebraic and analytic aspects of zeta functions and  $L$ -functions, *MSJ Mem.*, Vol. 21, (2010): 141-166.

- [22] Oda, T. *Convex bodies and algebraic geometry. An introduction to the theory of toric varieties*, Springer-Verlag, 1988.
- [23] Oka, M. *Non-degenerate complete intersection singularity*, Hermann, Paris (1997).
- [24] Varchenko, A. N. “Zeta-function of monodromy and Newton’s diagram”, *Invent. Math.*, 37 (1976): 253-262.
- [25] Veys, W. “Determination of the poles of the topological zeta functions for curves”, *Manuscr. Math.*, 87 (1995): 435-448.

NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS  
FACULTY OF MATHEMATICS NRU HSE, 6 USACHEVA, 119048, MOSCOW, RUSSIA  
*E-mail address:* aesterov@hse.ru

UNIVERSITÉ CÔTE D’AZUR, CNRS, LABORATOIRE J.-A. DIEUDONNÉ,  
UMR CNRS 7351, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE  
*E-mail address:* ann.lemahieu@unice.fr

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TSUKUBA, 1-1-1,  
TENNODAI, TSUKUBA, IBARAKI, 305-8571, JAPAN  
*E-mail address:* takemicro@nifty.com